

**Technical Report 1716**  
January 1996

**Environmental Analysis  
of U.S. Navy Shipboard  
Solid Waste Discharges**

**Appendices A-L**



Approved for public release; distribution is unlimited.

## **LIST OF APPENDICES**

<b>Appendix A</b>	Chemical Analysis Report
<b>Appendix B</b>	Pulped Material Particle Sizing Report
<b>Appendix C</b>	Liquid Phase Organism Toxicity Testing Report
<b>Appendix D</b>	Solid Phase Benthic Organism Toxicity Screening Report
<b>Appendix E</b>	Zooplankton Interaction Report
<b>Appendix F</b>	Fish Interaction Report
<b>Appendix G</b>	Shipboard Metal Waste Discharge Corrosion Report
<b>Appendix H</b>	Wake Dispersion Modeling Results
<b>Appendix I</b>	Ambient Dispersion Modeling Results
<b>Appendix J</b>	Field Studies Plan/Acquisition Report
<b>Appendix K</b>	Sewage Emissions Data
<b>Appendix L</b>	Bibliography



## SAMPLE CROSS REFERENCE

 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: (NONE)

 Report Date: August 29, 1994  
 ATI I.D. : 408242

SI #	Client Description	Matrix	Date Collected
	P1-1	WATER	16-AUG-94
	P1-2	WATER	16-AUG-94
	P1-3	WATER	16-AUG-94
	P1-4	WATER	16-AUG-94
	P2-1	WATER	16-AUG-94
	P2-2	WATER	16-AUG-94
	P2-3	WATER	16-AUG-94
	P2-4	WATER	16-AUG-94
	P5-1	WATER	16-AUG-94
3	P5-2	WATER	16-AUG-94
1	P5-3	WATER	16-AUG-94
2	P5-4	WATER	16-AUG-94
3	P6-1	WATER	16-AUG-94
4	P6-2	WATER	16-AUG-94
5	P6-3	WATER	16-AUG-94
6	P6-4	WATER	16-AUG-94
7	P8-1	WATER	16-AUG-94
8	P8-2	WATER	16-AUG-94
9	P8-3	WATER	16-AUG-94
0	P8-4	WATER	16-AUG-94
1	P1-1	SLUDGE	16-AUG-94
2	P1-2	SLUDGE	16-AUG-94
3	P1-3	SLUDGE	16-AUG-94
4	P1-4	SLUDGE	16-AUG-94
5	P2-1	SLUDGE	16-AUG-94
6	P2-2	SLUDGE	16-AUG-94
7	P2-3	SLUDGE	16-AUG-94
8	P2-4	SLUDGE	16-AUG-94
9	P5-1	SLUDGE	16-AUG-94
0	P5-2	SLUDGE	16-AUG-94
1	P5-3	SLUDGE	16-AUG-94
2	P5-4	SLUDGE	16-AUG-94
3	P6-1	SLUDGE	16-AUG-94
4	P6-2	SLUDGE	16-AUG-94
5	P6-3	SLUDGE	16-AUG-94
6	P6-4	SLUDGE	16-AUG-94
7	P8-1	SLUDGE	16-AUG-94
8	P8-2	SLUDGE	16-AUG-94
9	P8-3	SLUDGE	16-AUG-94
0	P8-4	SLUDGE	16-AUG-94

---TOTALS---

Matrix
# Samples

SLUDGE

20



SAMPLE CROSS REFERENCE

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: (NONE)

Report Date: August 29, 1994  
ATI I.D. : 408242

---TOTALS---

<u>Matrix</u>	<u># Samples</u>
WATER	20

ATI STANDARD DISPOSAL PRACTICE

The sample(s) from this project will be disposed of in twenty-one (21) days from the date of this report. If an extended storage period is required, please contact our sample control department before the scheduled disposal date.



ANALYTICAL SCHEDULE

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: (NONE)

ATI I.D.: 408242

Analysis	Technique/Description
SA 90-3.2 (TOTAL ORGANIC CARBON)	WALKLEY-BLACK
PA 160.3 (TOTAL SOLIDS)	GRAVIMETRIC
PA 351.2 (TOTAL KJELDAHL NITROGEN)	COLORIMETRIC
PA 365.2 (TOTAL PHOSPHATE AS PHOSPHORUS)	COLORIMETRIC
PA 405.1 (BIOCHEMICAL OXYGEN DEMAND)	ELECTRODE
PA 410.2 (CHEMICAL OXYGEN DEMAND)	TITRATION



GENERAL CHEMISTRY RESULTS

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: (NONE)

ATI I.D.: 40

Sample #	Client ID	Matrix	Date Sampled	Date Received
1	P1-1	WATER	16-AUG-94	17-AUG-94
2	P1-2	WATER	16-AUG-94	17-AUG-94
3	P1-3	WATER	16-AUG-94	17-AUG-94
4	P1-4	WATER	16-AUG-94	17-AUG-94
5	P2-1	WATER	16-AUG-94	17-AUG-94

Parameter	Units	1	2	3	4	5
BIOCHEMICAL OXYGEN DEMAND	MG/L	<5.0	<5.0	<5.0	<5.0	365
CHEMICAL OXYGEN DEMAND	MG/L	344	383	295	310	196
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	0.11	<0.10	<0.10	<0.10	0.16
TOTAL KJELDAHL NITROGEN	MG/L	1.1	1.1	1.3	1.3	2.5
TOTAL SOLIDS	MG/L	7210	7360	7250	7020	9510

**GENERAL CHEMISTRY RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: (NONE)

ATI I.D.: 408242

Sample #	Client ID	Matrix	Date Sampled	Date Received
	P2-2	WATER	16-AUG-94	17-AUG-94
	P2-3	WATER	16-AUG-94	17-AUG-94
	P2-4	WATER	16-AUG-94	17-AUG-94
	P5-1	WATER	16-AUG-94	17-AUG-94
0	P5-2	WATER	16-AUG-94	17-AUG-94

Parameter	Units	6	7	8	9	10
BIOCHEMICAL OXYGEN DEMAND	MG/L	340	343	273	187	193
CHEMICAL OXYGEN DEMAND	MG/L	595	501	3500	569	1550
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	0.31	0.32	<0.10	<0.10	0.11
TOTAL KJELDAHL NITROGEN	MG/L	2.2	1.3	1.4	1.9	2.3
TOTAL SOLIDS	MG/L	9980	10100	10400	9650	10400

**GENERAL CHEMISTRY RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: (NONE)

ATI I.D.: 408

Sample #	Client ID	Matrix	Date Sampled	Date Received
11	P5-3	WATER	16-AUG-94	17-AUG-94
12	P5-4	WATER	16-AUG-94	17-AUG-94
13	P6-1	WATER	16-AUG-94	17-AUG-94
14	P6-2	WATER	16-AUG-94	17-AUG-94
15	P6-3	WATER	16-AUG-94	17-AUG-94

Parameter	Units	11	12	13	14	15
BIOCHEMICAL OXYGEN DEMAND	MG/L	125	167	63.0	222	153
CHEMICAL OXYGEN DEMAND	MG/L	221	753	1300	344	993
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	<0.10	<0.10	0.39	<0.10	0.22
TOTAL KJELDAHL NITROGEN	MG/L	2.6	2.6	2.8	2.5	2.9
TOTAL SOLIDS	MG/L	9160	8970	10400	9900	10500

**GENERAL CHEMISTRY RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONON)  
 Project Name: (NONE)

ATI I.D.: 408242

Sample #	Client ID	Matrix	Date Sampled	Date Received
6	P6-4	WATER	16-AUG-94	17-AUG-94
7	P8-1	WATER	16-AUG-94	17-AUG-94
8	P8-2	WATER	16-AUG-94	17-AUG-94
9	P8-3	WATER	16-AUG-94	17-AUG-94
10	P8-4	WATER	16-AUG-94	17-AUG-94

Parameter	Units	16	17	18	19	20
BIOCHEMICAL OXYGEN DEMAND	MG/L	88.2	135	280	311	186
CHEMICAL OXYGEN DEMAND	MG/L	151	245	202	212	151
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	<0.10	0.37	<0.10	0.52	0.29
TOTAL KJELDAHL NITROGEN	MG/L	2.2	2.3	3.1	2.6	3.9
TOTAL SOLIDS	MG/L	9920	11700	12100	11700	11500



GENERAL CHEMISTRY RESULTS

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: (NONE)

ATI I.D.: 408

Sample #	Client ID	Matrix	Date Sampled	Date Received
21	P1-1	SLUDGE	16-AUG-94	17-AUG
22	P1-2	SLUDGE	16-AUG-94	17-AUG
23	P1-3	SLUDGE	16-AUG-94	17-AUG
24	P1-4	SLUDGE	16-AUG-94	17-AUG
25	P2-1	SLUDGE	16-AUG-94	17-AUG

Parameter	Units	21	22	23	24	25
TOTAL ORGANIC CARBON (WB)	%	0.020	<0.010	0.019	<0.010	0.15

## GENERAL CHEMISTRY RESULTS

Client : NCCOSC RDT&E DIVISION  
 Object # : (NONE)  
 Object Name: (NONE)

ATI I.D.: 408242

Sample Client ID #	Matrix	Date Sampled	Date Received
P2-2	SLUDGE	16-AUG-94	17-AUG-94
P2-3	SLUDGE	16-AUG-94	17-AUG-94
P2-4	SLUDGE	16-AUG-94	17-AUG-94
P5-1	SLUDGE	16-AUG-94	17-AUG-94
P5-2	SLUDGE	16-AUG-94	17-AUG-94

Parameter	Units	26	27	28	29	30
TOTAL ORGANIC CARBON (WB)	%	0.12	0.22	0.16	0.20	0.11



GENERAL CHEMISTRY RESULTS

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: (NONE)

ATI I.D.: 40

Sample #	Client ID	Matrix	Date Sampled	Date Received
31	P5-3	SLUDGE	16-AUG-94	17-AUG-94
32	P5-4	SLUDGE	16-AUG-94	17-AUG-94
33	P6-1	SLUDGE	16-AUG-94	17-AUG-94
34	P6-2	SLUDGE	16-AUG-94	17-AUG-94
35	P6-3	SLUDGE	16-AUG-94	17-AUG-94

Parameter	Units	31	32	33	34	35
TOTAL ORGANIC CARBON (WB)	%	0.11	0.16	0.096	0.25	0.24



GENERAL CHEMISTRY RESULTS

Client : NCCOSC RDT&E DIVISION
Project # : (NONE)
Project Name: (NONE)

ATI I.D.: 408242

Table with columns: Sample Client ID #, Matrix, Date Sampled, Date Received, Parameter, Units, 36, 37, 38, 39, 40. Includes rows for samples 6-10 and TOTAL ORGANIC CARBON (WB).



Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: (NONE)

ATI I.D. : 40:

Parameters	REF I.D.	Units	Sample Result	Dup Result	RPD	Spiked Sample	Spike Conc	R
BIOCHEMICAL OXYGEN DEMAND	408242-04	MG/L	<5.0	<5.0	0	N/A	N/A	N
BIOCHEMICAL OXYGEN DEMAND	408242-13	MG/L	63.0	62.3	1	N/A	N/A	N
CHEMICAL OXYGEN DEMAND	408239-01	MG/L	13	11	17	N/A	N/A	N
CHEMICAL OXYGEN DEMAND	408240-01	MG/L	<5	<5	0	N/A	N/A	N
CHEMICAL OXYGEN DEMAND	408219-03	MG/L	22	20	10	N/A	N/A	N
TOTAL KJELDAHL NITROGEN	408282-01	MG/L	0.65	0.70	7	1.8	1.0	1
TOTAL KJELDAHL NITROGEN	408266-06	MG/L	0.84	0.88	5	1.8	1.0	9
TOTAL ORGANIC CARBON (WB)	408242-25	%	0.15	0.15	0	0.68	0.50	1
TOTAL ORGANIC CARBON (WB)	408222-06	%	1.5	1.5	0	1.9	0.50	8
TOTAL ORGANIC CARBON (WB)	408242-40	%	0.27	0.25	8	0.76	0.48	1
TOTAL PHOSPHATE AS PHOSPHORUS	408156-01	MG/L	<0.10	<0.10	0	0.42	0.40	1
TOTAL PHOSPHATE AS PHOSPHORUS	408156-04	MG/L	<0.10	<0.10	0	0.45	0.40	1
TOTAL PHOSPHATE AS PHOSPHORUS	408244-02	MG/L	<0.10	<0.10	0	0.47	0.40	1
TOTAL SOLIDS	408242-20	MG/L	11500	11700	2	N/A	N/A	N
TOTAL SOLIDS	408242-08	MG/L	10400	9760	6	N/A	N/A	N
TOTAL SOLIDS	408242-18	MG/L	12100	11000	10	N/A	N/A	N

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration  
 RPD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result



BLANK SPIKE

Client : NCCOSC RDT&E DIVISION
Project # : (NONE)
Project Name: (NONE)

ATI I.D. : 408242

Table with 7 columns: Parameters, Blank Spike ID#, Units, Blank Result, Spiked Sample, Spike Conc., % Rec. Rows include TOTAL KJELDAHL NITROGEN, TOTAL ORGANIC CARBON (WB), and TOTAL PHOSPHATE AS PHOSPHORUS.

Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration
PD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result



Analytical**Technologies, Inc.**

Corporate Offices 5550 Morehouse Drive San Diego, CA 92121 (619) 458-9141

ATI I.D.: 501244

February 13, 1995

NCCOSC RDT&E DIVISION  
53475 STROTHER ROAD RM 267A  
SAN DIEGO, CA 92152

Project Name: SSWD  
Project # : (NONE)


Attention: STACEY CURTIS

Analytical Technologies, Inc. has received the following sample(s):

<u>Date Received</u>	<u>Quantity</u>	<u>Matrix</u>
January 27, 1995	30	WATER

The sample(s) were analyzed with EPA methodology or equivalent methods as specified in the enclosed analytical schedule. The symbol for "less than" indicates a value below the reportable detection limit. If any flags appear next to the analytical data in this report, please see the attached list of flag definitions.

The results of these analyses and the quality control data are enclosed. Please note that the Sample Condition Upon Receipt Checklist is included at the end of this report.

  
JULIO PAREDES  
PROJECT MANAGER

  
ALAN J. KLEINSCHMIDT  
LABORATORY MANAGER

**SAMPLE CROSS REFERENCE**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: SSWD

Report Date: February 13, 1995  
 ATI I.D. : 501244

ATI #	Client Description	Matrix	Date Collected
1	1	WATER	27-JAN-95
2	2	WATER	27-JAN-95
3	3	WATER	27-JAN-95
4	4	WATER	27-JAN-95
5	5	WATER	27-JAN-95
6	6	WATER	27-JAN-95
7	7	WATER	27-JAN-95
8	8	WATER	27-JAN-95
9	9	WATER	27-JAN-95
10	10	WATER	27-JAN-95
11	11	WATER	27-JAN-95
12	12	WATER	27-JAN-95
13	13	WATER	27-JAN-95
14	14	WATER	27-JAN-95
15	15	WATER	27-JAN-95
16	16	WATER	27-JAN-95
17	17	WATER	27-JAN-95
18	18	WATER	27-JAN-95
19	19	WATER	27-JAN-95
20	20	WATER	27-JAN-95
21	21	WATER	27-JAN-95
22	22	WATER	27-JAN-95
23	23	WATER	27-JAN-95
24	24	WATER	27-JAN-95
25	25	WATER	27-JAN-95
26	26	WATER	27-JAN-95
27	27	WATER	27-JAN-95
28	28	WATER	27-JAN-95
29	29	WATER	27-JAN-95
30	30	WATER	27-JAN-95

## ---TOTALS---

<u>Matrix</u>	<u># Samples</u>
WATER	30

ATI STANDARD DISPOSAL PRACTICE

The sample(s) from this project will be disposed of in twenty-one (21) days from the date of this report. If an extended storage period is required, please contact our sample control department before the scheduled disposal date.

## ANALYTICAL SCHEDULE

Page

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: SSWD

ATI I.D.: 501

Analysis	Technique/Description
EPA 160.2 (TOTAL SUSPENDED SOLIDS)	GRAVIMETRIC
EPA 350.1 (AMMONIA AS NITROGEN)	COLORIMETRIC
EPA 353.2 (NITRATE-NITRITE AS NITROGEN)	COLORIMETRIC
EPA 365.2 (TOTAL PHOSPHATE AS PHOSPHORUS)	COLORIMETRIC
EPA 405.1 (BIOCHEMICAL OXYGEN DEMAND)	ELECTRODE
EPA 415.2 (TOTAL ORGANIC CARBON)	TOTAL ORGANIC CARBON ANALYZER



GENERAL CHEMISTRY RESULTS

Client : NCCOSC RDT&E DIVISION
Project # : (NONE)
Project Name: SSWD

ATI I.D.: 501244

Table with 4 columns: Sample #, Client ID, Matrix, Date Sampled, Date Received. Rows 1-5 show water samples from client 1, 2, 3, 4, 5, all dated 27-JAN-95.

Table with 7 columns: Parameter, Units, 1, 2, 3, 4, 5. Rows include BIOCHEMICAL OXYGEN DEMAND, AMMONIA AS NITROGEN, NITRATE-NITRITE AS NITROGEN, TOTAL PHOSPHATE AS PHOSPHORUS, TOTAL ORGANIC CARBON, and TOTAL SUSPENDED SOLIDS.



GENERAL CHEMISTRY RESULTS

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: SSWD

Pa-  
 ATI I.D.: 50

Sample #	Client ID	Matrix	Date Sampled	Date Received
6	6	WATER	27-JAN-95	27-JAN-95
7	7	WATER	27-JAN-95	27-JAN-95
8	8	WATER	27-JAN-95	27-JAN-95
9	9	WATER	27-JAN-95	27-JAN-95
10	10	WATER	27-JAN-95	27-JAN-95

Parameter	Units	6	7	8	9	10
BIOCHEMICAL OXYGEN DEMAND	MG/L	<5.0	<5.0	<5.0	<5.0	<5.0
TOTAL ORGANIC CARBON	MG/L	1.4	1.1	1.6	1.3	1.6

**GENERAL CHEMISTRY RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: SSWD

ATI I.D.: 501244

Sample #	Client ID	Matrix	Date Sampled	Date Received
11	11	WATER	27-JAN-95	27-JAN-95
12	12	WATER	27-JAN-95	27-JAN-95
13	13	WATER	27-JAN-95	27-JAN-95
14	14	WATER	27-JAN-95	27-JAN-95
15	15	WATER	27-JAN-95	27-JAN-95

Parameter	Units	11	12	13	14	15
BIOCHEMICAL OXYGEN DEMAND	MG/L	8.7	<5.0	<5.0	<5.0	<5.0
AMMONIA AS NITROGEN	MG/L	-	-	-	<0.20	-
NITRATE-NITRITE AS NITROGEN	MG/L	-	-	-	0.05	-
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	-	-	-	<0.10	-
TOTAL ORGANIC CARBON	MG/L	1.8	1.9	1.9	3.7	1.5
TOTAL SUSPENDED SOLIDS	MG/L	-	-	-	<20	-

**GENERAL CHEMISTRY RESULTS**

**Client : NCCOSC RDT&E DIVISION**  
**Project # : (NONE)**  
**Project Name: SSWD**

**ATI I.D.: 501**

Sample #	Client ID #	Matrix	Date Sampled	Date Received
16	16	WATER	27-JAN-95	27-JAN-95
17	17	WATER	27-JAN-95	27-JAN-95
18	18	WATER	27-JAN-95	27-JAN-95
19	19	WATER	27-JAN-95	27-JAN-95
20	20	WATER	27-JAN-95	27-JAN-95

Parameter	Units	16	17	18	19	20
BIOCHEMICAL OXYGEN DEMAND	MG/L	<5.0	<5.0	<5.0	<5.0	<5.0
AMMONIA AS NITROGEN	MG/L	-	-	-	<0.20	-
NITRATE-NITRITE AS NITROGEN	MG/L	-	-	-	0.05	-
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	-	-	-	<0.10	-
TOTAL ORGANIC CARBON	MG/L	2.5	2.0	2.6	2.9	2.1
TOTAL SUSPENDED SOLIDS	MG/L	-	-	-	46	-

**GENERAL CHEMISTRY RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: SSWD

ATI I.D.: 501244

Sample #	Client ID	Matrix	Date Sampled	Date Received
21	21	WATER	27-JAN-95	27-JAN-95
22	22	WATER	27-JAN-95	27-JAN-95
23	23	WATER	27-JAN-95	27-JAN-95
24	24	WATER	27-JAN-95	27-JAN-95
25	25	WATER	27-JAN-95	27-JAN-95

Parameter	Units	21	22	23	24	25
BIOCHEMICAL OXYGEN DEMAND	MG/L	<5.0	<5.0	<5.0	<5.0	<5.0
AMMONIA AS NITROGEN	MG/L	-	-	-	-	<0.20
NITRATE-NITRITE AS NITROGEN	MG/L	-	-	-	-	<0.05
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	-	-	-	-	<0.10
TOTAL ORGANIC CARBON	MG/L	2.2	2.2	1.6	1.8	2.0
TOTAL SUSPENDED SOLIDS	MG/L	-	-	-	-	<20

**GENERAL CHEMISTRY RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: SSWD

ATI I.D.: 501

Sample #	Client ID #	Matrix	Date Sampled	Date Received
26	26	WATER	27-JAN-95	27-JAN-95
27	27	WATER	27-JAN-95	27-JAN-95
28	28	WATER	27-JAN-95	27-JAN-95
29	29	WATER	27-JAN-95	27-JAN-95
30	30	WATER	27-JAN-95	27-JAN-95

Parameter	Units	26	27	28	29	30
BIOCHEMICAL OXYGEN DEMAND	MG/L	<5.0	<5.0	<5.0	<5.0	<5.0
TOTAL ORGANIC CARBON	MG/L	1.9	2.5	2.3	3.1	1.9
TOTAL SUSPENDED SOLIDS	MG/L	-	<20	-	-	-

**GENERAL CHEMISTRY - QUALITY CONTROL**
**DUP/MS**
**Client : NCCOSC RDT&E DIVISION**  
**Project # : (NONE)**  
**Project Name: SSWD**
**ATI I.D. : 501244**

Parameters	REF I.D.	Units	Sample Result	Dup Result	RPD	Spiked Sample	Spike Conc	% Rec
AMMONIA AS NITROGEN	501256-01	MG/L	<0.20	<0.20	0	1.9	2.0	95
BIOCHEMICAL OXYGEN DEMAND	501244-01	MG/L	<5.0	<5.0	0	N/A	N/A	N/A
BIOCHEMICAL OXYGEN DEMAND	501244-12	MG/L	<5.0	<5.0	0	N/A	N/A	N/A
BIOCHEMICAL OXYGEN DEMAND	501244-21	MG/L	<5.0	<5.0	0	N/A	N/A	N/A
NITRATE-NITRITE AS NITROGEN	501244-25	MG/L	<0.05	<0.05	0	1.9	2.0	95
TOTAL ORGANIC CARBON	501244-02	MG/L	2.3	2.3	0	22.5	20.0	101
TOTAL ORGANIC CARBON	501244-12	MG/L	1.9	1.6	17	21.1	20.0	96
TOTAL ORGANIC CARBON	501244-22	MG/L	2.2	2.2	0	23.3	20.0	106
TOTAL ORGANIC CARBON	501266-01	MG/L	8.8	8.7	1	29.8	20.0	105
TOTAL PHOSPHATE AS PHOSPHORUS	502027-05	MG/L	<0.10	<0.10	0	0.45	0.40	113
TOTAL SUSPENDED SOLIDS	501244-27	MG/L	<20	<20	0	N/A	N/A	N/A

**% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration**
**RPD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result**

**GENERAL CHEMISTRY - QUALITY CONTROL**
**BLANK SPIKE**
**Page**
**Client : NCCOSC RDT&E DIVISION**  
**Project # : (NONE)**  
**Project Name: SSWD**
**ATI I.D. : 501**

Parameters	Blank Spike ID#	Units	Blank Result	Spiked Sample	Spike Conc.	% R <sub>c</sub>
AMMONIA AS NITROGEN	54123	MG/L	<0.20	1.9	2.0	95
NITRATE-NITRITE AS NITROGEN	54047	MG/L	<0.05	2.0	2.0	10
TOTAL ORGANIC CARBON	54176	MG/L	<0.5	20.9	20.0	1C
TOTAL ORGANIC CARBON	54358	MG/L	<0.5	20.8	20	1C
TOTAL ORGANIC CARBON	54359	MG/L	<0.5	20.7	20	1C
TOTAL PHOSPHATE AS PHOSPHORUS	54167	MG/L	<0.10	0.44	0.40	11

$$\% \text{ Recovery} = (\text{Spike Sample Result} - \text{Sample Result}) * 100 / \text{Spike Concentration}$$

$$\text{RPD (Relative \% Difference)} = (\text{Sample Result} - \text{Duplicate Result}) * 100 / \text{Average Result}$$

ACCESSION #: 501244

INITIALS: [Signature]

**SAMPLE CONDITION UPON RECEIPT CHECKLIST**  
(FOR RE-ACCESSIONS, COMPLETE #7 THRU #9)

1	Does this project require special handling according to NEESA Levels C, D, AFOEHL or CLP protocols? If yes, complete a) thru c) a) Cooler temperature _____ b) pH sample aliquoted: yes / no / n/a c) LOT #'s: _____	YES	<u>NO</u>
2	Are custody seals present on cooler? If yes, are seals intact? <u>N/A</u>	YES	<u>NO</u>
3	Are custody seals present on sample containers? If yes, are seals intact? <u>N/A</u>	YES	<u>NO</u>
4	Is there a Chain-Of-Custody (COC)*?	<u>YES</u>	NO
5	Is the COC* complete? Relinquished: <u>yes</u> /no Requested analysis: <u>yes</u> /no	<u>YES</u>	NO
6	Is the COC* in agreement with the samples received? # Samples: <u>yes</u> /no Sample ID's: <u>yes</u> /no Date sampled: <u>yes</u> /no Matrix: <u>yes</u> /no # containers: <u>yes</u> /no	<u>YES</u>	NO
7	Are the samples preserved correctly?	<u>YES</u>	NO
8	Is there enough sample for all the requested analyses?	<u>YES</u>	NO
9	Are all samples within holding times for the requested analyses?	<u>YES</u>	NO
10	Cooler temperature: <u>NO COOLER - Samples went to truck, straight from field.</u>		
11	Were all sample containers received intact (ie. not broken, leaking, etc.)?	<u>YES</u>	NO
12	Are samples requiring no headspace, headspace free? N/A	<u>YES</u>	<u>NO</u>
13	Are VOA 1st stickers required?	YES	<u>NO</u>
14	Are there special comments on the Chain of Custody which require client contact?	YES	<u>N/A</u>
15	If yes, was ATI Project Manager notified?	YES	NO

Describe "no" items: #3 07,13 have approximately 1/2" headspace for BOD  
#5 10,12 have approximately 1 1/2" headspace for BOD.  
#19 has approximately 2" headspace for BOD

Was client contacted? yes / no  
If yes, Date: \_\_\_\_\_ Name of Person contacted: \_\_\_\_\_

Describe actions taken or client instructions: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

\*Or other representative documents, letters, and/or shipping memos



Analytical**Technologies, Inc.**

Corporate Offices 5550 Morehouse Drive San Diego, CA 92121 (619) 458-9141

ATI I.D.: 502027

February 13, 1995

NCCOSC RDT&E DIVISION  
53475 STROTHER ROAD RM 267A  
SAN DIEGO, CA 92152

Project Name: (NONE)  
Project # : (NONE)

Attention: STACY CURTIS

Analytical Technologies, Inc. has received the following sample(s):

<u>Date Received</u>	<u>Quantity</u>	<u>Matrix</u>
February 02, 1995	5	WATER

The sample(s) were analyzed with EPA methodology or equivalent methods as specified in the enclosed analytical schedule. The symbol for "less than" indicates a value below the reportable detection limit. If any flags appear next to the analytical data in this report, please see the attached list of flag definitions.

The results of these analyses and the quality control data are enclosed. Please note that the Sample Condition Upon Receipt Checklist is included at the end of this report.

JULIO PAREDES  
PROJECT MANAGER

ALAN J. KLEINSCHMIDT  
LABORATORY MANAGER

## SAMPLE CROSS REFERENCE

Page 1

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: (NONE)

Report Date: February 13, 1995  
ATI I.D. : 502027

ATI #	Client Description	Matrix	Date Collected
1	P2	WATER	02-FEB-95
2	P5	WATER	02-FEB-95
3	P6	WATER	02-FEB-95
4	P8	WATER	02-FEB-95
5	PULPER 1-8	WATER	02-FEB-95

## ---TOTALS---

<u>Matrix</u>	<u># Samples</u>
WATER	5

ATI STANDARD DISPOSAL PRACTICE

The sample(s) from this project will be disposed of in twenty-one (21) days from the date of this report. If an extended storage period is required, please contact our sample control department before the scheduled disposal date.



## ANALYTICAL SCHEDULE

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: (NONE)

Pa.  
ATI I.D.: 50

Analysis	Technique/Description
EPA 350.1 (AMMONIA AS NITROGEN)	COLORIMETRIC
EPA 353.2 (NITRATE-NITRITE AS NITROGEN)	COLORIMETRIC
EPA 365.2 (TOTAL PHOSPHATE AS PHOSPHORUS)	COLORIMETRIC

**GENERAL CHEMISTRY RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: (NONE)

ATI I.D.: 502027

Sample #	Client ID	Matrix	Date Sampled	Date Received
1	P2	WATER	02-FEB-95	02-FEB-95
2	P5	WATER	02-FEB-95	02-FEB-95
3	P6	WATER	02-FEB-95	02-FEB-95
4	P8	WATER	02-FEB-95	02-FEB-95
5	PULPER 1-8	WATER	02-FEB-95	02-FEB-95

Parameter	Units	1	2	3	4	5
AMMONIA AS NITROGEN	MG/L	<0.20	<0.20	<0.20	<0.20	<0.20
NITRATE-NITRITE AS NITROGEN	MG/L	0.43	0.15	0.16	<0.05	0.14
TOTAL PHOSPHATE AS PHOSPHORUS	MG/L	0.19	0.18	0.21	0.43	<0.10



GENERAL CHEMISTRY - QUALITY CONTROL

DUP/MS

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: (NONE)

ATI I.D. : 50

Parameters	REF I.D.	Units	Sample Result	Dup Result	RPD	Spiked Sample	Spike Conc	R
AMMONIA AS NITROGEN	502028-03	MG/L	7.4	7.3	1	18.0	10.0	1
NITRATE-NITRITE AS NITROGEN	502049-02	MG/L	<0.05	<0.05	0	2.0	2.0	1
TOTAL PHOSPHATE AS PHOSPHORUS	502027-05	MG/L	<0.10	<0.10	0	0.45	0.40	1

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration  
 RPD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result

**GENERAL CHEMISTRY - QUALITY CONTROL**
**BLANK SPIKE**

**Client : NCCOSC RDT&E DIVISION**  
**Project # : (NONE)**  
**Project Name: (NONE)**

**ATI I.D. : 502027**

Parameters	Blank Spike ID#	Units	Blank Result	Spiked Sample	Spike Conc.	% Rec
AMMONIA AS NITROGEN	54170	MG/L	<0.20	1.9	2.0	95
NITRATE-NITRITE AS NITROGEN	54325	MG/L	<0.05	2.0	2.0	100
TOTAL PHOSPHATE AS PHOSPHORUS	54167	MG/L	<0.10	0.44	0.40	110

**% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration**  
**RPD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result**



Analytical **Technologies, Inc.**

Corporate Offices 5550 Morehouse Drive San Diego, CA 92121 (619) 458-9141

ATI I.D.: 506082

June 29, 1995

NCCOSC RDT&E DIVISION  
53475 STROTHER ROAD RM 267A  
SAN DIEGO, CA 92152

Project Name: PRIORITY POLLUTANTS  
Project # : (NONE)

Attention: STACY CURTIS


Analytical Technologies, Inc. has received the following sample(s):

<u>Date Received</u>	<u>Quantity</u>	<u>Matrix</u>
June 08, 1995	2	SOLID

The sample(s) were analyzed with EPA methodology or equivalent methods as specified in the enclosed analytical schedule. The symbol for "less than" indicates a value below the reportable detection limit. If any flags appear next to the analytical data in this report, please see the attached list of flag definitions.

The results of these analyses and the quality control data are enclosed. Please note that the Sample Condition Upon Receipt Checklist is included at the end of this report.

  
JULIO PAREDES  
PROJECT MANAGER

  
ALAN J. KLEINSCHMIDT  
FOR LABORATORY MANAGER

Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: PRIORITY POLLUTANTS

Report Date: June 29, 1995  
ATI I.D. : 506082

ATI #	Client Description	Matrix	Date Collected
1	PULPER PAPER 01	SOLID	08-JUN-95
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95

---TOTALS---

<u>Matrix</u>	<u># Samples</u>
SOLID	2

ATI STANDARD DISPOSAL PRACTICE

The sample(s) from this project will be disposed of in twenty-one (21) days from the date of this report. If an extended storage period is required, please contact our sample control department before the scheduled disposal date.



Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: PRIORITY POLLUTANTS

ATI I.D.: 50

Analysis	Technique/Description
EPA 6010 (ANTIMONY)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 6010 (BERYLLIUM)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 6010 (CHROMIUM)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 6010 (COPPER)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 6010 (LEAD)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 6010 (NICKEL)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 6010 (SILVER)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 6010 (ZINC)	INDUCTIVELY COUPLED ARGON PLASMA
EPA 7060 (ARSENIC)	ATOMIC ABSORPTION/GRAPHITE FURNACE
EPA 7131 (CADMIUM)	ATOMIC ABSORPTION/GRAPHITE FURNACE
EPA 7471 (NON AQUEOUS MERCURY)	ATOMIC ABSORPTION/COLD VAPOR
EPA 7740 (SELENIUM)	ATOMIC ABSORPTION/GRAPHITE FURNACE
EPA 7841 (THALLIUM)	ATOMIC ABSORPTION/GRAPHITE FURNACE
EPA 8080 (ORGANOCHLORINE PESTICIDES & PCB'S)	GC/ELECTRON CAPTURE DETECTOR
EPA 8240 (GC/MS FOR VOLATILE ORGANICS)	GC/MASS SPECTROMETER
EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)	GC/MASS SPECTROMETER
EPA 9012 (TOTAL CYANIDE)	COLORIMETRIC
EPA 9066 (PHENOLS, TOTAL)	COLORIMETRIC
METHOD 7-2.2, METHODS OF SOIL ANALYSIS(% MOISTURE)	GRAVIMETRIC



Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: PRIORITY POLLUTANTS

ATI I.D.: 506082

Sample #	Client ID	Matrix	Date Sampled	Date Received
1	PULPER PAPER 01	SOLID	08-JUN-95	08-JUN-95
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95	08-JUN-95

Parameter	Units	1	2
TOTAL CYANIDE	MG/KG	<0.10	<0.10
% MOISTURE	%	84.4	83.9
PHENOLS, TOTAL	MG/KG	<0.20	<0.20



DUP/MS

Pa

Client : NCCOSC RDT&E DIVISION
Project # : (NONE)
Project Name: PRIORITY POLLUTANTS

ATI I.D. : 500

Table with 8 columns: Parameters, REF I.D., Units, Sample Result, Dup Result, RPD, Spiked Sample, Spike Conc. Rows include % MOISTURE, PHENOLS, TOTAL, and TOTAL CYANIDE.

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration
RPD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result



BLANK SPIKE

Client : NCCOSC RDT&E DIVISION
Project # : (NONE)
Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082

Table with 7 columns: Parameters, Blank Spike ID#, Units, Blank Result, Spiked Sample, Spike Conc., % Rec. Rows include PHENOLS, TOTAL; TOTAL CYANIDE; TOTAL CYANIDE.

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration
RPD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result

**METALS RESULTS**

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D.: 506

Sample #	Client ID	Matrix	Date Sampled	Date Received
1	PULPER PAPER 01	SOLID	08-JUN-95	08-JUN-95
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95	08-JUN-95

Parameter	Units	1	2
SILVER	MG/KG	<1.0	<1.0
ARSENIC	MG/KG	<1.0	<1.0
BERYLLIUM	MG/KG	<0.5	<0.5
CADMIUM	MG/KG	<0.5	<0.5
CHROMIUM	MG/KG	<0.5	<0.5
COPPER	MG/KG	1.7	<1.0
MERCURY	MG/KG	<0.25	<0.25
NICKEL	MG/KG	<1.0	<1.0
LEAD	MG/KG	<1.5	<1.5
ANTIMONY	MG/KG	<3.0	<3.0
SELENIUM	MG/KG	<1.0	<1.0
THALLIUM	MG/KG	<1.0	<1.0
ZINC	MG/KG	5.5	5.8

**METALS - QUALITY CONTROL**

DUP/MS

Page 7

 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082

Parameters	REF I.D.	Units	Sample Result	Dup Result	RPD	Spiked Sample	Spike Conc	% Rec
ANTIMONY	505309-32	MG/KG	<3.0	<3.0	0	48.0	49.8	96@v
ARSENIC	505309-32	MG/KG	1.4	1.5	7	45.8	49.9	89
BERYLLIUM	505309-32	MG/KG	<0.5	<0.5	0	44.5	49.7	90
CADMIUM	505309-32	MG/KG	<0.5	<0.5	0	45.6	49.9	91
CHROMIUM	505309-32	MG/KG	3.5	3.4	3	46.0	49.7	86
COPPER	505309-32	MG/KG	11.2	12.4	10	70.7	49.7	120
LEAD	505309-32	MG/KG	5.7	6.3	10	49.9	49.7	89
MERCURY	506114-01	MG/KG	0.54	0.47	14	1.25	1.00	71
NICKEL	505309-32	MG/KG	<1.0	<1.0	N/A@S	43.8	49.7	88
SELENIUM	505309-32	MG/KG	<1.0	<1.0	0	26.5	29.9	87
SILVER	505309-32	MG/KG	<1.0	<1.0	0	46.5	49.7	94
THALLIUM	505309-32	MG/KG	<1.0	<1.0	0	48.1	49.9	96
ZINC	505309-32	MG/KG	13.4	14.4	7	57.9	49.7	90

$$\% \text{ Recovery} = (\text{Spike Sample Result} - \text{Sample Result}) * 100 / \text{Spike Concentration}$$

$$\text{RPD (Relative \% Difference)} = (\text{Sample Result} - \text{Duplicate Result}) * 100 / \text{Average Result}$$



METALS - QUALITY CONTROL

BLANK SPIKE

Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

Parameters	Blank Spike ID#	Units	Blank Result	Spiked Sample	Spike Conc.	% Recovery
ANTIMONY	57011	MG/KG	<3.0	46.7	50.0	93
ARSENIC	57029	MG/KG	<1.0	45.9	50.0	92
BERYLLIUM	57011	MG/KG	<0.5	45.9	50.0	92
CADMIUM	57033	MG/KG	<0.5	47.0	50.0	94
CHROMIUM	57011	MG/KG	<0.5	47.3	50.0	95
COPPER	57011	MG/KG	<1.0	47.6	50.0	95
LEAD	57011	MG/KG	<1.5	47.8	50.0	96
MERCURY	57034	MG/KG	<0.25	1.06	1.00	100
NICKEL	57011	MG/KG	<1.0	47.5	50.0	95
SELENIUM	57027	MG/KG	<1.0	26.5	30.0	88
SILVER	57011	MG/KG	<1.0	46.0	50.0	92
THALLIUM	57032	MG/KG	<1.0	48.0	50.0	96
ZINC	57011	MG/KG	<2.0	47.8	50.0	96

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration  
 RPD (Relative % Difference) = (Sample Result - Duplicate Result)\*100/Average Result



Test : EPA 8080 (ORGANOCHLORINE PESTICIDES & PCB'S)  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082

Sample #	Client ID	Matrix	Date Sampled	Date Extracted	Date Analyzed	Dil. Factor
1	PULPER PAPER 01	SOLID	08-JUN-95	12-JUN-95	21-JUN-95	1.00
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95	12-JUN-95	21-JUN-95	1.00

Parameter	Units	1	2
ALDRIN	MG/KG	<0.032	<0.031
ALPHA-BHC	MG/KG	<0.032	<0.031
BETA-BHC	MG/KG	<0.032	<0.031
GAMMA-BHC (LINDANE)	MG/KG	<0.032	<0.031
DELTA-BHC	MG/KG	<0.032	<0.031
CHLORDANE	MG/KG	<0.32	<0.31
2,4'-DDD	MG/KG	<0.064	<0.062
2,4'-DDE	MG/KG	<0.064	<0.062
2,4'-DDT	MG/KG	<0.064	<0.062
4,4'-DDD	MG/KG	<0.064	<0.062
4,4'-DDE	MG/KG	<0.064	<0.062
4,4'-DDT	MG/KG	<0.064	<0.062
DIELDRIN	MG/KG	<0.064	<0.062
ENDOSULFAN I	MG/KG	<0.032	<0.031
ENDOSULFAN II	MG/KG	<0.064	<0.062
ENDOSULFAN SULFATE	MG/KG	<0.064	<0.062
ENDRIN	MG/KG	<0.064	<0.062
ENDRIN KETONE	MG/KG	<0.064	<0.062
HEPTACHLOR	MG/KG	<0.032	<0.031
HEPTACHLOR EPOXIDE	MG/KG	<0.032	<0.031
METHOXYCHLOR	MG/KG	<0.32	<0.31
TOXAPHENE	MG/KG	<0.64	<0.62
AROCLOR-1016	MG/KG	<0.32	<0.31
AROCLOR-1221	MG/KG	<0.32	<0.31
AROCLOR-1232	MG/KG	<0.32	<0.31
AROCLOR-1242	MG/KG	<0.32	<0.31
AROCLOR-1248	MG/KG	<0.32	<0.31
AROCLOR-1254	MG/KG	<0.32	<0.31
AROCLOR-1260	MG/KG	<0.32	<0.31

SURROGATES

DBC	%	75	70
-----	---	----	----



## REAGENT BLANK

Page

Test : EPA 8080 (ORGANOCHLORINE PESTICIDES & PCB'S)  
 Blank I.D. : 35764  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082  
 Date Extracted: 12-JUN-95  
 Date Analyzed : 20-JUN-95  
 Dil. Factor : 1.00

Parameters	Units	Results
ALDRIN	MG/KG	<0.0050
ALPHA-BHC	MG/KG	<0.0050
BETA-BHC	MG/KG	<0.0050
GAMMA-BHC (LINDANE)	MG/KG	<0.0050
DELTA-BHC	MG/KG	<0.0050
CHLORDANE	MG/KG	<0.050
2,4'-DDD	MG/KG	<0.010
2,4'-DDE	MG/KG	<0.010
2,4'-DDT	MG/KG	<0.010
4,4'-DDD	MG/KG	<0.010
4,4'-DDE	MG/KG	<0.010
4,4'-DDT	MG/KG	<0.010
DIELDRIN	MG/KG	<0.010
ENDOSULFAN I	MG/KG	<0.0050
ENDOSULFAN II	MG/KG	<0.010
ENDOSULFAN SULFATE	MG/KG	<0.010
ENDRIN	MG/KG	<0.010
ENDRIN KETONE	MG/KG	<0.010
HEPTACHLOR	MG/KG	<0.0050
HEPTACHLOR EPOXIDE	MG/KG	<0.0050
METHOXYCHLOR	MG/KG	<0.050
TOXAPHENE	MG/KG	<0.10
AROCLOR-1016	MG/KG	<0.050
AROCLOR-1221	MG/KG	<0.050
AROCLOR-1232	MG/KG	<0.050
AROCLOR-1242	MG/KG	<0.050
AROCLOR-1248	MG/KG	<0.050
AROCLOR-1254	MG/KG	<0.050
AROCLOR-1260	MG/KG	<0.050

SURROGATES

DBC

%

74



MSMSD

Test : EPA 8080 (ORGANOCHLORINE PESTICIDES & PCB'S)  
 MSMSD # : 76499  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082  
 Date Extracted: 12-JUN-95  
 Date Analyzed : 20-JUN-95  
 Sample Matrix : SOIL  
 REF I.D. : 506082-01

Parameters	Units	Sample Result	Conc Spike	Spiked Sample	% Rec	Dup Spike	Dup % Rec	RPD
ALDRIN	MG/KG	<0.0050	0.21	0.19	90	0.17	81	11
GAMMA-BHC (LINDANE)	MG/KG	<0.0050	0.21	0.15	71	0.15	71	0
4,4'-DDT	MG/KG	<0.010	0.43	0.35	81	0.29	67	19
DIELDRIN	MG/KG	<0.010	0.43	0.34	79	0.29	67	16
ENDRIN	MG/KG	<0.010	0.43	0.36	84	0.32	74	12
HEPTACHLOR	MG/KG	<0.0050	0.21	0.17	81	0.15	71	13

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration

RPD (Relative % Difference) = (Spiked Sample Result - Duplicate Spike Result)\*100/Average Result



## BLANK SPIKE

Pag

Test : EPA 8080 (ORGANOCHLORINE PESTICIDES & PCB'S)  
Blank Spike #: 57184  
Client : NCCOSC RDT&E DIVISION  
Project #: (NONE)  
Project Name : PRIORITY POLLUTANTS

ATI I.D. : 506082  
Date Extracted: 12-JUN-95  
Date Analyzed : 20-JUN-95  
Sample Matrix : SOIL

Parameters	Units	Blank Result	Spiked Sample	Spike Conc.	R
ALDRIN	MG/KG	<0.0050	0.023	0.033	7
GAMMA-BHC (LINDANE)	MG/KG	<0.0050	0.024	0.033	7
4,4'-DDT	MG/KG	<0.010	0.059	0.067	8
DIELDRIN	MG/KG	<0.010	0.054	0.067	8
ENDRIN	MG/KG	<0.010	0.057	0.067	8
HEPTACHLOR	MG/KG	<0.0050	0.027	0.033	8

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration

RPD (Relative % Difference) = (Spiked Sample - Blank Result)\*100/Average Result



Test : EPA 8240 (GC/MS FOR VOLATILE ORGANICS)  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082

Sample #	Client ID	Matrix	Date Sampled	Date Extracted	Date Analyzed	Dil. Factor
1	PULPER PAPER 01	SOLID	08-JUN-95	09-JUN-95	19-JUN-95	1.00
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95	09-JUN-95	19-JUN-95	1.00

Parameter	Units	1	2
CHLOROMETHANE	MG/KG	<0.5	<0.5
VINYL CHLORIDE	MG/KG	<0.3	<0.3
BROMOMETHANE	MG/KG	<0.5	<0.5
CHLOROETHANE	MG/KG	<0.3	<0.3
ACETONE	MG/KG	2.1	1.2
1,1-DICHLOROETHENE	MG/KG	<0.05	<0.05
METHYLENE CHLORIDE	MG/KG	<0.3	<0.3
CARBON DISULFIDE	MG/KG	<0.1	<0.1
TRANS-1,2-DICHLOROETHENE	MG/KG	<0.05	<0.05
1,1-DICHLOROETHANE	MG/KG	<0.05	<0.05
CIS-1,2-DICHLOROETHENE	MG/KG	<0.05	<0.05
CHLOROFORM	MG/KG	<0.05	<0.05
2-BUTANONE (MEK)	MG/KG	<0.5	<0.5
1,1,1-TRICHLOROETHANE	MG/KG	<0.05	<0.05
CARBON TETRACHLORIDE	MG/KG	<0.05	<0.05
1,2-DICHLOROETHANE	MG/KG	<0.05	<0.05
BENZENE	MG/KG	<0.05	<0.05
TRICHLOROETHENE	MG/KG	<0.05	<0.05
1,2-DICHLOROPROPANE	MG/KG	<0.05	<0.05
BROMODICHLOROMETHANE	MG/KG	<0.05	<0.05
4-METHYL-2-PENTANONE (MIBK)	MG/KG	<0.5	<0.5
CIS-1,3-DICHLOROPROPENE	MG/KG	<0.05	<0.05
TOLUENE	MG/KG	<0.1	<0.1
TRANS-1,3-DICHLOROPROPENE	MG/KG	<0.05	<0.05
2-HEXANONE (MBK)	MG/KG	<0.5	<0.5
1,1,2-TRICHLOROETHANE	MG/KG	<0.05	<0.05
TETRACHLOROETHENE	MG/KG	<0.05	<0.05
DIBROMOCHLOROMETHANE	MG/KG	<0.05	<0.05
CHLOROBENZENE	MG/KG	<0.05	<0.05
ETHYLBENZENE	MG/KG	<0.05	<0.05
XYLENES (TOTAL)	MG/KG	<0.05	<0.05
STYRENE	MG/KG	<0.05	<0.05
BROMOFORM	MG/KG	<0.3	<0.3
1,1,2,2-TETRACHLOROETHANE	MG/KG	<0.1	<0.1
DICHLORODIFLUOROMETHANE	MG/KG	<0.5	<0.5
TRICHLOROFLUOROMETHANE	MG/KG	<0.3	<0.3
1,1,2-TRICHLORO-1,2,2-TRIFLUOROETHANE	MG/KG	<0.3	<0.3
1,2-DICHLOROBENZENE	MG/KG	<0.3	<0.3



Test : EPA 8240 (GC/MS FOR VOLATILE ORGANICS)  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082

Sample #	Client ID	Matrix	Date Sampled	Date Extracted	Date Analyzed	Dil. Factor
1	PULPER PAPER 01	SOLID	08-JUN-95	09-JUN-95	19-JUN-95	1.00
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95	09-JUN-95	19-JUN-95	1.00

Parameter	Units	1	2
1,3-DICHLOROBENZENE	MG/KG	<0.3	<0.3
1,4-DICHLOROBENZENE	MG/KG	<0.3	<0.3
<b>SURROGATES</b>			
1,2-DICHLOROETHANE-D4	%	48@H	56@H
TOLUENE-D8	%	49@H	61
BFB	%	47@H	57



ADDITIONAL COMPOUNDS (SEMI-QUANTITATED)

Method : EPA 8240 (GC/MS FOR VOLATILE ORGANICS)  
Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: PRIORITY POLLUTANTS

SOLID  
ATI I.D.: 506082

Sample Parameters	Units	Results
1 UNKNOWN HYDROCARBON	MG/KG	0.3
METHYL ACETATE	MG/KG	0.6
2 UNKNOWN HYDROCARBON	MG/KG	0.3
METHYL ACETATE	MG/KG	0.7



## REAGENT BLANK

Page

Test : EPA 8240 (GC/MS FOR VOLATILE ORGANICS)  
Blank I.D. : 35756  
Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082  
Date Extracted: 09-JUN-95  
Date Analyzed : 19-JUN-95  
Dil. Factor : 1.00

Parameters	Units	Results
CHLOROMETHANE	MG/KG	<0.5
VINYL CHLORIDE	MG/KG	<0.3
BROMOMETHANE	MG/KG	<0.5
CHLOROETHANE	MG/KG	<0.3
ACETONE	MG/KG	<0.5
1,1-DICHLOROETHENE	MG/KG	<0.05
METHYLENE CHLORIDE	MG/KG	<0.3
CARBON DISULFIDE	MG/KG	<0.1
TRANS-1,2-DICHLOROETHENE	MG/KG	<0.05
1,1-DICHLOROETHANE	MG/KG	<0.05
CIS-1,2-DICHLOROETHENE	MG/KG	<0.05
CHLOROFORM	MG/KG	<0.05
2-BUTANONE (MEK)	MG/KG	<0.5
1,1,1-TRICHLOROETHANE	MG/KG	<0.05
CARBON TETRACHLORIDE	MG/KG	<0.05
1,2-DICHLOROETHANE	MG/KG	<0.05
BENZENE	MG/KG	<0.05
TRICHLOROETHENE	MG/KG	<0.05
1,2-DICHLOROPROPANE	MG/KG	<0.05
BROMODICHLOROMETHANE	MG/KG	<0.05
4-METHYL-2-PENTANONE (MIBK)	MG/KG	<0.5
CIS-1,3-DICHLOROPROPENE	MG/KG	<0.05
TOLUENE	MG/KG	<0.1
TRANS-1,3-DICHLOROPROPENE	MG/KG	<0.05
2-HEXANONE (MBK)	MG/KG	<0.5
1,1,2-TRICHLOROETHANE	MG/KG	<0.05
TETRACHLOROETHENE	MG/KG	<0.05
DIBROMOCHLOROMETHANE	MG/KG	<0.05
CHLOROBENZENE	MG/KG	<0.05
ETHYLBENZENE	MG/KG	<0.05
XYLENES (TOTAL)	MG/KG	<0.05
STYRENE	MG/KG	<0.05
BROMOFORM	MG/KG	<0.3
1,1,2,2-TETRACHLOROETHANE	MG/KG	<0.1
DICHLORODIFLUOROMETHANE	MG/KG	<0.5
TRICHLOROFLUOROMETHANE	MG/KG	<0.3
1,1,2-TRICHLORO-1,2,2-TRIFLUOROETHANE	MG/KG	<0.3
1,2-DICHLOROBENZENE	MG/KG	<0.3
1,3-DICHLOROBENZENE	MG/KG	<0.3
1,4-DICHLOROBENZENE	MG/KG	<0.3
<u>SURROGATES</u>		
1,2-DICHLOROETHANE-D4	%	88
TOLUENE-D8	%	91
BFB	%	91



REAGENT BLANK  
ADDITIONAL COMPOUNDS (SEMI-QUANTITATED)

Test : EPA 8240 (GC/MS FOR VOLATILE ORGANICS)

Blank I.D. : 35756

ATI I.D. : 506082

Client : NCCOSC RDT&E DIVISION

Project # : (NONE)

Project Name: PRIORITY POLLUTANTS

Parameters	Units	Results
NONE DETECTED	N/A	N/A



MSMSD

Test : EPA 8240 (GC/MS FOR VOLATILE ORGANICS)  
 MSMSD # : 76484  
 Client : NCCOSC RDT&E DIVISION

ATI I.D. : 506082  
 Date Extracted: 09-JUN-95  
 Date Analyzed : 19-JUN-95  
 Sample Matrix : SOIL  
 REF I.D. : 506082-02

Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

Parameters	Units	Sample Result	Conc Spike	Spiked Sample	% Rec	Dup Spike	Dup % Rec	RF
1,1-DICHLOROETHENE	MG/KG	<0.05	2.5	0.81*H	32	0.84*H	34	4
BENZENE	MG/KG	<0.05	2.5	1.31*H	52	1.42*H	57	8
TRICHLOROETHENE	MG/KG	<0.05	2.5	1.24*H	50	1.33*H	53	7
TOLUENE	MG/KG	<0.1	2.5	1.49*H	60	1.56*H	62	5
CHLOROBENZENE	MG/KG	<0.05	2.5	1.60*H	64	1.65*H	66	3

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration  
 RPD (Relative % Difference) = (Spiked Sample Result - Duplicate Spike Result)\*100/Average Re-



## BLANK SPIKE

Page 19

Test : EPA 8240 (GC/MS FOR VOLATILE ORGANICS)  
Blank Spike #: 57172  
Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name : PRIORITY POLLUTANTS

ATI I.D. : 506082  
Date Extracted: 09-JUN-95  
Date Analyzed : 19-JUN-95  
Sample Matrix : SOIL

Parameters	Units	Blank Result	Spiked Sample	Spike Conc.	% Rec
1,1-DICHLOROETHENE	MG/KG	<0.05	1.6	2.5	64
BENZENE	MG/KG	<0.05	2.4	2.5	96
TRICHLOROETHENE	MG/KG	<0.05	2.3	2.5	92
TOLUENE	MG/KG	<0.1	2.7	2.5	108
CHLOROBENZENE	MG/KG	<0.05	2.8	2.5	112

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration

RPD (Relative % Difference) = (Spiked Sample - Blank Result)\*100/Average Result



Test : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082

Sample #	Client ID	Matrix	Date Sampled	Date Extracted	Date Analyzed	Dil. Factor
1	PULPER PAPER 01	SOLID	08-JUN-95	12-JUN-95	20-JUN-95	1.00
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95	12-JUN-95	17-JUN-95	5.00

Parameter	Units	1	2
N-NITROSODIMETHYLAMINE	MG/KG	<0.17	<0.17
PYRIDINE	MG/KG	<0.17	<0.17
PHENOL	MG/KG	<0.17	<0.17
ANILINE	MG/KG	<0.34	<0.34
BIS(2-CHLOROETHYL) ETHER	MG/KG	<0.17	<0.17
2-CHLOROPHENOL	MG/KG	<0.17	<0.17
1,3-DICHLOROBENZENE	MG/KG	<0.17	<0.17
1,4-DICHLOROBENZENE	MG/KG	<0.17	<0.17
BENZYL ALCOHOL	MG/KG	<0.17	<0.17
1,2-DICHLOROBENZENE	MG/KG	<0.17	<0.17
2-METHYLPHENOL	MG/KG	<0.17	<0.17
BIS(2-CHLOROISOPROPYL) ETHER	MG/KG	<0.17	<0.17
4-METHYLPHENOL	MG/KG	<0.17	<0.17
N-NITroso-DI-N-PROPYLAMINE	MG/KG	<0.17	<0.17
HEXACHLOROETHANE	MG/KG	<0.17	<0.17
NITROBENZENE	MG/KG	<0.17	<0.17
ISOPHORONE	MG/KG	<0.17	<0.17
2-NITROPHENOL	MG/KG	<0.17	<0.17
2,4-DIMETHYLPHENOL	MG/KG	<0.17	<0.17
BENZOIC ACID	MG/KG	<0.85	<0.85
BIS(2-CHLOROETHOXY) METHANE	MG/KG	<0.17	<0.17
2,4-DICHLOROPHENOL	MG/KG	<0.17	<0.17
1,2,4-TRICHLOROBENZENE	MG/KG	<0.17	<0.17
NAPHTHALENE	MG/KG	<0.17	<0.17
4-CHLOROANILINE	MG/KG	<0.50	<0.50
HEXACHLOROBUTADIENE	MG/KG	<0.17	<0.17
4-CHLORO-3-METHYLPHENOL	MG/KG	<0.17	<0.17
2-METHYLNAPHTHALENE	MG/KG	<0.17	<0.17
HEXACHLOROCYCLOPENTADIENE	MG/KG	<0.17	<0.17
2,4,6-TRICHLOROPHENOL	MG/KG	<0.17	<0.17
2,4,5-TRICHLOROPHENOL	MG/KG	<0.85	<0.85
2-CHLORONAPHTHALENE	MG/KG	<0.17	<0.17
2-NITROANILINE	MG/KG	<0.85	<0.85
DIMETHYLPHTHALATE	MG/KG	0.29	<0.17
ACENAPHTHYLENE	MG/KG	<0.17	<0.17
2,6-DINITROTOLUENE	MG/KG	<0.17	<0.17
3-NITROANILINE	MG/KG	<0.85	<0.85
ACENAPHTHENE	MG/KG	<0.17	<0.17



Test : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082

Sample #	Client ID	Matrix	Date Sampled	Date Extracted	Date Analyzed	Dil. Factor
1	PULPER PAPER 01	SOLID	08-JUN-95	12-JUN-95	20-JUN-95	1.00
2	PULPER PAPER 01/DUPLICATE	SOLID	08-JUN-95	12-JUN-95	17-JUN-95	5.00

Parameter	Units	1	2
2,4-DINITROPHENOL	MG/KG	<0.85	<0.85
4-NITROPHENOL	MG/KG	<0.85	<0.85
DIBENZOFURAN	MG/KG	<0.17	<0.17
2,4-DINITROTOLUENE	MG/KG	<0.17	<0.17
DIETHYLPHTHALATE	MG/KG	<0.17	<0.17
4-CHLOROPHENYL-PHENYLETHER	MG/KG	<0.17	<0.17
FLUORENE	MG/KG	<0.17	<0.17
4-NITROANILINE	MG/KG	<0.85	<0.85
2-METHYL-4,6-DINITROPHENOL	MG/KG	<0.85	<0.85
N-NITROSODIPHENYLAMINE	MG/KG	<0.17	<0.17
4-BROMOPHENYL-PHENYLETHER	MG/KG	<0.17	<0.17
HEXACHLOROBENZENE	MG/KG	<0.17	<0.17
PENTACHLOROPHENOL	MG/KG	<0.85	<0.85
PHENANTHRENE	MG/KG	<0.17	<0.17
ANTHRACENE	MG/KG	<0.17	<0.17
DI-N-BUTYLPHTHALATE	MG/KG	<0.17	<0.17
FLUORANTHENE	MG/KG	<0.17	<0.17
PYRENE	MG/KG	<0.17	<0.17
BUTYLBENZYLPHTHALATE	MG/KG	<0.17	<0.17
3,3'-DICHLOROBENZIDINE	MG/KG	<0.34	<0.34
BENZO(a)ANTHRACENE	MG/KG	<0.17	<0.17
CHRYSENE	MG/KG	<0.17	<0.17
BIS(2-ETHYLHEXYL) PHTHALATE	MG/KG	<0.17	<0.17
DI-N-OCTYLPHTHALATE	MG/KG	<0.17	<0.17
BENZO(b) FLUORANTHENE	MG/KG	<0.17	<0.17
BENZO(k) FLUORANTHENE	MG/KG	<0.17	<0.17
BENZO(a) PYRENE	MG/KG	<0.17	<0.17
INDENO(1,2,3-cd) PYRENE	MG/KG	<0.17	<0.17
DIBENZ(a,h) ANTHRACENE	MG/KG	<0.17	<0.17
BENZO(g,h,i) PERYLENE	MG/KG	<0.17	<0.17

SURROGATES

NITROBENZENE-D5	%	76	81
2-FLUOROBIPHENYL	%	72	94
TERPHENYL-D14	%	79	79
PHENOL-D6	%	81	83
2-FLUOROPHENOL	%	66	68
2,4,6-TRIBROMOPHENOL	%	88	98



Method : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)  
Client : NCCOSC RDT&E DIVISION  
Project # : (NONE)  
Project Name: PRIORITY POLLUTANTS

SOLID  
ATI I.D.: 506082

Sample Parameters	Units	Results
1 ALIPHATIC HYDROCARBONS (C-16)	MG/KG	2
ALIPHATIC HYDROCARBONS (C-16, C-17)	MG/KG	0.6
ALIPHATIC HYDROCARBONS (C-17)	MG/KG	0.5
ALIPHATIC HYDROCARBONS (C-18, C-19)	MG/KG	0.7
ALIPHATIC HYDROCARBONS (C-20)	MG/KG	0.4
2 ALIPHATIC HYDROCARBONS (C-16)	MG/KG	1
ALIPHATIC HYDROCARBONS (C-16, C-17)	MG/KG	2
ALIPHATIC HYDROCARBONS (C-17)	MG/KG	2
ALIPHATIC HYDROCARBONS (C-18, C-19)	MG/KG	2
ALIPHATIC HYDROCARBONS (C-20)	MG/KG	1

**REAGENT BLANK**

Test : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)  
 Blank I.D. : 35693  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082  
 Date Extracted: 12-JUN-95  
 Date Analyzed : 15-JUN-95  
 Dil. Factor : 1.00

Parameters	Units	Results
N-NITROSODIMETHYLAMINE	MG/KG	<0.17
PYRIDINE	MG/KG	<0.17
PHENOL	MG/KG	<0.17
ANILINE	MG/KG	<0.34
BIS(2-CHLOROETHYL) ETHER	MG/KG	<0.17
2-CHLOROPHENOL	MG/KG	<0.17
1,3-DICHLOROBENZENE	MG/KG	<0.17
1,4-DICHLOROBENZENE	MG/KG	<0.17
BENZYL ALCOHOL	MG/KG	<0.17
1,2-DICHLOROBENZENE	MG/KG	<0.17
2-METHYLPHENOL	MG/KG	<0.17
BIS(2-CHLOROISOPROPYL) ETHER	MG/KG	<0.17
4-METHYLPHENOL	MG/KG	<0.17
N-NITROSO-DI-N-PROPYLAMINE	MG/KG	<0.17
HEXACHLOROETHANE	MG/KG	<0.17
NITROBENZENE	MG/KG	<0.17
ISOPHORONE	MG/KG	<0.17
2-NITROPHENOL	MG/KG	<0.17
2,4-DIMETHYLPHENOL	MG/KG	<0.17
BENZOIC ACID	MG/KG	<0.85
BIS(2-CHLOROETHOXY) METHANE	MG/KG	<0.17
2,4-DICHLOROPHENOL	MG/KG	<0.17
1,2,4-TRICHLOROBENZENE	MG/KG	<0.17
NAPHTHALENE	MG/KG	<0.17
4-CHLOROANILINE	MG/KG	<0.50
HEXACHLOROBUTADIENE	MG/KG	<0.17
4-CHLORO-3-METHYLPHENOL	MG/KG	<0.17
2-METHYLNAPHTHALENE	MG/KG	<0.17
HEXACHLOROCYCLOPENTADIENE	MG/KG	<0.17
2,4,6-TRICHLOROPHENOL	MG/KG	<0.17
2,4,5-TRICHLOROPHENOL	MG/KG	<0.85
2-CHLORONAPHTHALENE	MG/KG	<0.17
2-NITROANILINE	MG/KG	<0.85
DIMETHYLPHTHALATE	MG/KG	<0.17
ACENAPHTHYLENE	MG/KG	<0.17
2,6-DINITROTOLUENE	MG/KG	<0.17
3-NITROANILINE	MG/KG	<0.85
ACENAPHTHENE	MG/KG	<0.17
2,4-DINITROPHENOL	MG/KG	<0.85
4-NITROPHENOL	MG/KG	<0.85
DIBENZOFURAN	MG/KG	<0.17
2,4-DINITROTOLUENE	MG/KG	<0.17
DIETHYLPHTHALATE	MG/KG	<0.17
4-CHLOROPHENYL-PHENYLEETHER	MG/KG	<0.17
FLUORENE	MG/KG	<0.17
4-NITROANILINE	MG/KG	<0.85



## REAGENT BLANK

Page:

Test : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)  
 Blank I.D. : 35693  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082  
 Date Extracted: 12-JUN-95  
 Date Analyzed : 15-JUN-95  
 Dil. Factor : 1.00

Parameters	Units	Results
2-METHYL-4,6-DINITROPHENOL	MG/KG	<0.85
N-NITROSODIPHENYLAMINE	MG/KG	<0.17
4-BROMOPHENYL-PHENYLETHER	MG/KG	<0.17
HEXACHLOROBENZENE	MG/KG	<0.17
PENTACHLOROPHENOL	MG/KG	<0.85
PHENANTHRENE	MG/KG	<0.17
ANTHRACENE	MG/KG	<0.17
DI-N-BUTYLPHTHALATE	MG/KG	<0.17
FLUORANTHENE	MG/KG	<0.17
PYRENE	MG/KG	<0.17
BUTYLBENZYLPHTHALATE	MG/KG	<0.17
3,3'-DICHLOROBENZIDINE	MG/KG	<0.34
BENZO(a)ANTHRACENE	MG/KG	<0.17
CHRYSENE	MG/KG	<0.17
BIS(2-ETHYLHEXYL) PHTHALATE	MG/KG	<0.17
DI-N-OCTYLPHTHALATE	MG/KG	<0.17
BENZO(b)FLUORANTHENE	MG/KG	<0.17
BENZO(k)FLUORANTHENE	MG/KG	<0.17
BENZO(a)PYRENE	MG/KG	<0.17
INDENO(1,2,3-cd)PYRENE	MG/KG	<0.17
DIBENZ(a,h)ANTHRACENE	MG/KG	<0.17
BENZO(g,h,i)PERYLENE	MG/KG	<0.17
<u>SURROGATES</u>		
NITROBENZENE-D5	%	68
2-FLUOROBIPHENYL	%	72
TERPHENYL-D14	%	66
PHENOL-D6	%	67
2-FLUOROPHENOL	%	59
2,4,6-TRIBROMOPHENOL	%	81



Analytical **Technologies, Inc**

GAS CHROMATOGRAPHY/MASS SPECTROSCOPY - QUALITY CONTROL

REAGENT BLANK  
ADDITIONAL COMPOUNDS (SEMI-QUANTITATED)

Page 25

Test : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)

Blank I.D. : 35693

ATI I.D.

: 506082

Client : NCCOSC RDT&E DIVISION

Project # : (NONE)

Project Name: PRIORITY POLLUTANTS

Parameters	Units	Results
UNKNOWN HYDROCARBONS	MG/KG	0.2
UNKNOWN HYDROCARBONS	MG/KG	0.2



## MSMSD

Page:

Test : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)  
 MSMSD # : 76364  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name: PRIORITY POLLUTANTS

ATI I.D. : 506082  
 Date Extracted: 12-JUN-95  
 Date Analyzed : 20-JUN-95  
 Sample Matrix : SOIL  
 REF I.D. : 506082-02

Parameters	Units	Sample Result	Conc Spike	Spiked Sample	% Rec	Dup Spike	Dup % Rec	R'
PHENOL	MG/KG	<0.85	5.0	4.7	94	3.9	78	1'
2-CHLOROPHENOL	MG/KG	<0.85	5.0	4.0	80	3.3	66	1'
1,4-DICHLOROBENZENE	MG/KG	<0.85	3.3	2.5	76	2.2	67	1
N-NITROSO-DI-N-PROPYLAMINE	MG/KG	<0.85	3.3	3.3	100	2.7	82	2
1,2,4-TRICHLOROBENZENE	MG/KG	<0.85	3.3	3.1	94	2.5	76	2
4-CHLORO-3-METHYLPHENOL	MG/KG	<0.85	5.0	4.6	92	3.9	78	1
ACENAPHTHENE	MG/KG	<0.85	3.3	3.9	118	3.2	97	2
4-NITROPHENOL	MG/KG	<4.3	5.0	3.4	68	2.7	54	2
2,4-DINITROTOLUENE	MG/KG	<0.85	3.3	2.6	79	2.1	64	2
PENTACHLOROPHENOL	MG/KG	<4.3	5.0	3.3	66	2.6	52	2
PYRENE	MG/KG	<0.85	3.3	2.8	85	2.2	67	2

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration

RPD (Relative % Difference) = (Spiked Sample Result - Duplicate Spike Result)\*100/Average Re



BLANK SPIKE

Test : EPA 8270 (GC/MS FOR SEMIVOLATILE ORGANICS)  
 Blank Spike #: 57065  
 Client : NCCOSC RDT&E DIVISION  
 Project # : (NONE)  
 Project Name : PRIORITY POLLUTANTS

ATI I.D. : 506082  
 Date Extracted: 12-JUN-95  
 Date Analyzed : 15-JUN-95  
 Sample Matrix : SOIL

Parameters	Units	Blank Result	Spiked Sample	Spike Conc.	% Rec
PHENOL	MG/KG	<0.17	2.7	5.0	54
2-CHLOROPHENOL	MG/KG	<0.17	3.0	5.0	60
1,4-DICHLOROBENZENE	MG/KG	<0.17	2.1	3.3	64
N-NITROSO-DI-N-PROPYLAMINE	MG/KG	<0.17	2.2	3.3	67
1,2,4-TRICHLOROBENZENE	MG/KG	<0.17	2.3	3.3	70
4-CHLORO-3-METHYLPHENOL	MG/KG	<0.17	3.2	5.0	64
ACENAPHTHENE	MG/KG	<0.17	2.5	3.3	76
4-NITROPHENOL	MG/KG	<0.85	3.6	5.0	72
2,4-DINITROTOLUENE	MG/KG	<0.17	2.3	3.3	70
PENTACHLOROPHENOL	MG/KG	<0.85	3.8	5.0	76
PYRENE	MG/KG	<0.17	2.4	3.3	73

% Recovery = (Spike Sample Result - Sample Result)\*100/Spike Concentration  
 RPD (Relative % Difference) = (Spiked Sample - Blank Result)\*100/Average Result

ANALYTICAL TECHNOLOGIES, INC.  
SAN DIEGO  
FLAGS

INORGANICS

FLAG MESSAGE DESCRIPTION

B ABSOLUTE VALUE OF ANALYTE CONCENTRATION IS < CRDL BUT  $\geq$  THE IDL  
BB RESULT BETWEEN IDL AND LOQ  
D POST DIGESTION SPIKE FOR GFAA OUTSIDE LIMITS AFTER 1:25 DILUTION. SAMPLE REPORTED AT ORIGINAL CONCENTRATION.  
E ESTIMATED VALUE DUE TO INTERFERENCE  
M DUPLICATE INJECTION PRECISION NOT MET  
N SPIKED SAMPLE RECOVERY NOT WITHIN CONTROL LIMITS  
S REPORTED VALUE WAS DETERMINED BY METHOD OF STANDARD ADDITIONS  
U COMPOUND WAS ANALYZED FOR BUT NOT DETECTED  
W POST DIGESTION SPIKE OUT OF CONTROL LIMITS; SAMPLE ABSORBANCE < 50% OF SPIKE ABSORBANCE FOR GF/AA  
X ABSOLUTE VALUE OF ANALYTE CONCENTRATION IS LESS THAN 3 TIMES THE MDL  
\* DUPLICATE ANALYSIS NOT WITHIN CONTROL LIMITS  
+ CORRELATION COEFFICIENT FOR MSA IS LESS THAN 0.995  
\*H RESULTS OUTSIDE OF LIMITS DUE TO SAMPLE MATRIX INTERFERENCE  
\*Q INSUFFICIENT SAMPLE FOR ANALYSIS  
\*R DATA IS NOT USABLE  
\*V SAMPLE RESULT IS >4X SPIKED CONCENTRATION, THEREFORE SPIKE IS NOT DETECTABLE  
\*Y RESULT NOT ATTAINABLE DUE TO SAMPLE MATRIX INTERFERENCE  
@C *VARIABLE MESSAGE*  
@H DETECTION LIMIT ELEVATED DUE TO MATRIX INTERFERENCE  
@Q DETECTION LIMIT ELEVATED DUE TO LIMITED SAMPLE FOR ANALYSIS  
@R RPD LIMIT IS 67% FOR INORGANIC RESULTS LESS THAN TEN TIMES THE REPORTING DETECTION LIMIT  
@S RPD: ONE RESULT ABOVE AND ONE RESULT BELOW REPORTING LIMIT (RL). RESULT ABOVE SHOULD BE < 5 TIMES RL TO BE IN CONTROL.  
@V PRE-DIGEST SPIKE OUT OF LIMITS. POST DIGESTION SPIKE YIELDED ACCEPTABLE RESULTS  
@W DETECTION LIMIT ELEVATED DUE TO REDUCED SAMPLE WEIGHT  
@Y ION BALANCE OUTSIDE OF ATI'S ACCEPTANCE LIMITS; REANALYSIS CONFIRMED ORIGINAL RESULT  
@X RESULTS VERIFIED BY REDIGESTION AND REANALYSIS

ANALYTICAL TECHNOLOGIES, INC.  
SAN DIEGO  
FLAGS

ORGANICS

FLAG	MESSAGE DESCRIPTION
A	A TIC IS A SUSPECTED ALDOL-CONDENSATION PRODUCT
B	ANALYTE FOUND IN THE ASSOCIATED REAGENT BLANK
C	PESTICIDE, WHERE THE IDENTIFICATION WAS CONFIRMED BY GC/MS
CO	THESE COMPOUNDS CO-ELUTE AND ARE QUANTITATED AS ONE PEAK
D	COMPOUND IDENTIFIED IN AN ANALYSIS AT SECONDARY DILUTION
E	ANALYTE AMOUNT EXCEEDS THE CALIBRATION RANGE
J	ESTIMATED VALUE
H	QUANTIFIED AS DIESEL BUT CHROMATOGRAPHIC PATTERN DOES NOT MATCH THAT OF DIESEL
K	QUANTIFIED AS KEROSENE BUT CHROMATOGRAPHIC PATTERN DOES NOT MATCH THAT OF KEROSENE
L	QUANTIFIED AS GASOLINE BUT CHROMATOGRAPHIC PATTERN DOES NOT MATCH THAT OF GASOLINE
N	PRESUMPTIVE EVIDENCE OF A COMPOUND
P	PESTICIDE/AROCLOR TARGET ANALYTE, WHERE THERE IS GREATER THAN 25% DIFFERENCE FOR DETECTED CONCENTRATION BETWEEN 2 GC COLUMNS
TR	COMPOUND DETECTED AT AN UNQUANTIFIABLE TRACE LEVEL
U	COMPOUND WAS ANALYZED FOR BUT NOT DETECTED
X	SEE CASE NARRATIVE
Y	SEE CASE NARRATIVE
Z	SEE CASE NARRATIVE
*	OUTSIDE OF QUALITY CONTROL LIMITS
*D	COMPOUND ANALYZED FROM A SECONDARY ANALYSIS
*F	RESULT OUTSIDE OF ATT'S QUALITY CONTROL LIMITS
*G	RESULT OUTSIDE QUALITY CONTROL LIMITS. INSUFFICIENT SAMPLE FOR RE-EXTRACTION/ANALYSIS
*H	RESULT OUTSIDE OF LIMITS DUE TO SAMPLE MATRIX INTERFERENCE
*I	BECAUSE OF NECESSARY SAMPLE DILUTION, VALUE WAS OUTSIDE QC LIMITS
*K	DUE TO THE NECESSARY DILUTION OF THE SAMPLE, RESULT WAS NOT ATTAINABLE
*L	ANALYTE IS A SUSPECTED LAB CONTAMINANT
*P	A STANDARD WAS USED TO QUANTITATE THIS VALUE
*R	DATA IS NOT USABLE
*T	SURROGATE RECOVERY IS OUTSIDE QC CONTROL LIMITS. NO CORRECTIVE ACTION INDICATED BY METHOD
*V	SAMPLE RESULT IS >4X SPIKED CONCENTRATION, THEREFORE SPIKE IS NOT DETECTABLE
*Y	RESULT NOT ATTAINABLE DUE TO SAMPLE MATRIX INTERFERENCE
@A	RESULTS OUT OF LIMITS DUE TO SAMPLE NON-HOMOGENEITY
@C	VARIABLE MESSAGE
@D	RESULT COULD NOT BE CONFIRMED DUE TO MATRIX INTERFERENCE ON THE CONFIRMATION COLUMN
@E	RESULT MAY BE FALSELY ELEVATED DUE TO SAMPLE MATRIX INTERFERENCE
@F	RESULT OUTSIDE OF CONTRACT SPECIFIED QUALITY CONTROL LIMITS
@G	RESULT OUTSIDE OF CONTRACT SPECIFIED ADVISORY LIMITS
@H	DETECTION LIMIT ELEVATED DUE TO MATRIX INTERFERENCE
@M	RESULT NOT CONFIRMED BY U.V. DUE TO SAMPLE MATRIX INTERFERENCE
@N	RESULT NOT CONFIRMED BY FLUORESCENCE DUE TO SAMPLE MATRIX INTERFERENCE
@P	RESULT QUANTITATED USING FLUORESCENCE ONLY DUE TO THE LOW CONCENTRATION
@Q	DETECTION LIMIT ELEVATED DUE TO LIMITED SAMPLE FOR ANALYSIS
@T	RESULT DUE TO TCLP EXTRACTION MATRIX INTERFERENCE. NO QC LIMITS HAVE BEEN ESTABLISHED
@U	SAMPLE CHROMATOGRAM DOES NOT RESEMBLE COMMON FUEL HYDROCARBON FINGERPRINTS
@Z	SAMPLE CHROMATOGRAM DOES NOT RESEMBLE A FUEL HYDROCARBON

## **APPENDIX B**

### **PULPED MATERIAL PARTICLE SIZING REPORT**

Source: Pulped Material Particle Sizing.  
San Diego, California  
Environmental Testing Associates (ETA), 1994

## NUMERICAL SIZE DISTRIBUTION ANALYSIS (Summary Report)

**Client Name:** NRaD

**Analysis Date :** 9/5/94

**Contact :** Stacey Curtis

**ETA Project # :** 94-4274

**Client Address:** NCCOSC RDTE Division, San Diego, CA 92152

**ETA Sample # :** 4274-1

**Client Project#:** Paper sizing

**Client Sample # :** P2-1 (slides E-H)

**Sample Description :** White paper slurry (final dilution = 0.00008)

**Analysis Requested :** Size distribution analysis

**Analysis Method :** Polarized Light Microscopy

**Magnification(x):** 50

**Scale (µm/division):** 9.90

**Total particles counted:** 100

HYDRODYNAMIC SIZE DISTRIBUTION AND MORPHOLOGY STATISTICS (all paper particles)											
Description	Mean	Std. Dev.	95% C.L.	Description	Mean	Std. Dev.	95% C.L.				
Hydrodynamic Diameter (µm)	183	±239	±47	Fibers / Structure	1.40	±2.15	±0.42				
X-Section Diameter (µm)	330	±520	±102	Paper Fiber Diameter (µm)	21.30	±13.22	±2.59				
Median (µm)	76			Aspect Ratio (all particles)	17.07	±0.19	±0.04				
Mode (size category)	≥31			Structure Sphericity	0.47	±0.18	±0.04				
Skewness	2.6 (positive)			Surface Area/particle (mm <sup>2</sup> )	0.28						
Kurtosis	7.7 (peaked)			Total Surface Area / Volume Ratio	0.01						
HYDRODYNAMIC SIZE DISTRIBUTION (µm ≥ stated size)											
Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	92	63	39	21	9	3		
Individual Count			8	29	24	18	12	6	3		
Individual Numerical %			8.0%	29.0%	24.0%	18.0%	12.0%	6.0%	3.0%		
Cumulative Numerical %			8.0%	37.0%	61.0%	79.0%	91.0%	97.0%	100.0%		
Estimated Volume (Mass Equivalent) Distribution											
Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Individual Volume %			0.0%	0.0%	0.2%	1.8%	7.9%	22.2%	67.8%		
Cumulative Volume %			0.0%	0.0%	0.2%	2.1%	10.0%	32.2%	100.0%		

CROSS-SECTION SIZE DISTRIBUTION (µm ≥ stated size)											
Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	92	64	45	32	20	9	2	
Individual Count			8	28	19	13	12	11	7	2	
Individual Numerical %			8.0%	28.0%	19.0%	13.0%	12.0%	11.0%	7.0%	2.0%	
Cumulative Numerical %			8.0%	36.0%	55.0%	68.0%	80.0%	91.0%	98.0%	100.0%	

Particle Category	Count %	Estimated Volume %	Ave. Hydrodynamic Size (µm)	Ave. X-section Size (µm)	Ave. Aspect Ratio
paper particle	25.0%	0.0%	38	42	2.2
fiber	68.0%	47.3%	197	367	31.6
bundle	4.0%	3.3%	296	500	11.1
matrix	3.0%	49.3%	926	1667	47.7
non-paper					

**Analyst :** \_\_\_\_\_ **Date :** \_\_\_\_ / \_\_\_\_ / \_\_\_\_

## COMPOSITION DISTRIBUTION ANALYSIS (Summary Report)

**Client Name:** NRaD

**Contact :** Stacey Curtis

**Client Address:** NCCOSC RDTE Division, San Diego, CA 92152

**Client Project#:** Paper sizing

**Client Sample # :** P2-1 (slides E-H)

**Sample Description :** White paper slurry (final dilution = 0.00008)

**Analysis Requested :** Size distribution analysis

**Analysis Method :** Polarized Light Microscopy

**Analysis Date :** 9/5/94

**ETA Project # :** 94-4274

**ETA Sample # :** 4274-1

**Magnification(x):** 50

**Scale (µm/div.):** 9.90

**Total particles counted:** 100

Particle Category	Numerical Count	Individual Count % ≥ "Hydrodynamic" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	25			6%	17%	2%						
fiber	68			2%	12%	21%	17%	10%	5%	1%		
bundle	4					1%	1%	1%	1%			
matrix	3							1%		2%		
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ Stated "Hydrodynamic" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	25			6%	23%	25%						
fiber	68			2%	14%	35%	52%	62%	67%	68%		
bundle	4					1%	2%	3%	4%			
matrix	3							1%	1%	3%		
non-paper												

Particle Category	Numerical Count	Individual Count % ≥ "Cross-section" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	25			6%	17%	2%						
fiber	68			2%	11%	16%	12%	12%	9%	5%	1%	
bundle	4					1%	1%		1%	1%		
matrix	3								1%	1%	1%	
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ stated "Cross-section" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	25			6%	23%	25%						
fiber	68			2%	13%	29%	41%	53%	62%	67%	68%	
bundle	4					1%	2%	2%	3%	4%		
matrix	3								1%	2%	3%	
non-paper												

Particle Category	Normalized Count	Individual Hydrodynamic Normalized Count % < maximum stripped size										<2000µm	>4000µm	
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000			
paper particle	25			6%	17%	2%								
fiber	68			2%	12%	21%	17%	10%	5%	1%				
bundle	4					1%	1%	1%	1%					
matrix	3							1%		2%				
non-paper														

Numerical percent of distribution <2000µm = 100%

\* Specific Gravity and thickness to diameter ratios utilized in mass / volume calculations.

Category	paper particle	fiber	bundle	matrix	non-paper
Thickness : diameter ratio	0.80	1.00	0.75	0.50	0.70
Specific Gravity	1.40	1.40	1.40	1.40	1.40

**INDIVIDUAL SIZE DISTRIBUTION COUNT DATA**

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274  
 Sample Description : White paper slurry (final dilution = 0.00008)  
 Analysis Requested : Size distribution analysis  
 Analysis Method : Polarized Light Microscopy  
 Magnification(x): 50  
 Conversion (µm / div.) : 9.90

Client Sample # : P2-1 (slides E-H)  
 ETA Sample # : 4274-1

Total particles counted: 100

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia (µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
1	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
2	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
3	f	1188	20	20	20	1	604	303	60.0	0.26	0.289	0.020
4	b	1782	297	20	134	9	1040	576	6.0	0.32	1.042	0.010
5	f	1485	20	20	20	1	752	352	75.0	0.24	0.390	0.017
6	f	624	30	30	30	1	327	226	21.0	0.36	0.161	0.027
7	f	89	30	30	30	1	59	62	3.0	0.69	0.012	0.097
8	f	693	15	15	15	1	354	192	46.7	0.28	0.116	0.031
9	f	5940	40	40	40	1	2990	1118	150.0	0.19	3.927	0.005
10	f	3119	40	40	40	1	1579	728	78.8	0.23	1.663	0.008
11	f	1188	20	20	20	1	604	303	60.0	0.26	0.289	0.020
12	p	30	15	15	12	1	22	20	2.0	0.68	0.001	0.295
13	p	30	20	20	16	1	25	22	1.5	0.75	0.002	0.268
14	f	990	20	20	20	1	505	269	50.0	0.27	0.227	0.022
15	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
16	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
17	f	2475	40	40	40	1	1257	624	62.5	0.25	1.222	0.010
18	f	248	5	5	5	1	126	67	50.0	0.27	0.014	0.089
19	f	594	20	20	20	1	307	191	30.0	0.32	0.115	0.031
20	p	59	20	20	16	1	40	35	3.0	0.60	0.004	0.169
21	p	30	30	30	24	1	30	26	1.0	0.86	0.002	0.234
22	f	594	20	20	20	1	307	191	30.0	0.32	0.115	0.031
23	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.079
24	f	248	20	20	20	1	134	107	12.5	0.43	0.036	0.056
25	f	50	10	10	10	1	30	29	5.0	0.58	0.003	0.207
26	f	743	20	20	20	1	381	222	37.5	0.30	0.155	0.027
27	m	2970	990	40	416	21	1980	1155	3.0	0.39	4.190	0.005
28	f	69	10	10	10	1	40	36	7.0	0.52	0.004	0.166
29	f	50	10	10	10	1	30	29	5.0	0.58	0.003	0.207
30	f	50	20	20	20	1	35	36	2.5	0.74	0.004	0.165
31	p	119	40	40	32	1	79	71	3.0	0.60	0.016	0.085
32	p	89	20	20	16	1	54	47	4.5	0.52	0.007	0.129
33	f	89	15	15	15	1	52	49	6.0	0.55	0.008	0.122
34	f	228	10	10	10	1	119	80	23.0	0.35	0.020	0.075
35	f	79	20	20	20	1	50	50	4.0	0.63	0.008	0.120
36	p	69	30	30	24	1	50	45	2.3	0.65	0.006	0.133
37	p	79	30	30	24	1	54	49	2.7	0.62	0.008	0.122
38	p	69	30	30	24	1	50	45	2.3	0.65	0.006	0.133
39	f	1188	79	79	79	1	634	482	15.0	0.41	0.729	0.012
40	f	347	79	79	79	1	213	212	4.4	0.61	0.141	0.028

Note: Thickness measurements are based on estimated thickness to diameter ratios for each structure type.

Structure Type Codes			
p	paper particle	m	matrix
f	fiber	n	non-paper
b	bundle		

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P2-1 (slides E-1  
 ETA Sample # : 4274-1

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area Vol. Ratio
41	f	99	30	30	30	1	64	66	3.3	0.67	0.014	0.09
42	f	297	10	10	10	1	153	96	30.0	0.32	0.029	0.06
43	f	149	10	10	10	1	79	60	15.0	0.41	0.011	0.10
44	f	396	30	30	30	1	213	167	13.3	0.42	0.088	0.00
45	f	267	20	20	20	1	144	112	13.5	0.42	0.040	0.05
46	p	20	20	20	16	1	20	17	1.0	0.86	0.001	0.35
47	f	1584	20	20	20	1	802	368	80.0	0.23	0.425	0.01
48	b	1188	59	15	45	4	624	361	20.0	0.30	0.410	0.01
49	f	208	20	20	20	1	114	95	10.5	0.46	0.028	0.06
50	f	842	20	20	20	1	431	241	42.5	0.29	0.183	0.02
51	m	1188	59	30	45	3	624	361	20.0	0.30	0.410	0.01
52	p	30	30	30	24	1	30	26	1.0	0.86	0.002	0.22
53	f	376	20	20	20	1	198	141	19.0	0.37	0.062	0.04
54	p	40	20	20	16	1	30	27	2.0	0.68	0.002	0.22
55	f	99	20	20	20	1	59	58	5.0	0.58	0.011	0.10
56	f	99	20	20	20	1	59	58	5.0	0.58	0.011	0.10
57	f	129	30	30	30	1	79	79	4.3	0.61	0.020	0.07
58	m	4752	40	30	59	4	2396	1262	120.0	0.27	5.007	0.00
59	p	50	40	40	32	1	45	40	1.3	0.80	0.005	0.15
60	f	891	20	20	20	1	455	251	45.0	0.28	0.197	0.02
61	f	149	15	15	15	1	82	69	10.0	0.46	0.015	0.08
62	f	198	10	10	10	1	104	73	20.0	0.37	0.017	0.08
63	f	1485	20	20	20	1	752	352	75.0	0.24	0.390	0.01
64	f	693	20	20	20	1	356	212	35.0	0.31	0.141	0.02
65	f	644	10	10	10	1	327	160	65.0	0.25	0.080	0.02
66	f	297	15	15	15	1	156	109	20.0	0.37	0.038	0.05
67	f	842	20	20	20	1	431	241	42.5	0.29	0.183	0.02
68	p	50	40	40	32	1	45	40	1.3	0.80	0.005	0.15
69	f	495	20	20	20	1	257	169	25.0	0.34	0.090	0.05
70	f	99	20	20	20	1	59	58	5.0	0.58	0.011	0.10
71	p	50	40	40	32	1	45	40	1.3	0.80	0.005	0.15
72	p	79	20	20	16	1	50	43	4.0	0.54	0.006	0.10
73	f	3069	30	30	30	1	1549	654	103.3	0.21	1.344	0.00
74	f	99	8	8	8	1	53	43	12.5	0.43	0.006	0.12
75	f	1218	10	10	10	1	614	245	123.0	0.20	0.188	0.02
76	f	178	8	8	8	1	93	63	22.5	0.35	0.013	0.05
77	p	50	20	20	16	1	35	31	2.5	0.63	0.003	0.10
78	f	347	20	20	20	1	183	133	17.5	0.39	0.056	0.07
79	p	59	20	20	16	1	40	35	3.0	0.60	0.004	0.15
80	p	89	20	20	16	1	54	47	4.5	0.52	0.007	0.12
81	p	50	20	20	16	1	35	31	2.5	0.63	0.003	0.10
82	f	3515	30	30	30	1	1772	716	118.3	0.20	1.610	0.00
83	f	2277	30	30	30	1	1153	536	76.7	0.24	0.903	0.01
84	f	396	10	10	10	1	203	116	40.0	0.29	0.042	0.06
85	f	446	10	10	10	1	228	125	45.0	0.28	0.049	0.06
86	b	396	30	8	18	3	213	119	13.3	0.30	0.044	0.06
87	f	475	20	20	20	1	248	165	24.0	0.35	0.085	0.06

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P2-1 (slides E-H)  
 ETA Sample # : 4274-1

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
88	b	208	40	20	45	3	124	129	5.3	0.62	0.053	0.046
89	p	50	40	40	32	1	45	40	1.3	0.80	0.005	0.152
90	f	89	15	15	15	1	52	49	6.0	0.55	0.008	0.122
91	f	218	10	10	10	1	114	78	22.0	0.36	0.019	0.077
92	p	50	30	30	24	1	40	36	1.7	0.73	0.004	0.167
93	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.079
94	p	50	30	30	24	1	40	36	1.7	0.73	0.004	0.167
95	f	1564	50	50	50	1	807	495	31.6	0.32	0.769	0.012
96	f	99	10	10	10	1	54	46	10.0	0.46	0.007	0.131
97	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
98	p	99	40	40	32	1	69	63	2.5	0.63	0.012	0.095
99	p	50	30	30	24	1	40	36	1.7	0.73	0.004	0.167
100	f	891	30	30	30	1	460	287	30.0	0.32	0.258	0.021

## NUMERICAL SIZE DISTRIBUTION ANALYSIS (Summary Report)

**Client Name:** NRaD

**Analysis Date :** 9/5/94

**Contact :** Stacey Curtis

**ETA Project # :** 94-4274

**Client Address:** NCCOSC RDTE Division, San Diego, CA 92152

**ETA Sample # :** 4274-3

**Client Project#:** Paper sizing

**Client Sample # :** P5-1

**Sample Description :** Brown paper / cardboard

**Analysis Requested :** Size and shape distribution analysis

**Analysis Method :** Polarized Light Microscopy

**Magnification(x):** 50

**Scale (µm/division):** 9.90

**Total particles counted:** 100

### HYDRODYNAMIC SIZE DISTRIBUTION AND MORPHOLOGY STATISTICS (all paper particles)

Description	Mean	Std. Dev.	95% C.L.	Description	Mean	Std. Dev.	95% C.L.
Hydrodynamic Diameter (µm)	192	±245	±48	Fibers / Structure	1.55	±3.74	±0.73
X-Section Diameter (µm)	358	±529	±104	Paper Fiber Diameter (µm)	23.52	±19.90	±3.90
Median (µm)	84			Aspect Ratio (all particles)	18.02	±0.18	±0.04
Mode (size category)	≥31			Structure Sphericity	0.44	±0.18	±0.04
Skewness	2.9 (positive)			Surface Area/particle (mm <sup>2</sup> )	0.30		
Kurtosis	10.9 (peaked)			Total Surface Area / Volume Ratio	0.01		

### HYDRODYNAMIC SIZE DISTRIBUTION (µm ≥ stated size)

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	99	66	37	22	13	1		
Individual Count			1	33	29	15	9	12	1		
Individual Numerical %			1.0%	33.0%	29.0%	15.0%	9.0%	12.0%	1.0%		
Cumulative Numerical %			1.0%	34.0%	63.0%	78.0%	87.0%	99.0%	100.0%		

### Estimated Volume (Mass Equivalent) Distribution

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Individual Volume %			0.0%	0.0%	0.2%	1.2%	4.6%	44.4%	49.5%		
Cumulative Volume %			0.0%	0.0%	0.3%	1.5%	6.1%	50.5%	100.0%		

### CROSS-SECTION SIZE DISTRIBUTION (µm ≥ stated size)

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	99	74	46	33	19	14	2	
Individual Count			1	25	28	13	14	5	12	2	
Individual Numerical %			1.0%	25.0%	28.0%	13.0%	14.0%	5.0%	12.0%	2.0%	
Cumulative Numerical %			1.0%	26.0%	54.0%	67.0%	81.0%	86.0%	98.0%	100.0%	

Particle Category	Count %	Estimated Volume %	Ave. Hydrodynamic Size (µm)	Ave. X-section Size (µm)	Ave. Aspect Ratio
paper particle	27.0%	0.2%	61	69	2.6
fiber	67.0%	42.4%	213	433	42.8
bundle	1.0%	4.9%	740	1460	13.8
matrix	5.0%	52.6%	497	708	5.6
non-paper					

**Analyst :** \_\_\_\_\_ **Date :** \_\_\_\_ / \_\_\_\_ / \_\_\_\_

**COMPOSITION DISTRIBUTION ANALYSIS  
(Summary Report)**

**Client Name:** NRaD

**Analysis Date :** 9/5/94

**Contact :** Stacey Curtis

**ETA Project # :** 94-4274

**Client Address:** NCCOSC RDTE Division, San Diego, CA 92152

**ETA Sample # :** 4274-3

**Client Project#:** Paper sizing

**Client Sample # :** P5-1

**Sample Description :** Brown paper / cardboard

**Magnification(x):** 50

**Analysis Requested :** Size and shape distribution analysis

**Scale (µm/div.):** 9.90

**Analysis Method :** Polarized Light Microscopy

**Total particles counted:** 100

Particle Category	Numerical Count	Individual Count % ≥ "Hydrodynamic" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	27				20%	5%	2%					
fiber	67			1%	11%	24%	12%	9%	10%			
bundle	1								1%			
matrix	5				2%		1%		1%	1%		
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ Stated "Hydrodynamic" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	27				20%	25%	27%					
fiber	67			1%	12%	36%	48%	57%	67%			
bundle	1								1%			
matrix	5				2%	2%	3%	3%	4%	5%		
non-paper												

Particle Category	Numerical Count	Individual Count % ≥ "Cross-section" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	27				19%	6%	2%					
fiber	67			1%	6%	20%	11%	13%	5%	10%	1%	
bundle	1									1%		
matrix	5					2%		1%		1%	1%	
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ stated "Cross-section" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	27				19%	25%	27%					
fiber	67			1%	7%	27%	38%	51%	56%	66%	67%	
bundle	1									1%		
matrix	5					2%	2%	3%	3%	4%	5%	
non-paper												

Particle Category	Normalized Count	Individual Hydrodynamic Normalized Count % < maximum stripped size										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	<2000µm ≥4000
paper particle	27				20%	5%	2%					
fiber	67			1%	11%	24%	12%	9%	10%			
bundle	1								1%			
matrix	5				2%		1%		1%	1%		
non-paper												

*Numerical percent of distribution <2000µm = 100%*

\* Specific Gravity and thickness to diameter ratios utilized in mass / volume calculations.

Category	paper particle	fiber	bundle	matrix	non-paper
Thickness : diameter ratio	0.80	1.00	0.75	0.50	0.70
Specific Gravity	1.40	1.40	1.40	1.40	1.40

**INDIVIDUAL SIZE DISTRIBUTION COUNT DATA**

Client Name: NRaD

Client Project#: Paper sizing

ETA Project # : 94-4274

Client Sample # : P5-1

ETA Sample # : 4274-3

Sample Description : Brown paper / cardboard

Analysis Requested : Size and shape distribution analysis

Analysis Method : Polarized Light Microscopy

Magnification(x): 50

Total particles counted: 100

Conversion (µm / div.) : 9.90

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia (µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
1	m	3218	842	30	579	39	2030	1604	3.8	0.50	8.082	0.004
2	f	2376	30	30	30	1	1203	551	80.0	0.23	0.955	0.011
3	f	238	5	5	5	1	121	65	48.0	0.28	0.013	0.092
4	f	248	10	10	10	1	129	85	25.0	0.34	0.023	0.071
5	f	158	10	10	10	1	84	63	16.0	0.40	0.012	0.095
6	f	178	5	5	5	1	92	54	36.0	0.30	0.009	0.111
7	f	89	10	10	10	1	50	43	9.0	0.48	0.006	0.140
8	f	743	20	20	20	1	381	222	37.5	0.30	0.155	0.027
9	f	2376	30	30	30	1	1203	551	80.0	0.23	0.955	0.011
10	f	2079	30	30	30	1	1054	504	70.0	0.24	0.799	0.012
11	p	69	30	30	24	1	50	45	2.3	0.65	0.006	0.133
12	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.079
13	f	277	10	10	10	1	144	91	28.0	0.33	0.026	0.066
14	f	69	8	8	8	1	39	34	8.8	0.49	0.004	0.178
15	f	257	5	5	5	1	131	69	52.0	0.27	0.015	0.087
16	f	792	20	20	20	1	406	232	40.0	0.29	0.168	0.026
17	p	79	40	40	32	1	59	54	2.0	0.68	0.009	0.111
18	m	119	50	10	15	3	84	40	2.4	0.33	0.005	0.151
19	f	743	30	30	30	1	386	254	25.0	0.34	0.203	0.024
20	p	99	30	30	24	1	64	57	3.3	0.58	0.010	0.105
21	f	178	20	20	20	1	99	86	9.0	0.48	0.023	0.070
22	f	941	10	10	10	1	475	206	95.0	0.22	0.133	0.029
23	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
24	f	50	5	5	5	1	27	23	10.0	0.46	0.002	0.261
25	f	198	10	10	10	1	104	73	20.0	0.37	0.017	0.082
26	b	2723	198	20	104	7	1460	740	13.8	0.27	1.718	0.008
27	f	1386	20	20	20	1	703	336	70.0	0.24	0.355	0.018
28	p	59	59	59	48	1	59	51	1.0	0.86	0.008	0.117
29	f	842	30	30	30	1	436	276	28.3	0.33	0.239	0.022
30	f	99	10	10	10	1	54	46	10.0	0.46	0.007	0.131
31	p	119	30	30	24	1	74	64	4.0	0.54	0.013	0.093
32	m	1931	149	50	99	4	1040	627	13.0	0.32	1.233	0.010
33	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.079
34	p	89	30	30	24	1	59	53	3.0	0.60	0.009	0.113
35	p	79	30	30	24	1	54	49	2.7	0.62	0.008	0.122
36	p	79	30	30	24	1	54	49	2.7	0.62	0.008	0.122
37	p	59	50	50	40	1	54	48	1.2	0.81	0.007	0.125
38	p	50	50	50	40	1	50	43	1.0	0.86	0.006	0.141
39	f	2129	20	20	20	1	1074	448	107.5	0.21	0.630	0.013
40	p	99	20	20	16	1	59	50	5.0	0.50	0.008	0.120

Note: Thickness measurements are based on estimated thickness to diameter ratios for each structure type.

Structure Type Codes			
p	paper particle	m	matrix
f	fiber	n	non-paper
b	bundle		

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P5-1  
 ETA Sample # : 4274-3

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area Vol. Rat
41	p	69	20	20	16	1	45	39	3.5	0.57	0.005	0.1
42	f	149	10	10	10	1	79	60	15.0	0.41	0.011	0.1
43	p	69	50	50	40	1	59	53	1.4	0.77	0.009	0.1
44	f	149	30	30	30	1	89	87	5.0	0.58	0.024	0.0
45	f	495	30	30	30	1	262	194	16.7	0.39	0.118	0.0
46	f	178	20	20	20	1	99	86	9.0	0.48	0.023	0.0
47	f	644	20	20	20	1	332	202	32.5	0.31	0.128	0.0
48	f	3762	20	20	20	1	1891	654	190.0	0.17	1.345	0.0
49	f	495	10	10	10	1	252	134	50.0	0.27	0.057	0.0
50	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.0
51	p	99	40	40	32	1	69	63	2.5	0.63	0.012	0.0
52	p	297	149	149	119	1	223	203	2.0	0.68	0.130	0.0
53	p	69	50	50	40	1	59	53	1.4	0.77	0.009	0.1
54	f	69	10	10	10	1	40	36	7.0	0.52	0.004	0.1
55	f	198	5	5	5	1	101	58	40.0	0.29	0.011	0.1
56	f	3416	30	30	30	1	1723	702	115.0	0.21	1.550	0.0
57	f	891	20	20	20	1	455	251	45.0	0.28	0.197	0.0
58	f	198	20	20	20	1	109	92	10.0	0.46	0.027	0.0
59	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.0
60	f	3069	20	20	20	1	1544	571	155.0	0.19	1.026	0.0
61	p	89	59	59	48	1	74	67	1.5	0.75	0.014	0.0
62	f	3049	30	30	30	1	1539	651	102.7	0.21	1.332	0.0
63	f	594	20	20	20	1	307	191	30.0	0.32	0.115	0.0
64	f	416	10	10	10	1	213	120	42.0	0.29	0.045	0.0
65	f	267	20	20	20	1	144	112	13.5	0.42	0.040	0.0
66	p	59	40	40	32	1	50	45	1.5	0.75	0.006	0.1
67	f	1436	10	10	10	1	723	273	145.0	0.19	0.235	0.0
68	f	495	30	30	30	1	262	194	16.7	0.39	0.118	0.0
69	p	99	79	79	63	1	89	79	1.3	0.80	0.020	0.0
70	m	545	79	20	40	4	312	180	6.9	0.33	0.102	0.0
71	f	2525	30	30	30	1	1277	574	85.0	0.23	1.036	0.0
72	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.0
73	f	327	20	20	20	1	173	128	16.5	0.39	0.052	0.0
74	p	89	20	20	16	1	54	47	4.5	0.52	0.007	0.1
75	p	149	20	20	16	1	84	65	7.5	0.44	0.013	0.0
76	p	79	30	30	24	1	54	49	2.7	0.62	0.008	0.1
77	f	1752	20	20	20	1	886	393	88.5	0.22	0.486	0.0
78	p	59	50	50	40	1	54	48	1.2	0.81	0.007	0.1
79	f	178	30	30	30	1	104	98	6.0	0.55	0.030	0.0
80	p	238	79	79	63	1	158	142	3.0	0.60	0.063	0.0
81	f	990	25	25	25	1	507	289	40.0	0.29	0.263	0.0
82	f	347	5	5	5	1	176	84	70.0	0.24	0.022	0.0
83	f	743	20	20	20	1	381	222	37.5	0.30	0.155	0.0
84	f	79	10	10	10	1	45	40	8.0	0.50	0.005	0.1
85	f	218	5	5	5	1	111	62	44.0	0.28	0.012	0.0
86	m	99	50	8	16	4	74	37	2.0	0.37	0.004	0.1
87	f	5643	30	30	30	1	2836	982	190.0	0.17	3.027	0.0

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P5-1  
 ETA Sample # : 4274-3

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
88	f	79	5	5	5	1	42	31	16.0	0.40	0.003	0.191
89	p	50	40	40	32	1	45	40	1.3	0.80	0.005	0.152
90	p	79	20	20	16	1	50	43	4.0	0.54	0.006	0.140
91	f	218	20	20	20	1	119	98	11.0	0.45	0.030	0.061
92	f	317	20	20	20	1	168	126	16.0	0.40	0.050	0.048
93	f	3000	40	40	40	1	1520	709	75.8	0.24	1.579	0.008
94	f	396	5	5	5	1	200	92	80.0	0.23	0.027	0.065
95	f	297	20	20	20	1	158	120	15.0	0.41	0.046	0.050
96	f	109	20	20	20	1	64	62	5.5	0.57	0.012	0.097
97	f	1386	50	50	50	1	718	456	28.0	0.33	0.655	0.013
98	f	297	20	20	20	1	158	120	15.0	0.41	0.046	0.050
99	f	614	20	20	20	1	317	195	31.0	0.32	0.120	0.031
100	p	59	30	30	24	1	45	41	2.0	0.68	0.005	0.148

## NUMERICAL SIZE DISTRIBUTION ANALYSIS (Summary Report)

**Client Name:** NRaD

**Analysis Date :** 9/5/94

**Contact :** Stacey Curtis

**ETA Project # :** 94-4274

**Client Address:** NCCOSC RDTE Division, San Diego, CA 92152

**ETA Sample # :** 4274-2

**Client Project#:** Paper sizing

**Client Sample # :** P6-1

**Sample Description :** White paper slurry

**Analysis Requested :** Size and shape distribution analysis

**Analysis Method :** Polarized Light Microscopy

**Magnification(x):** 50

**Scale (µm/division):** 9.90

**Total particles counted:** 100

### HYDRODYNAMIC SIZE DISTRIBUTION AND MORPHOLOGY STATISTICS (all paper particles)

Description	Mean	Std. Dev.	95% C.L.	Description	Mean	Std. Dev.	95% C.L.
Hydrodynamic Diameter (µm)	130	±179	±35	Fibers / Structure	1.38	±1.79	±0.35
X-Section Diameter (µm)	227	±402	±79	Paper Fiber Diameter (µm)	19.30	±13.57	±2.66
Median (µm)	71			Aspect Ratio (all particles)	13.76	±0.15	±0.03
Mode (size category)	≥31			Structure Sphericity	0.46	±0.17	±0.03
Skewness	4.4 (positive)			Surface Area/particle (mm <sup>2</sup> )	0.15		
Kurtosis	25.9 (peaked)			Total Surface Area / Volume Ratio	0.01		

### HYDRODYNAMIC SIZE DISTRIBUTION (µm ≥ stated size)

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	93	54	29	13	3	1		
Individual Count			7	39	25	16	10	2	1		
Individual Numerical %			7.0%	39.0%	25.0%	16.0%	10.0%	2.0%	1.0%		
Cumulative Numerical %			7.0%	46.0%	71.0%	87.0%	97.0%	99.0%	100.0%		

### Estimated Volume (Mass Equivalent) Distribution

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Individual Volume %			0.0%	0.1%	0.4%	2.5%	10.4%	18.7%	67.9%		
Cumulative Volume %			0.0%	0.1%	0.5%	3.0%	13.4%	32.1%	100.0%		

### CROSS-SECTION SIZE DISTRIBUTION (µm ≥ stated size)

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	96	59	39	20	12	3	1	
Individual Count			4	37	20	19	8	9	2	1	
Individual Numerical %			4.0%	37.0%	20.0%	19.0%	8.0%	9.0%	2.0%	1.0%	
Cumulative Numerical %			4.0%	41.0%	61.0%	80.0%	88.0%	97.0%	99.0%	100.0%	

Particle Category	Count %	Estimated Volume %	Ave. Hydrodynamic Size (µm)	Ave. X-section Size (µm)	Ave. Aspect Ratio
paper particle	28.0%	4.7%	62	116	9.4
fiber	66.0%	27.1%	140	233	27.5
bundle	1.0%	0.0%	79	218	10.0
matrix	5.0%	68.2%	379	770	5.4
non-paper					

Analyst : \_\_\_\_\_ Date : \_\_\_\_ / \_\_\_\_ / \_\_\_\_

## COMPOSITION DISTRIBUTION ANALYSIS (Summary Report)

**Client Name:** NRaD

**Contact :** Stacey Curtis

**Client Address:** NCCOSC RDTE Division, San Diego, CA 92152

**Client Project#:** Paper sizing

**Client Sample # :** P6-1

**Sample Description :** White paper slurry

**Analysis Requested :** Size and shape distribution analysis

**Analysis Method :** Polarized Light Microscopy

**Analysis Date :** 9/5/94

**ETA Project # :** 94-4274

**ETA Sample # :** 4274-2

**Magnification(x):** 50

**Scale (µm/div.):** 9.90

**Total particles counted:** 100

Particle Category	Numerical Count	Individual Count % ≥ "Hydrodynamic" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	28			4%	21%	2%			1%			
fiber	66			3%	18%	19%	15%	10%	1%			
bundle	1					1%						
matrix	5					3%	1%			1%		
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ Stated "Hydrodynamic" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	28			4%	25%	27%	27%	27%	28%			
fiber	66			3%	21%	40%	55%	65%	66%			
bundle	1					1%						
matrix	5					3%	4%	4%	4%	5%		
non-paper												

Particle Category	Numerical Count	Individual Count % ≥ "Cross-section" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	28			2%	22%	3%				1%		
fiber	66			2%	15%	17%	15%	7%	9%	1%		
bundle	1						1%					
matrix	5						3%	1%			1%	
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ stated "Cross-section" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	28			2%	24%	27%	27%	27%	27%	28%		
fiber	66			2%	17%	34%	49%	56%	65%	66%		
bundle	1						1%					
matrix	5						3%	4%	4%	4%	5%	
non-paper												

Particle Category	Normalized Count	Individual Hydrodynamic Normalized Count % < maximum stripped size										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	<2000µm
paper particle	28			4%	21%	2%			1%			
fiber	66			3%	18%	19%	15%	10%	1%			
bundle	1					1%						
matrix	5					3%	1%			1%		
non-paper												

*Numerical percent of distribution <2000µm = 100%*

\* Specific Gravity and thickness to diameter ratios utilized in mass / volume calculations.

Category	paper particle	fiber	bundle	matrix	non-paper
Thickness : diameter ratio	0.80	1.00	0.75	0.50	0.70
Specific Gravity	0.50	0.50	0.50	0.50	0.50

**INDIVIDUAL SIZE DISTRIBUTION COUNT DATA**

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274  
 Sample Description : White paper slurry  
 Analysis Requested : Size and shape distribution analysis  
 Analysis Method : Polarized Light Microscopy  
 Magnification(x): 50  
 Conversion (µm / div.): 9.90

Client Sample # : P6-1  
 ETA Sample # : 4274-2

Total particles counted: 100

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
1	f	30	10	10	10	1	20	21	3.0	0.69	0.001	0.291
2	f	594	30	30	30	1	312	219	20.0	0.37	0.150	0.027
3	p	40	10	10	8	1	25	21	4.0	0.54	0.001	0.279
4	f	396	5	5	5	1	200	92	80.0	0.23	0.027	0.065
5	f	79	5	5	5	1	42	31	16.0	0.40	0.003	0.191
6	f	139	20	20	20	1	79	72	7.0	0.52	0.016	0.083
7	f	79	10	10	10	1	45	40	8.0	0.50	0.005	0.152
8	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.079
9	f	376	30	30	30	1	203	161	12.7	0.43	0.082	0.037
10	f	99	10	10	10	1	54	46	10.0	0.46	0.007	0.131
11	p	59	30	30	24	1	45	41	2.0	0.68	0.005	0.148
12	m	327	149	30	119	8	238	216	2.2	0.66	0.147	0.028
13	f	297	10	10	10	1	153	96	30.0	0.32	0.029	0.063
14	p	89	30	30	24	1	59	53	3.0	0.60	0.009	0.113
15	p	40	30	30	24	1	35	31	1.3	0.78	0.003	0.194
16	m	297	248	8	36	9	272	77	1.2	0.26	0.019	0.078
17	f	792	20	20	20	1	406	232	40.0	0.29	0.168	0.026
18	f	990	15	15	15	1	502	244	66.7	0.25	0.187	0.025
19	p	59	30	30	24	1	45	41	2.0	0.68	0.005	0.148
20	p	79	40	40	32	1	59	54	2.0	0.68	0.009	0.111
21	p	3812	20	20	16	1	1916	569	192.5	0.15	1.017	0.011
22	f	50	5	5	5	1	27	23	10.0	0.46	0.002	0.261
23	f	99	5	5	5	1	52	36	20.0	0.37	0.004	0.165
24	f	238	8	8	8	1	123	76	30.0	0.32	0.018	0.078
25	f	297	40	40	40	1	168	152	7.5	0.51	0.072	0.040
26	f	238	40	40	40	1	139	131	6.0	0.55	0.054	0.046
27	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
28	p	30	30	30	24	1	30	26	1.0	0.86	0.002	0.234
29	p	89	20	20	16	1	54	47	4.5	0.52	0.007	0.129
30	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.079
31	p	40	30	30	24	1	35	31	1.3	0.78	0.003	0.194
32	p	50	40	40	32	1	45	40	1.3	0.80	0.005	0.152
33	f	198	10	10	10	1	104	73	20.0	0.37	0.017	0.082
34	f	1485	15	15	15	1	750	320	100.0	0.22	0.322	0.019
35	f	218	10	10	10	1	114	78	22.0	0.36	0.019	0.077
36	f	178	5	5	5	1	92	54	36.0	0.30	0.009	0.111
37	b	396	40	5	11	3	218	79	10.0	0.20	0.020	0.076
38	p	99	79	79	63	1	89	79	1.3	0.80	0.020	0.076
39	f	396	50	50	50	1	223	198	8.0	0.50	0.123	0.030
40	f	89	5	5	5	1	47	34	18.0	0.38	0.004	0.176

Note: Thickness measurements are based on estimated thickness to diameter ratios for each structure type.

Structure Type Codes			
p	paper particle	m	matrix
f	fiber	n	non-paper
b	bundle		

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P6-1  
 ETA Sample # : 4274-2

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area Vol. Ratio
41	f	3762	40	40	40	1	1901	824	95.0	0.22	2.136	0.00
42	f	396	20	20	20	1	208	146	20.0	0.37	0.067	0.04
43	f	198	10	10	10	1	104	73	20.0	0.37	0.017	0.09
44	f	1683	20	20	20	1	851	383	85.0	0.23	0.460	0.01
45	p	59	20	20	16	1	40	35	3.0	0.60	0.004	0.10
46	f	139	5	5	5	1	72	46	28.0	0.33	0.007	0.10
47	f	1049	20	20	20	1	535	279	53.0	0.27	0.245	0.02
48	f	99	5	5	5	1	52	36	20.0	0.37	0.004	0.10
49	f	743	5	5	5	1	374	140	150.0	0.19	0.061	0.04
50	p	40	40	40	32	1	40	34	1.0	0.86	0.004	0.17
51	m	5198	495	30	223	15	2846	1394	10.5	0.27	6.103	0.00
52	p	79	30	30	24	1	54	49	2.7	0.62	0.008	0.10
53	p	99	20	20	16	1	59	50	5.0	0.50	0.008	0.10
54	f	89	15	15	15	1	52	49	6.0	0.55	0.008	0.10
55	f	347	30	30	30	1	188	153	11.7	0.44	0.073	0.00
56	f	1337	30	30	30	1	683	376	45.0	0.28	0.444	0.01
57	f	792	40	40	40	1	416	292	20.0	0.37	0.267	0.02
58	f	99	8	8	8	1	53	43	12.5	0.43	0.006	0.10
59	f	99	10	10	10	1	54	46	10.0	0.46	0.007	0.10
60	f	69	10	10	10	1	40	36	7.0	0.52	0.004	0.10
61	p	89	30	30	24	1	59	53	3.0	0.60	0.009	0.11
62	f	99	20	20	20	1	59	58	5.0	0.58	0.011	0.10
63	f	1238	20	20	20	1	629	312	62.5	0.25	0.305	0.00
64	f	149	30	30	30	1	89	87	5.0	0.58	0.024	0.00
65	p	89	20	20	16	1	54	47	4.5	0.52	0.007	0.10
66	f	792	30	30	30	1	411	265	26.7	0.33	0.221	0.00
67	f	1089	40	40	40	1	564	361	27.5	0.33	0.409	0.00
68	f	594	20	20	20	1	307	191	30.0	0.32	0.115	0.00
69	f	99	10	10	10	1	54	46	10.0	0.46	0.007	0.10
70	f	297	5	5	5	1	151	76	60.0	0.26	0.018	0.00
71	f	99	20	20	20	1	59	58	5.0	0.58	0.011	0.10
72	f	79	5	5	5	1	42	31	16.0	0.40	0.003	0.10
73	p	89	20	20	16	1	54	47	4.5	0.52	0.007	0.10
74	f	743	20	20	20	1	381	222	37.5	0.30	0.155	0.00
75	f	129	20	20	20	1	74	69	6.5	0.54	0.015	0.00
76	f	89	3	3	3	1	46	29	30.0	0.32	0.003	0.20
77	p	50	20	20	16	1	35	31	2.5	0.63	0.003	0.10
78	p	50	30	30	24	1	40	36	1.7	0.73	0.004	0.10
79	f	248	20	20	20	1	134	107	12.5	0.43	0.036	0.00
80	p	50	20	20	16	1	35	31	2.5	0.63	0.003	0.10
81	f	248	3	3	3	1	125	57	83.3	0.23	0.010	0.10
82	f	208	5	5	5	1	106	60	42.0	0.29	0.011	0.10
83	f	337	20	20	20	1	178	131	17.0	0.39	0.054	0.00
84	p	99	30	30	24	1	64	57	3.3	0.58	0.010	0.10
85	f	297	15	15	15	1	156	109	20.0	0.37	0.038	0.00
86	f	208	20	20	20	1	114	95	10.5	0.46	0.028	0.00
87	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.00

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P6-1  
 ETA Sample # : 4274-2

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
88	p	99	50	50	40	1	74	68	2.0	0.68	0.014	0.089
89	p	79	30	30	24	1	54	49	2.7	0.62	0.008	0.122
90	m	396	99	10	30	6	248	112	4.0	0.28	0.039	0.054
91	m	446	50	10	15	3	248	96	9.0	0.22	0.029	0.063
92	f	396	20	20	20	1	208	146	20.0	0.37	0.067	0.041
93	f	416	20	20	20	1	218	151	21.0	0.36	0.071	0.040
94	f	1337	50	50	50	1	693	446	27.0	0.33	0.624	0.013
95	p	69	30	30	24	1	50	45	2.3	0.65	0.006	0.133
96	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
97	f	1188	30	30	30	1	609	347	40.0	0.29	0.379	0.017
98	f	208	20	20	20	1	114	95	10.5	0.46	0.028	0.063
99	p	79	20	20	16	1	50	43	4.0	0.54	0.006	0.140
100	p	59	20	20	16	1	40	35	3.0	0.60	0.004	0.169

## COMPOSITION DISTRIBUTION ANALYSIS (Summary Report)

**Client Name:** NRaD

**Analysis Date :** 9/5/94

**Contact :** Stacey Curtis

**ETA Project # :** 94-4274

**Client Address:** NCCOSC RDTE Division, San Diego, CA 92152

**ETA Sample # :** 4274-4

**Client Project#:** Paper sizing

**Client Sample # :** P8-1

**Sample Description :** Mixed paper

**Magnification(x):** 50

**Analysis Requested :** Size and shape distribution analysis

**Scale (µm/div.):** 9.90

**Analysis Method :** Polarized Light Microscopy

**Total particles counted:** 100

Particle Category	Numerical Count	Individual Count % ≥ "Hydrodynamic" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	26				22%	3%	1%					
fiber	68			1%	11%	21%	19%	12%	4%			
bundle	2					1%	1%					
matrix	4				1%	1%			1%	1%		
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ Stated "Hydrodynamic" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	26				22%	25%	26%					
fiber	68			1%	12%	33%	52%	64%	68%			
bundle	2					1%	2%					
matrix	4				1%	2%	2%	2%	3%	4%		
non-paper												

Particle Category	Numerical Count	Individual Count % ≥ "Cross-section" Stated Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	26				20%	5%			1%			
fiber	68				5%	23%	8%	18%	10%	4%		
bundle	2						2%					
matrix	4					1%	1%				2%	
non-paper												

Particle Category	Numerical Count	Cumulative Count % ≥ stated "Cross-section" Size(µm)										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	26				20%	25%	25%	25%	26%			
fiber	68				5%	28%	36%	54%	64%	68%		
bundle	2						2%					
matrix	4					1%	2%	2%	2%	2%	4%	
non-paper												

Particle Category	Normalized Count	Individual Hydrodynamic Normalized Count % < maximum stripped size <2000µm										
		<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
paper particle	26				22%	3%	1%					
fiber	68			1%	11%	21%	19%	12%	4%			
bundle	2					1%	1%					
matrix	4				1%	1%			1%	1%		
non-paper												

Numerical percent of distribution <2000µm = 100%

\* Specific Gravity and thickness to diameter ratios utilized in mass / volume calculations.

Category	paper particle	fiber	bundle	matrix	non-paper
Thickness : diameter ratio	0.80	1.00	0.75	0.50	0.70
Specific Gravity	0.50	0.50	0.50	0.50	0.50

## NUMERICAL SIZE DISTRIBUTION ANALYSIS (Summary Report)

<b>Client Name:</b> NRaD	<b>Analysis Date :</b> 9/5/94
<b>Contact :</b> Stacey Curtis	<b>ETA Project # :</b> 94-4274
<b>Client Address:</b> NCCOSC RDTE Division, San Diego, CA 92152	<b>ETA Sample # :</b> 4274-4
<b>Client Project#:</b> Paper sizing	
<b>Client Sample # :</b> P8-1	
<b>Sample Description :</b> Mixed paper	
<b>Analysis Requested :</b> Size and shape distribution analysis	
<b>Analysis Method :</b> Polarized Light Microscopy	
<b>Magnification(x):</b> 50	
<b>Scale (µm/division):</b> 9.90	
<b>Total particles counted:</b> 100	

### HYDRODYNAMIC SIZE DISTRIBUTION AND MORPHOLOGY STATISTICS (all paper particles)

Description	Mean	Std. Dev.	95% C.L.	Description	Mean	Std. Dev.	95% C.L.
Hydrodynamic Diameter (µm)	175	±232	±46	Fibers / Structure	1.65	±4.45	±0.87
X-Section Diameter (µm)	305	±455	±89	Paper Fiber Diameter (µm)	22.23	±15.99	±3.13
Median (µm)	87			Aspect Ratio (all particles)	13.21	±0.16	±0.03
Mode (size category)	≥31			Structure Sphericity	0.44	±0.16	±0.03
Skewness	4.5 (positive)			Surface Area/particle (mm <sup>2</sup> )	0.27		
Kurtosis	28.1 (peaked)			Total Surface Area / Volume Ratio	0.01		

### HYDRODYNAMIC SIZE DISTRIBUTION (µm ≥ stated size)

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	99	65	39	18	6	1		
Individual Count			1	34	26	21	12	5	1		
Individual Numerical %			1.0%	34.0%	26.0%	21.0%	12.0%	5.0%	1.0%		
Cumulative Numerical %			1.0%	35.0%	61.0%	82.0%	94.0%	99.0%	100.0%		

### Estimated Volume (Mass Equivalent) Distribution

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Individual Volume %			0.0%	0.0%	0.2%	2.1%	5.7%	19.4%	72.6%		
Cumulative Volume %			0.0%	0.0%	0.2%	2.3%	8.0%	27.4%	100.0%		

### CROSS-SECTION SIZE DISTRIBUTION (µm ≥ stated size)

Particle Size (µm)	<8	≥8	≥16	≥31	≥63	≥125	≥250	≥500	≥1000	≥2000	≥4000
Midpoint size (µm)	6	12	23	47	94	188	375	750	1500	3000	≥4000
Numerical Count ≥	100	100	100	100	75	46	35	17	6	2	
Individual Count				25	29	11	18	11	4	2	
Individual Numerical %				25.0%	29.0%	11.0%	18.0%	11.0%	4.0%	2.0%	
Cumulative Numerical %				25.0%	54.0%	65.0%	83.0%	94.0%	98.0%	100.0%	

Particle Category	Count %	Estimated Volume %	Ave. Hydrodynamic Size (µm)	Ave. X-section Size (µm)	Ave. Aspect Ratio
paper particle	26.0%	0.2%	56	74	4.6
fiber	68.0%	23.1%	191	338	32.7
bundle	2.0%	0.1%	136	196	8.0
matrix	4.0%	76.6%	680	1297	4.2
non-paper					

Analyst : \_\_\_\_\_ Date : \_\_\_\_ / \_\_\_\_ / \_\_\_\_

**INDIVIDUAL SIZE DISTRIBUTION COUNT DATA**

Client Name: NRaD

Client Project#: Paper sizing

ETA Project # : 94-4274

Sample Description : Mixed paper

Analysis Requested : Size and shape distribution analysis

Analysis Method : Polarized Light Microscopy

Magnification(x): 50

Client Sample # : P8-1

ETA Sample # : 4274-4

Total particles counted: 100

Conversion (µm / div.) : 9.90

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
1	f	277	20	20	20	1	149	115	14.0	0.41	0.042	0.052
2	p	69	20	20	16	1	45	39	3.5	0.57	0.005	0.153
3	p	1089	20	20	16	1	554	247	55.0	0.23	0.191	0.024
4	f	792	20	20	20	1	406	232	40.0	0.29	0.168	0.026
5	p	59	30	30	24	1	45	41	2.0	0.68	0.005	0.148
6	f	515	20	20	20	1	267	174	26.0	0.34	0.095	0.035
7	p	79	40	40	32	1	59	54	2.0	0.68	0.009	0.111
8	f	69	10	10	10	1	40	36	7.0	0.52	0.004	0.166
9	m	4455	990	25	569	46	2723	1866	4.5	0.42	10.938	0.003
10	f	178	10	10	10	1	94	68	18.0	0.38	0.015	0.088
11	f	673	20	20	20	1	347	208	34.0	0.31	0.136	0.029
12	f	743	20	20	20	1	381	222	37.5	0.30	0.155	0.027
13	p	50	30	30	24	1	40	36	1.7	0.73	0.004	0.167
14	p	99	30	30	24	1	64	57	3.3	0.58	0.010	0.105
15	f	277	20	20	20	1	149	115	14.0	0.41	0.042	0.052
16	p	59	20	20	16	1	40	35	3.0	0.60	0.004	0.169
17	f	941	40	40	40	1	490	327	23.8	0.35	0.336	0.018
18	f	109	20	20	20	1	64	62	5.5	0.57	0.012	0.097
19	f	248	10	10	10	1	129	85	25.0	0.34	0.023	0.071
20	f	644	15	15	15	1	329	183	43.3	0.28	0.105	0.033
21	f	614	20	20	20	1	317	195	31.0	0.32	0.120	0.031
22	f	1109	59	59	59	1	584	418	18.7	0.38	0.549	0.014
23	f	693	40	40	40	1	366	267	17.5	0.39	0.224	0.022
24	p	59	30	30	24	1	45	41	2.0	0.68	0.005	0.148
25	f	178	30	30	30	1	104	98	6.0	0.55	0.030	0.061
26	m	3564	792	30	149	10	2178	707	4.5	0.20	1.571	0.008
27	p	89	25	25	20	1	57	50	3.6	0.56	0.008	0.120
28	f	158	20	20	20	1	89	79	8.0	0.50	0.020	0.076
29	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.092
30	f	198	20	20	20	1	109	92	10.0	0.46	0.027	0.065
31	p	50	40	40	32	1	45	40	1.3	0.80	0.005	0.152
32	f	218	15	15	15	1	116	89	14.7	0.41	0.025	0.067
33	f	970	20	20	20	1	495	265	49.0	0.27	0.221	0.023
34	m	277	59	20	30	3	168	104	4.7	0.38	0.034	0.057
35	f	792	20	20	20	1	406	232	40.0	0.29	0.168	0.026
36	p	50	50	50	40	1	50	43	1.0	0.86	0.006	0.141
37	f	3861	40	40	40	1	1950	839	97.5	0.22	2.211	0.007
38	f	79	10	10	10	1	45	40	8.0	0.50	0.005	0.152
39	f	178	20	20	20	1	99	86	9.0	0.48	0.023	0.070
40	f	1782	20	20	20	1	901	398	90.0	0.22	0.497	0.015

Note: Thickness measurements are based on estimated thickness to diameter ratios for each structure type.

Structure Type Codes			
p	paper particle	m	matrix
f	fiber	n	non-paper
b	bundle		

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P8-1  
 ETA Sample # : 4274-4

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area. Vol. Ratio
41	f	2059	30	30	30	1	1044	501	69.3	0.24	0.789	0.01
42	f	1634	35	35	35	1	834	452	47.1	0.28	0.642	0.01
43	p	99	40	40	32	1	69	63	2.5	0.63	0.012	0.05
44	f	139	20	20	20	1	79	72	7.0	0.52	0.016	0.08
45	f	267	20	20	20	1	144	112	13.5	0.42	0.040	0.05
46	p	69	30	30	24	1	50	45	2.3	0.65	0.006	0.13
47	p	99	20	20	16	1	59	50	5.0	0.50	0.008	0.12
48	p	59	50	50	40	1	54	48	1.2	0.81	0.007	0.12
49	f	119	15	15	15	1	67	59	8.0	0.50	0.011	0.10
50	f	1040	10	10	10	1	525	220	105.0	0.21	0.153	0.02
51	p	69	20	20	16	1	45	39	3.5	0.57	0.005	0.15
52	f	178	15	15	15	1	97	78	12.0	0.44	0.019	0.07
53	f	99	20	20	20	1	59	58	5.0	0.58	0.011	0.10
54	f	1287	20	20	20	1	653	320	65.0	0.25	0.322	0.01
55	f	3812	30	30	30	1	1921	756	128.3	0.20	1.794	0.00
56	f	891	15	15	15	1	453	228	60.0	0.26	0.163	0.02
57	f	396	50	50	50	1	223	198	8.0	0.50	0.123	0.03
58	f	554	20	20	20	1	287	183	28.0	0.33	0.105	0.03
59	f	59	5	5	5	1	32	26	12.0	0.44	0.002	0.23
60	f	188	20	20	20	1	104	89	9.5	0.47	0.025	0.06
61	m	178	59	5	12	5	119	43	3.0	0.24	0.006	0.13
62	f	198	20	20	20	1	109	92	10.0	0.46	0.027	0.06
63	f	614	20	20	20	1	317	195	31.0	0.32	0.120	0.03
64	f	594	59	59	59	1	327	276	10.0	0.46	0.239	0.02
65	p	69	20	20	16	1	45	39	3.5	0.57	0.005	0.15
66	p	69	30	30	24	1	50	45	2.3	0.65	0.006	0.13
67	f	198	5	5	5	1	101	58	40.0	0.29	0.011	0.10
68	f	1238	10	10	10	1	624	248	125.0	0.20	0.192	0.02
69	p	119	40	40	32	1	79	71	3.0	0.60	0.016	0.06
70	f	297	30	30	30	1	163	138	10.0	0.46	0.060	0.04
71	f	644	15	15	15	1	329	183	43.3	0.28	0.105	0.03
72	f	149	20	20	20	1	84	76	7.5	0.51	0.018	0.07
73	f	1208	40	40	40	1	624	387	30.5	0.32	0.469	0.01
74	p	69	40	40	32	1	54	50	1.8	0.72	0.008	0.12
75	f	178	5	5	5	1	92	54	36.0	0.30	0.009	0.11
76	f	178	20	20	20	1	99	86	9.0	0.48	0.023	0.07
77	f	376	20	20	20	1	198	141	19.0	0.37	0.062	0.04
78	p	69	20	20	16	1	45	39	3.5	0.57	0.005	0.15
79	b	297	50	5	15	4	173	73	6.0	0.25	0.017	0.06
80	f	743	10	10	10	1	376	176	75.0	0.24	0.097	0.02
81	f	446	5	5	5	1	225	99	90.0	0.22	0.031	0.06
82	f	495	40	40	40	1	267	213	12.5	0.43	0.143	0.02
83	f	2158	50	50	50	1	1104	613	43.6	0.28	1.181	0.01
84	f	188	15	15	15	1	101	81	12.7	0.43	0.020	0.07
85	b	396	40	20	45	3	218	199	10.0	0.50	0.124	0.00
86	p	119	119	119	95	1	119	102	1.0	0.86	0.033	0.05
87	f	119	20	20	20	1	69	65	6.0	0.55	0.013	0.09

INDIVIDUAL SIZE DISTRIBUTION COUNT DATA

Client Name: NRaD  
 Client Project#: Paper sizing  
 ETA Project # : 94-4274

Client Sample # : P8-1  
 ETA Sample # : 4274-4

Particle Number	Particle Type	Length (µm)	Structure Dia.(µm)	Fiber Dia.(µm)	Thickness (µm)	# of fibers in struc.	X-section Dia.(µm)	Hydro. Dia.(µm)	Aspect Ratio	Particle Sphericity	Surface Area(mm2)	Sur.Area / Vol. Ratio
88	p	79	30	30	24	1	54	49	2.7	0.62	0.008	0.122
89	f	208	5	5	5	1	106	60	42.0	0.29	0.011	0.100
90	f	129	10	10	10	1	69	55	13.0	0.43	0.009	0.110
91	p	109	30	30	24	1	69	61	3.7	0.56	0.012	0.099
92	f	218	10	10	10	1	114	78	22.0	0.36	0.019	0.077
93	f	198	5	5	5	1	101	58	40.0	0.29	0.011	0.104
94	f	1089	20	20	20	1	554	286	55.0	0.26	0.258	0.021
95	f	99	20	20	20	1	59	58	5.0	0.58	0.011	0.104
96	f	1337	20	20	20	1	678	328	67.5	0.25	0.339	0.018
97	p	50	30	30	24	1	40	36	1.7	0.73	0.004	0.167
98	f	1386	20	20	20	1	703	336	70.0	0.24	0.355	0.018
99	p	69	20	20	16	1	45	39	3.5	0.57	0.005	0.153
100	f	673	25	25	25	1	349	224	27.2	0.33	0.157	0.027

## MICROSCOPIC SIZE ANALYSIS DEFINITIONS AND RULES

### Structure definitions:

Structure :	An individual particle composed of one or more paper fibers.
Paper particle :	Paper particle with an aspect ratio of $< 5 : 1$
Fiber:	Paper particle with an aspect ratio of $\geq 5 : 1$
Bundle :	3 or more fibers contacting each other and in parallel arrangement
Matrix :	3 or more fibers intersecting or attached to a central point

### Counting rules:

A total of 100 "paper" structures greater than 5 divisions (49.5 $\mu$ m) are counted at a magnification of 50X  
 The largest structure on the first slide analyzed is sized first to account for a negative bias in counting large structures  
 Fields are established using the the "left" edge of a grid line, tabulating every 3rd structure crossing the vertical cross-hair.  
 After every 5th structure counted, the slide is moved to the next horizontal grid square.  
 After every 25 structures counted, a new slide (comprising a different quarter section of the same filter) is counted.  
 All numerical size distribution statistics are based on the arithmetic mean diameter.  
 The estimated mass distribution is based on particle volume (formula for a sphere) in each size category, and does not take into account particle specific gravity.

### Statistical Parameter Definitions & Formulas :

Hydrodynamic Diameter	Arithmetic mean of "structure" length, width, and approximate thickness using the sphericity coefficient. Diameter = Structure length * Sphericity
Cross-section Diameter	Arithmetic mean of "structure" length and width not accounting for particle thickness. Diameter = (Structure length + structure diam.) / 2
Median	Number in the middle of a distribution; that is, half the values are greater than the median, and half the values below.
Mode	Most frequently occurring range in a size distribution. The largest size category is reported in bimodal distributions.
Skewness	Degree of symmetry of a population around its mean. Positive skewness indicates a distribution with an asymmetric tail towards more positive values. Negative skewness indicates an asymmetric tail towards more negative values.
Kurtosis	Relative peakedness or flatness of a distribution compared to the normal distribution. Positive kurtosis indicates a relatively peaked distribution. Negative kurtosis indicates a relatively flat distribution.
95% C.L.	95% Confidence Limit (i.e. probability that 95% of time the mean value will fall within the specified size range).
Aspect Ratio	Ratio of particle length divided by the particle width
Particle Sphericity	Measure of effective particle size based on the formula (thickness <sup>2</sup> / (length*width)) <sup>0.333</sup>
Surface Area	$4\pi r^2$
Volume	$1.33\pi r^3$

## APPENDIX C

### LIQUID PHASE ORGANISM TOXICITY TESTING REPORT

Source: Marine Acute Toxicity Test Results of Pulped Paper/Cardboard and Shredded Metal to Mysid Shrimp (*Mysidopsis bahia*), Silverside Minnows (*Menidia beryllina*), Marine Dinoflagellates (*Gonyaulax polyedra*), Bacteriam (*Photobacterium phosphoreum*), and Chain Diatoms (*Skeletonema costatum*).  
San Diego, California  
Naval Command, Control & Ocean Surveillance Center, RDTE  
Division, Code 522, 1995

**Marine Acute Toxicity Test Results of Paper Pulp and Shredded Metal to Mysid Shrimp (*Mysidopsis bahia*), Silverside Minnows (*Menidia beryllina*), A Marine Dinoflagellate (*Gonyaulax polyedra*), a bacterium (*Photobacterium phosphoreum*), and a Chain Diatom (*Skeletonema costatum*)**

By

David Lapota<sup>1</sup>, Debbie Duckworth<sup>1</sup>, Gunther Rosen<sup>2</sup>,  
Jon Groves<sup>2</sup>, and Bradley Davidson<sup>2</sup>

<sup>1</sup>Naval Command, Control & Ocean Surveillance Center, RDTE Division, Code 522  
53475 Strothe Road, San Diego, CA 92152-6310  
Phone: (619) 553-2798; 2815  
FAX: (619) 553-6305  
email: lapota@nosc.mil

<sup>2</sup>Computer Sciences Corporation  
4045 Hancock Street, San Diego, CA 92110-5164

July 1995

A report prepared for:

Bart Chadwick

## EXECUTIVE SUMMARY

A series of static-renewal EPA acceptable bioassays were conducted to estimate the potential toxicity of 2 leachable materials. The materials, labelled Paper Pulp and Shredded Metal, were tested on the mysid shrimp (*Mysidopsis bahia*), the minnow (*Menidia beryllina*), the marine chain diatom (*Skeletonema costatum*, clone "Skel"), the Microtox bioassay (*Photobacterium phosphoreum*), and the Qwiklite Bioassay System, which uses the bioluminescent dinoflagellate, *Gonyaulax polyedra*. The marine diatom was used for the chlorophyll assays. Bioassay organisms representing different phyla were chosen and tested to represent a potential "risk" to the marine environment. *Mysidopsis bahia* was chosen to represent a benthic or bottom dwelling animal response, while the minnow *Menidia beryllina* was chosen to represent a pelagic or swimming animal response. The phytoplankton chain diatom species, *Skeletonema costatum* and the bioluminescent dinoflagellate, *Gonyaulax polyedra*, were used to observe any potential effect on the primary producers in marine waters. The endpoints measured were the concentration at which 50% of test organisms were affected (LC50/IC50) and the concentration at which no observable effect occurred (NOEC). The effects measured varied depending on the test species and were: survival in the mysids and minnows (LC50s), inhibition of bioluminescence of *G. polyedra* and the bacterium (IC50), and biomass or chlorophyll fluorescence (IC50) in the diatom tests.

Toxicity was observed in the mysid when exposed to a 5% leachate of Paper Pulp. No NOEC or LC50 value could be determined as significant toxicity was observed at the lowest leachate concentration and did not follow a dose response curve. Assays where the 5% leachate of Paper Pulp was centrifuged, less toxicity was observed and a dose response occurred. In only one mysid assay was a NOEC value observed at 6.25% (5%) leachate. No toxicity was observed in fish when exposed to 5% leachate of Paper Pulp. Each of the fish assays resulted in NOEC values of 100% (5%) Paper Pulp leachate. When tested with the diatom, *S. costatum*, a 5% leachate of Paper Pulp resulted in an IC50 value of between 12.5 and 50% leachate and NOEC values of 12.5% and 25% leachate respectively. The 5% Paper Pulp caused a dose response in the dinoflagellate, *G. polyedra*, where the IC50 was 27.7% and a NOEC was not applicable after 96 hours of exposure. A 0.01% leachate Paper Pulp resulted in an NOEC value of 100% due to variable bioluminescence levels which did not indicate a dose response curve to either hormesis or inhibition in the dinoflagellate, *G. polyedra*. There was up to 31% reduction of light output from bacteria after 5 minutes of exposure to 5% Paper leachate and an average of 20% reduction of light output from the bacterium after 5 minutes of exposure to 0.01% Paper leachate. The NOEC value after 96 hours of exposure to 5% leachate was 50% (5%) leachate and a NOEC value was not applicable after exposure to 0.01% leachate.

5% leachate of Shredded Metal had no adverse effects on mysid or fish. An IC50 value of 58.7% and an NOEC value of 25% was observed when *S. costatum* was exposed to 25% leachate of Shredded Metal. The 5% leachate Shredded Metal had no

adverse effect on the diatom resulting in an NOEC value of 100% (5%) leachate after 96 hours. When tested with the bioluminescent dinoflagellate, the 25% and 5% leachate resulted in similar IC50 values, 18.8% and 18.7% respectively. A NOEC value was not applicable when the dinoflagellate was exposed to 5% leachate but equal to 6.25% when tested with (25%) leachate. The 25% Metal leachate reduced light output from the bacterium by 37% after 5 minutes while the 5% leachate reduced light output by 23% after 5 minutes of exposure. NOEC values were not applicable for either concentration of leachate in the Microtox assays.

**Complete Progress of Bioassays  
for  
Paper Pulp & Shredded Metal Study**

	<i>Mysidopsis bahia</i>	<i>Menidia beryllina</i>	<i>Skeletonema costatum</i>	<i>Gonyaulax polyedra</i>	<i>Photo-bacterium</i>
Paper Pulp					
25%					
5%	XXX	XXX	XXX	X	XXX
0.01%			X	X	XXX
Shredded Metal					
25%			X	X	XXX
5%	XXX	XXX	X	X	XXX

## TABLE OF CONTENTS

	Page
EXECUTIVE SUMMARY .....	2
INTRODUCTION .....	6
MATERIALS AND METHODS .....	6
RESULTS .....	12
FIGURES .....	18-47
RAW DATA, FISH & MYSID ASSAYS .....	48-67
RAW DATA, DIATOM ASSAYS .....	68-72
RAW DATA, MICROTOX ASSAYS .....	73-77
RAW DATA, QWIKLITE ASSAYS .....	78-93

## INTRODUCTION

A series of static-renewal EPA (Environmental Protection Agency) acceptable bioassays, in addition to a suite of Microtox and Qwiklite Bioluminescence Assays, were conducted to estimate the potential toxicity of 2 materials. The materials, labelled Paper Pulp and Shredded Metal, were tested on the mysid shrimp (*Mysidopsis bahia*), the minnow (*Menidia beryllina*), the bioluminescent dinoflagellate (*Gonyaulax polyedra*), the bioluminescent bacteria (*Photobacterium phosphoreum*), and the marine chain diatom (*Skeletonema costatum*, clone "Skel"). The marine diatom was used for the chlorophyll assays. Bioassay organisms representing different phyla were chosen and tested to represent a potential "risk" to the marine environment. *Mysidopsis bahia* was chosen to represent a benthic or bottom dwelling animal response, while the minnow *Menidia beryllina* was chosen to represent a pelagic or swimming animal response. The phytoplankton chain diatom species and the dinoflagellate were used to observe any potential effect on the primary producers in marine waters. The endpoints measured were the concentration at which 50% of test organisms were affected (LC50/IC50) and the concentration at which no observable effect occurred (NOEC). The effects measured varied depending on the test species and were: survival in the mysids and minnows (LC50s), inhibition of bioluminescence of *G. polyedra* and the bacterium (IC50), and biomass or chlorophyll fluorescence (IC50) in the diatom tests.

## MATERIALS AND METHODS

### Test Equipment Preparation for Mysids and Minnows

All test chambers were constructed from borosilicate glass beakers with lids. All beakers were washed with a critical cleaner and rinsed with 10% nitric acid. Three deionized water rinses followed each cleaning procedure. All acute toxicity tests with the mysids were conducted in 300 ml beakers with 200 ml of dilution water. The minnows were maintained in 400 ml beakers with 250 ml of dilution water. The diatom assay required 125 ml Erlenmeyer flasks containing 25 ml of test solution. The bioluminescent assays used spectrophotometric grade cuvettes to contain approximately three ml of test solution.

#### Source and Acclimation of Test Species

Several day old *M. bahia* and *M. beryllina* were shipped overnight from Aquatic Indicators, St. Augustine, FL, to our laboratory. Both the mysids and minnows were slowly acclimated in a 25°C water bath and transferred by pipette to several holding tanks with filtered (0.45  $\mu\text{m}$ ) seawater. The dilution water used in testing was obtained from the NCCOSC Biological Effects Program bioassay facility located near the mouth of San Diego Bay. Water was filtered through a coarse sand filter prior to final filtration (0.45  $\mu\text{m}$ ). The test animals were slowly acclimated to the test water salinity of 33 parts-per-thousand over several days. All test animals were fed daily with freshly hatched *Artemia* brine shrimp.

The marine diatom, *Skeletonema costatum*, clone "Skel" was obtained from the UCLA, Hopkins Marine Station. The cultures were maintained on an enriched seawater medium (ESM) using filtered (0.20  $\mu\text{m}$ ) seawater collected from the Scripps Institute of Oceanography pier pump system in La Jolla. Samples of the stock were routinely

aliquoted into fresh media to maintain high cell densities. The diatom was cultured at room temperature ( $\sim 25^{\circ}\text{C}$ ) under cool white fluorescent bulbs at a light intensity of approximately 4000 lux for 12 hours per day.

### **Sea Water Extraction of Paper Pulp and Shredded Metal Materials**

Test Solution was attained by leaching each material in filtered sea water for one and a half hours, in which 30 min of mixing period was followed by one hour of settling period (Elutriate Preparation, EPA protocol, 1991).

A 25% elutriate was prepared (EPA protocol, 1991) by subsampling 1 L of filtered sea water exposed to 250 grams of the homogenized material (1 : 4 ratio). This 25% elutriate was then used as 100% test solution. For testing purposes and to determine a dose response curve, the 100% test solution was diluted with filtered sea water by half until 6.25%. The test solutions for every assay ranged between 100% and 6.25% elutriate. In cases of expected extreme toxicity, a 5% elutriate was prepared (50 gm/1 L). The Paper Pulp was subject to a dry : wet weight conversion factor (1 : 6.3) due to the high percentage of water in the material. Additional assays with the Paper Pulp were conducted to observe a no effect level of exposure using a 0.01% elutriate (0.630 gm/1 L). The supernatant was carefully removed from the material with the use of a mesh filter. The elutriate resulting from leaching was then used to make dilutions of the test solution. Following the first three assays exposing mysids to an elutriate of Paper Pulp, suspended solids were suspected of causing the toxicity and the elutriates were then centrifuged following the leaching procedure. Centrifugation was for 7 minutes at 1800 rpm at  $25^{\circ}\text{C}$  on a Damon, IEC Centra-8R Centrifuge.

## Experimental Test Design and Procedure for *Mysidopsis bahia* and *Menidia beryllina*

Toxicity testing of Paper Pulp and Shredded Metal consisted of 96 hour static renewal acute tests. These assays were conducted to test for potential toxicity arising from exposure to the leachates.

Environmental Protection Agency (EPA) test protocols were followed for the mysid and minnow bioassays (U.S. EPA, 1988). For acute bioassays, test chamber sizes for the mysids and minnows were typically 300 and 400 ml beakers filled with 200 ml and 250 ml of solution, respectively. The average age of the animals were 5 days and 13 days at the start of the bioassays for mysids and minnows, respectively. The mysids and minnows were set up at 10 animals per beaker with two replicates for each concentration. Each assay began when test species were distributed to test beakers with 50 ml of filtered seawater (0.45  $\mu\text{m}$ ). Animals were then pipetted from holding tanks into test beakers. Dilutions of the material leachate were added to each beaker to a final volume of either 200 or 250 ml. All animals were fed daily newly hatched *Artemia* brine shrimp. The test beakers were covered with glass lids and placed in a temperature controlled bath at 25°C. Solutions were renewed every 24 hours at which time fecal material was removed and seawater chemistry measurements were recorded. Survival was recorded every 24 hours. Seawater parameters measured daily were dissolved oxygen, pH, and temperature. Minimum requirements for test acceptability for dissolved oxygen are 40% saturation for acute tests and the seawater temperature must not fluctuate more than  $\pm 2^\circ\text{C}$ .

Test concentrations of 100%, 50%, 25%, 12.5%, and 6.25% with a seawater control were used for both materials in the mysid acute tests and the minnow acute tests.

Percent survival was calculated and graphed. A probit analysis was performed to estimate  $LC_{50,s}$  (lethal concentration to cause mortality in 50% of the tested population), where appropriate. All data were analyzed using Toxis II and Prodas statistical programs.

#### **Experimental Test Design and Procedure for Diatom Biomass (Fluorescence) in *Skeletonema costatum* (Clone "Skel")**

Prior to testing, monocultures of *Skeletonema* were maintained in enriched seawater medium (ESM) in 2 L borosilicate Erlenmeyer flasks under a light regime of 12:12 hours (light : dark) at a light intensity of approximately 4000 lux from cool white fluorescent bulbs. Culture temperature was maintained near 19°C. This bioassay was conducted in accordance with the American Society for Testing and Materials Standard Guide for Conducting Static 96-hr Toxicity Test with Microalgae (E 1218) (ASTM, 1992). At the beginning of each bioassay, 400  $\mu$ l of diatom stock was introduced into three replicate Erlenmeyer flasks containing a combined 150 ml of leachate and filtered seawater for the controls and different concentrations of the elutriate. The dilution water was collected from the pumped seawater system at the pier of the Scripps Institute of Oceanography in La Jolla. All seawater was filtered with membrane filters to 0.2  $\mu$ m and enriched as the stock cultures. Both the Paper Pulp and Shredded Metal materials were tested at concentrations of 100%, 50%, 25%, 12.5%, and 6.25% with seawater controls. All elutriate concentrations were nominal values. The control groups

received no addition of elutriate and did not exhibit background fluorescence. A Turner Model 112 fluorometer was used to measure *in-vivo* fluorescence from the diatom cells. The fluorometer was equipped with a combination T-5 lamp, a red-sensitive photomultiplier tube (R-136), a blue (5-60) excitation filter, and a red (2-64) filter to detect fluorescence at wavelengths > 640 nanometers (nm). Chlorophyll *a* fluorescence has maximum emission at 663 nm. The instrument was blanked between readings with filtered (0.45  $\mu\text{m}$ ) seawater. All flasks were read within 1 hour after the introduction of the diatoms into the flasks and at 24-hour intervals for a period of 96 hours. The measured fluorescence is directly related to cell number and to the presence of viable diatom cells relative to the leachate concentration. Mean relative fluorescence, standard deviation, and the coefficient of variation were calculated for each control and leachate concentration. Relative fluorescence, calculated as a percentage of control values, was plotted over time during the test.

#### **Experimental Test Design for Microtox (Bioluminescence) Assay**

The Microtox Bioassay System is an acute toxicity test utilizing a specially cultured bioluminescent bacteria (*Photobacterium phosphoreum*). The test is employed for the determination of a dose response curve, from which the inhibition concentration (IC) of test solution causing a specified effect is found. The method measures the effect on the bioluminescent light output of the bacteria as they are challenged by the test solution. Observations of light output are recorded at 5 and 15 minutes of exposure. This test is usually used as a screen test for toxic effects. Three trials of the Paper Pulp assay were performed with four dilutions and a control. Five minute and 15-minute readings were

taken. Both the EC20 and EC50 were determined graphing the calculated Microtox statistic on log/log paper. Also, the percent reduction of light output at the 100% leachate dilution was calculated.

#### **Experimental Test Design for QWIKLITE (Bioluminescence) Bioassay System**

The QWIKLITE Bioassay also measures the inhibition of light emitted by the bioluminescent dinoflagellate, *Gonyaulax polyedra*, exposed to a test solution. The test lasts 96 hours and results are expressed as the percent of control in which all dilutions are compared to the controls. Toxicity results are reported as the IC50 when a dose response is the effect of exposure.

Testing of the dinoflagellates is accomplished by placing individual cuvettes containing the test material, media, and cells into a darkened test chamber which is attached to a photomultiplier tube (PMT). We have used our QWIKLITE bioassay system which uses a 2-inch diameter 8575 PMT with an S-20 response used in the photon count mode. The top of the test chamber is removable and houses a small adjustable motor which drives a stainless steel shaft terminating in a plastic propeller. The propeller is seated into the cuvette and as the contents are stirred, bioluminescence is generated and measured by the PMT. Each test period is completed at 24 hour intervals thereafter until completion of the bioassay. Mean light output (PMT counts) is calculated for each experimental group and control. Light output means are then graphed as light output (percent of control) as a function of time. All graphs represent the data collected at 96 hours of exposure.

## RESULTS

### Effects of Paper Pulp and Shredded Metal to *Mysidopsis bahia*

#### Paper Pulp - 5% Leachate

Three assays resulted in 15% to 60% mortality at the lowest concentration (6.25%) and lethality in all higher concentrations of test solution (Figure 1. - 3). No dose response was observable. Suspended solids in the elutriate were suspected of contributing to toxicity. Consequently, no LC<sub>50</sub> was observed in *Mysidopsis bahia* from this material leachate and further assays would require centrifugation of the elutriate prior to becoming a test solution. A NOEC value was not applicable due to the observed effects.

#### Paper Pulp - 5% Leachate. Centrifuged

Two of three assays resulted in LC50 values of 22% and 32% test solution at 96 hours of exposure (Figure 4.- 5.). The third assay resulted in total lethality (97.5% mortality) in concentrations 12.5 thru 100% (Figure 6.) No LC50 was observable in *Mysidopsis bahia* from the third assay of this leachate. After 96 hours of exposure, a NOEC value was not applicable to two of the three assays due to observed effects but equal to 6.25% (5%) leachate in the third assay.

#### Shredded Metal - 5% Leachate. Centrifuged

Three assays conducted resulted in no significant mortality. No dose response observed (figure 19.-21). No LC50 was observable in *Mysidopsis bahia* from the three assays of this leachate. After 96 hours of exposure, the NOEC value was 100% leachate in each of the three assays.

## Effects of Paper Pulp and Shredded Metal to *Menidia beryllina*

### Paper Pulp - 5% Leachate, Centrifuged

In three assays conducted, no significant mortality occurred in any test concentration (Figure 7. - 9.). No LC<sub>50</sub> was observed in *Menidia beryllina* from this material leachate. After 96 hours of exposure, the NOEC value was 100% in each of the three assays.

### Shredded Metal - 5% Leachate, Centrifuged

Three assays conducted resulted in no mortality in any test concentration (Figure 22. - 24.). No LC<sub>50</sub> was observed in *Menidia beryllina* from this material leachate. After 96 hours of exposure, the NOEC value was 100% in each of the three assays.

## Effects of Paper Pulp and Shredded Metal to *Skeletonema costatum* (Clone "Skel")

### Paper Pulp - 5% Leachate, Centrifuged

Three assays resulted in dose responses where the IC50 values at 96 hours of exposure observed to be: 12.5% - 25%, 25%, and 25 - 50% respectively (Figure 10. - 12.). A decline in plant biomass with increased concentration of test solution was consistently observed daily until the assays ended at 96 hours. After 96 hours of exposure, the NOEC value in two of the three assays was 12.5% (5%) leachate and 25% (5%) leachate in the third assay.

### Paper Pulp - 0.01% Leachate, Centrifuged

One assay conducted resulted in no decline of biomass (Figure 13.). A slight enhancement of growth was observed in conjunction with increased test solution. No IC50 value was observable in *Skeletonema costatum* from this leachate. After 96 hours of

exposure, the NOEC value was 100% (0.01%) leachate.

#### Shredded Metal - 25% Leachate, Centrifuged

A 25% Shredded Metal leachate assay resulted in a dose response and an IC50 value of 59% leachate (Figure 14.). After 96 hours of exposure, the NOEC value was equal to 25% (25%) leachate.

#### Shredded Metal - 5% Leachate, Centrifuged

One assay resulted in no decline or enhancement of plant biomass (Figure 15.). No IC50 was observable in *Skeletonema costatum* from this leachate. After 96 hours of exposure, the NOEC value was 100% (5%) leachate.

#### **Effects of Paper Pulp and Shredded Metal to *Gonyaulax polyedra***

#### Paper Pulp - 5% Leachate, Centrifuged

One assay resulted in a dose response curve where after 95 hours of exposure, no NOEC value was applicable and an IC50 value of 27.7% (5%) leachate was observed (Figure 16.).

#### Paper Pulp - 0.01% Leachate, Centrifuged

One assay resulted in variable levels of bioluminescence (Figure 17.). A poor dose response resulted in no observable IC50 value. After 96 hours of exposure, a NOEC value was observed at 100% (0.01%) leachate.

#### Shredded Metal - 25% Leachate, Centrifuged

One assay resulted in a dose response curve and an IC50 value at 96 hours at an 18.8% leachate (Figure 18.). After 96 hours of exposure, the NOEC value was 6.25% (25%) leachate.

### Shredded Metal - 5% Leachate, Centrifuged

One assay conducted resulted in a dose response curve and an IC50 value at 96 hours at an of 18.7% leachate (Figure 25.). An NOEC value was not applicable after 96 hours of exposure.

### **Effects of Paper Pulp and Shredded Metal to *Photobacterium phosphoreum* (Microtox)**

#### Paper Pulp - 5% Leachate, Centrifuged

Trial 1 and 2 showed a 5-minute EC20 of 76% and 98%, respectively. The five minute EC50 in both those trials were at or exceeded 100%, the maximum dilution tested. The third trial showed no toxicity as the control and the 100% leachate reading were essentially the same. The 15-minute readings on trial 1 were inconclusive for determining EC values because all mean readings for the dilutions except 100% exceeded the control mean (Figure 26.). This yielded only one usable point, and a dose response curve could not be plotted. After 15 minutes of exposure, the NOEC value was 100% (5%) leachate in each of the three trials.

#### Paper Pulp - 0.01% Leachate, Centrifuged

A 0.01% Paper Pulp leachate was tested using four concentrations and a control for 5 and 15 minutes of exposure (Figure 28.). After 5 minutes, an EC20 value was observed at 90% (0.01%) leachate. No EC50 value was noted, although 20% reduction of light output occurred at 100% leachate. After 15 minutes, an EC20 value of 60% leachate was observed and 13% reduction of light output at 100% leachate. After 15 minutes of exposure, a NOEC value was not applicable due to effects observed.

#### Shredded Metal - 25% Leachate, Centrifuged & Non-centrifuged


A 25% Metal leachate, centrifuged and non-centrifuged, were tested using four dilutions, and a control, for 5 and 15 minutes of exposure time (Figure 27.). Only one trial was performed. The centrifuged sample appeared more toxic than the uncentrifuged. This may be due to enhanced stimulation in the uncentrifuged sample due to a white residue, resembling vegetable shortening, observed on the metal pieces used for the leachate. The 5 and 15 minute EC20 for the centrifuged sample are 27.5 and 46%, respectively. Both EC50 values exceeded 100% leachate. The 5 and 15 minute EC20 values for the uncentrifuged sample were 89 and 80% respectively. The EC50 values for this sample both exceeded 100%. A NOEC value was not applicable after 15 minutes of exposure due to an effect observed in the lowest concentration tested.

#### Shredded Metal - 5% Leachate, Centrifuged


A 5% Metal leachate, centrifuged, was tested using four dilutions and a control for 5 and 15 minutes of exposure (Figure 29.). After 5 minutes, an EC20 value at 44% leachate was observed. No EC50 was noted although there was a 23% reduction in light output at 100% leachate. After 15 minutes exposure, no EC20 or EC50 value was noted and only 13% reduction of light output occurred in 100% (5%) leachate. A NOEC value was not applicable after 5 or 15 minutes of exposure due to effects observed in the lowest concentration tested.

**Figure 1. Paper Pulp - 5% Leachate**  
*Mysidopsis bahia* - Trial #1


Poor Dose Response  
NOEC not applicable



**Figure 2. Paper Pulp - 5% Leachate**  
*Mysidopsis bahia* - Trial #2




**Figure 3. Paper Pulp - 5% Leachate**  
*Mysidopsis bahia* - Trial #3



# Figure 4. Paper Pulp - 5% Leachate Centrifuged

## *Mysidopsis bahia* - Trial #1

LC50 = 32% of 5% Leachate, NOEC = 6.25%




# Figure 5. Paper Pulp - 5% Leachate

## Centrifuged


### *Mysidopsis bahia* - Trial #2

LC50 = 22% of 5% Leachate

NOEC not applicable




**Figure 6. Paper Pulp - 5% Leachate  
Centrifuged  
*Mysidopsis bahia* - Trial #3**




# Figure 7. Paper Pulp - 5% Leachate Centrifuged

## *Menidia beryllina* - Trial #1




**Figure 8. Paper Pulp - 5% Leachate  
Centrifuged  
*Menidia beryllina* - Trial #2**



# Figure 9. Paper Pulp - 5% Leachate Centrifuged


## *Meridia beryllina* - Trial #3



# Figure 10. Paper Pulp - 5% Leachate Centrifuged

## *Skeletonema costatum* - Trial #1

IC50 = 12.5% < x < 25% (5% centrifuged) Leachate  
NOEC = 12.5% (5%) Leachate




# Figure 11. Paper Pulp - 5% Leachate Centrifuged

## *Skeletonema costatum* - Trial #2

IC50 = 25% (5%) Leachate


NOEC = 12.5% (5%) Leachate



# Figure 12. Paper Pulp - 5% Leachate Centrifuged

## *Skeletonema costatum* - Trial #3

IC50 = 25% < x < 50% (5%) Leachate  
NOEC = 25% (5%) Leachate




# Figure 13. Paper Pulp - 0.01% Leachate Centrifuged


## *Skeletonema costatum* - Trial #1

No Toxicity Noted

NOEC = 100% (0.01%) Leachate



**Figure 14. Shredded Metal - 25%  
Leachate, Centrifuged  
*Skeletonema costatum* - Trial #1**



**Figure 15. Shredded Metal - 5%  
Leachate, Centrifuged  
*Skeletonema costatum* - Trial #1**





Figure 16. Paper Pulp - 5% Leachate,  
Centrifuged


*Gonyaulax polyedra* - Trial #1




# Figure 17. Paper Pulp - 0.01% Leachate

## Centrifuged


### *Gonyaulax polyedra* - Trial #1




**Figure 18. Shredded Metal - 25%  
Leachate, Centrifuged  
*Gonyaulax polyedra* - Trial #1**




**Figure 19. Shredded Metal  
5% Leachate, Centrifuged  
*Mysidopsis bahia* - Trial #1**




**Figure 20. Shredded Metal  
5% Leachate, Centrifuged  
*Mysidopsis bahia* - Trial #2**




**Figure 21. Shredded Metal  
5% Leachate, Centrifuged  
*Mysidopsis bahia* - Trial #3**




**Figure 22. Shredded Metal  
5% Leachate, Centrifuged  
*Menidia beryllina* - Trial #1**




**Figure 23. Shredded Metal  
5% Leachate, Centrifuged  
*Menidia beryllina* - Trial #2**




**Figure 24. Shredded Metal  
5% Leachate, Centrifuged  
*Menidia beryllina* - Trial #3**




**Figure 25. Shredded Metal  
5% Leachate, Centrifuged  
*Gonyaulax polyedra* - Trial #1**




**Figure 26. Paper Pulp  
5% Leachate, Centrifuged  
*Photobacterium*, Microtox - Trial #1-3**




**Figure 27. Shredded Metal  
25% Leachate, Centrifuged  
*Photobacterium*, Microtox - Trial #1**




# Figure 28. Paper Pulp 0.01% Leachate, Centrifuged *Photobacterium*, Microtox - Trial #1




# Figure 29. Shredded Metal 5% Leachate, Centrifuged *Photobacterium*, Microtox - Trial #1




# Figure 30. Paper Pulp 0.01% Leachate, Centrifuged *Mysidopsis bahia*, - Trial #1



**Figure 31. Paper Pulp  
0.01% Leachate, Centrifuged  
*Mysidopsis bahia*, - Trial #2**



**Figure 32. Paper Pulp  
0.01% Leachate, Centrifuged  
*Mysidopsis bahia*, - Trial #3**



RAW DATA, Refer to  
Figure #1

TEST DATE

TEST NUMBER

Start: 25-Apr-95

0000004024

SPECIES: Mysidopsis bahia

TEST MATERIAL: Paper Pulp

SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

TEST TYPE: (96) hr acute ( ) chronic

REFERENCE TOXICANT TEST:

MYSID TEST DATA

Test Number: 0000004024

( ) Chronic (x) Acute 96 hours

Test Date: 25-Apr-95

Source:

Test Material: BPP (%)

Conc	Rep No.	Cont.	Start	Daily Survival						Prop	Females	Proc	Weight	
				1	2	3	4	5	6	End	Alive	w/eggs	w/eggs	/Mysid
0.000	1	6	10	9	9	9	9				.90			
0.000	2	12	10	10	9	8	8				.80			
6.250	1	9	10	7	7	7	7				.70			
6.250	2	8	10	5	5	5	5				.50			
12.500	1	10	10	0	0	0	0				0.00			
12.500	2	2	10	3	3	2	2				.20			
25.000	1	11	10	0	0	0	0				0.00			
25.000	2	7	10	0	0	0	0				0.00			
50.000	1	1	10	0	0	0	0				0.00			
50.000	2	3	10	0	0	0	0				0.00			
100.000	1	5	10	0	0	0	0				0.00			
100.000	2	4	10	0	0	0	0				0.00			

RAW DATA, REFER TO FIG. 2

TEST DATE

TEST NUMBER

Start: 25-Apr-95

0000004025

SPECIES: Mysidopsis bahia

TEST MATERIAL: Paper Pulp

SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

TEST TYPE: (96) hr acute ( ) chronic

REFERENCE TOXICANT TEST:

MYSID TEST DATA

Test Number: 0000004025

( ) Chronic (x) Acute 96 hours

Test Date: 25-Apr-95

Source:

Test Material: BPP (%)

Conc	Rep No.	Cont.	Start	Daily Survival						Proc Alive	Females w/eggs	Proc w/eggs	Weight /Mysid
				1	2	3	4	5	6 End				
0.000	1	3	10	9	9	8	8		.80				
0.000	2	2	10	9	9	9	9		.90				
6.250	1	4	10	0	0	0	0		0.00				
6.250	2	5	10	3	3	3	3		.30				
12.500	1	10	10	0	0	0	0		0.00				
12.500	2	6	10	6	6	6	6		.60				
25.000	1	9	10	6	6	6	6		.60				
25.000	2	11	10	0	0	0	0		0.00				
50.000	1	8	10	0	0	0	0		0.00				
50.000	2	12	10	3	3	3	3		.30				
100.000	1	7	10	2	2	1	1		.10				
100.000	2	1	10	3	1	1	1		.10				

RAW DATA REFER TO FIG. 3

TEST DATE

TEST NUMBER

Start: 25-Apr-95

0000004026

SPECIES: Mysidopsis bahia

TEST MATERIAL: Paper Pulp

SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

TEST TYPE: (96) hr acute ( ) chronic

REFERENCE TOXICANT TEST:

=====

MYSID TEST DATA

=====

Test Number: 0000004026 ( ) Chronic (x) Acute 96 hours

Test Date: 25-Apr-95

Source:

Test Material: BPP (%)

Conc	Rep No.	Cont.	Start	Daily Survival						Proc Alive	Females w/eggs	Proc w/eggs	Weight /Mysid
				1	2	3	4	5	6				
0.000	1	4	10	10	10	10	10		1.00				
0.000	2	5	10	10	10	8	8		.80				
6.250	1	2	10	6	6	6	6		.60				
6.250	2	9	10	3	2	2	2		.20				
12.500	1	1	10	5	5	6	6		.60				
12.500	2	6	10	0	0	0	0		0.00				
25.000	1	3	10	0	0	0	0		0.00				
25.000	2	12	10	0	0	0	0		0.00				
50.000	1	10	10	4	4	4	4		.40				
50.000	2	11	10	5	5	4	4		.40				
100.000	1	7	10	2	2	2	2		.20				
100.000	2	8	10	4	4	4	4		.40				

RAW DATA, REFER TO FIG. 4

TEST DATE	TEST NUMBER
Start: 2-May-95	0000004028

SPECIES: Mysidopsis bahia  
 TEST MATERIAL: Paper Pulp  
 SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

TEST TYPE: (96) hr acute ( ) chronic

REFERENCE TOXICANT TEST:

=====

MYSID TEST DATA

=====

Test Number: 0000004028 ( ) Chronic (x) Acute 96 hours  
 Test Date: 2-May-95  
 Source: Test Material: BPP (X)

Conc	Rep No.	Cont. Start	Daily Survival						Prop Alive	Females w/eggs	Proo w/eggs	Weight /Mysid
			1	2	3	4	5	6				
0.000	1	10	10	10	10	10	10	1.00				
0.000	2	1	10	9	9	9	8	.80				
6.250	1	2	10	8	8	8	8	.80				
6.250	2	3	10	9	9	9	9	.90				
12.500	1	12	10	5	5	5	5	.50				
12.500	2	8	10	5	4	4	4	.40				
25.000	1	6	10	8	8	7	7	.70				
25.000	2	9	10	8	6	6	5	.50				
50.000	1	7	10	1	1	1	1	.10				
50.000	2	4	10	6	2	2	2	.20				
100.000	1	5	10	7	4	3	3	.30				
100.000	2	11	10	0	0	0	0	0.00				

WATER QUALITY

Test Number: 0000004028

Test Date: 2-May-1995

Source:

	pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
Test											
Minimum	7.70	.5	32.00	25.0							
Maximum	8.00	6.3	32.00	26.0							

	pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
Day: 0											
Minimum	7.90	6.1	32.00	25.5							
Maximum	8.00	6.3	32.00	25.5							

Day: 1											
Minimum	7.70	.5	32.00	26.0							
Maximum	7.90	4.8	32.00	26.0							

Day: 2											
Minimum	7.70	1.1	32.00	25.5							
Maximum	7.90	4.4	32.00	25.5							

Day: 3											
Minimum	7.70	1.3	32.00	25.0							
Maximum	7.90	5.1	32.00	25.0							

Day: 4											
Minimum	7.80	2.3	32.00	25.5							
Maximum	7.90	5.0	32.00	25.5							

WATER QUALITY

Test Number: 0000004028

Test Date: 2-May-1995

Source:

Container	Conc	pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
-----------	------	----	----	-------	------	------	------	-----	-----	-------	-----	----

Day: 0 Time: 1400

0.00 D	8.00	6.29	32.00	25.5
6.25 D	8.00	6.17	32.00	25.5
12.50 D	8.00	6.11	32.00	25.5
25.00 D	8.00	6.29	32.00	25.5
50.00 D	7.90	6.05	32.00	25.5
100.00 D	7.90	6.31	32.00	25.5

Day: 1 Time: 1400

0.00 D	7.90	4.80	32.00	26.0
6.25 D	7.70	3.98	32.00	26.0
12.50 D	7.70	1.82	32.00	26.0
25.00 D	7.80	2.00	32.00	26.0
50.00 D	7.80	1.76	32.00	26.0
100.00 D	7.80	.51	32.00	26.0

Day: 2 Time: 1400

0.00 D	7.90	4.40	32.00	25.5
6.25 D	7.90	4.20	32.00	25.5
12.50 D	7.80	3.89	32.00	25.5
25.00 D	7.80	2.40	32.00	25.5
50.00 D	7.80	1.20	32.00	25.5
100.00 D	7.70	1.14	32.00	25.5

Day: 3 Time: 1400

0.00 D	7.90	5.08	32.00	25.0
6.25 D	7.80	4.93	32.00	25.0
12.50 D	7.80	4.54	32.00	25.0
25.00 D	7.80	4.36	32.00	25.0
50.00 D	7.80	2.48	32.00	25.0
100.00 D	7.70	1.30	32.00	25.0

Day: 4 Time: 1400

0.00 D	7.90	5.00	32.00	25.5
6.25 D	7.90	4.95	32.00	25.5
12.50 D	7.90	4.77	32.00	25.5
25.00 D	7.80	4.15	32.00	25.5
50.00 D	7.80	3.56	32.00	25.5
100.00 D	7.90	2.28	32.00	25.5

WATER QUALITY

Test Number: 0000004028

Test Date: 2-May-1995

Source:

Container	pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
-----------	----	----	-------	------	------	------	-----	-----	-------	-----	----

Concentration: 0.00 D

Day/Time		pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
0	1400	8.00	6.29	32.00	25.5							
1	1400	7.90	4.80	32.00	26.0							
2	1400	7.90	4.40	32.00	25.5							
3	1400	7.90	5.08	32.00	25.0							
4	1400	7.90	5.00	32.00	25.5							

Concentration: 6.25 D

Day/Time		pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
0	1400	8.00	6.17	32.00	25.5							
1	1400	7.70	3.98	32.00	26.0							
2	1400	7.90	4.20	32.00	25.5							
3	1400	7.80	4.93	32.00	25.0							
4	1400	7.90	4.95	32.00	25.5							

Concentration: 12.50 D

Day/Time		pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
0	1400	8.00	6.11	32.00	25.5							
1	1400	7.70	1.82	32.00	26.0							
2	1400	7.80	3.89	32.00	25.5							
3	1400	7.80	4.54	32.00	25.0							
4	1400	7.90	4.77	32.00	25.5							

WATER QUALITY

Test Number: 0000004028

Test Date: 2-May-1995

Source:

Container	pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
-----------	----	----	-------	------	------	------	-----	-----	-------	-----	----

Concentration: 25.00 0

Day/Time		pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
0	1400	8.00	6.29	32.00	25.5							
1	1400	7.80	2.00	32.00	25.0							
2	1400	7.80	2.40	32.00	25.5							
3	1400	7.80	4.36	32.00	25.0							
4	1400	7.80	4.15	32.00	25.5							

Concentration: 50.00 0

Day/Time		pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
0	1400	7.90	6.05	32.00	25.5							
1	1400	7.80	1.76	32.00	26.0							
2	1400	7.80	1.20	32.00	25.5							
3	1400	7.80	2.48	32.00	25.0							
4	1400	7.80	3.56	32.00	25.5							

Concentration: 100.00 0

Day/Time		pH	DO	Salin	Temp	Cond	Hard	Alk	NH3	Chlor	SO3	TS
0	1400	7.90	6.31	32.00	25.5							
1	1400	7.80	.51	32.00	26.0							
2	1400	7.70	1.14	32.00	25.5							
3	1400	7.70	1.30	32.00	25.0							
4	1400	7.90	2.28	32.00	25.5							

y > hour=96

Iteration	Intercept	Slope	Mu	Sigma
0	5.58164339	-0.01818490	31.98495866	-54.99066801
1	5.71721008	-0.02212948	32.40970527	-45.18857999
2	5.72236353	-0.02236584	32.29761989	-44.71103310
3	5.72238119	-0.02236690	32.29688793	-44.70892691
4	5.72238119	-0.02236690	32.29688791	-44.70892686

Covariance Matrix

	Intercept	Slope
Intercept	0.07172943	-0.00122280
Slope	-0.00122280	0.00004437

Covariance Matrix

	Mu	Sigma
Mu	78.01023950	-18.78825153
Sigma	-18.78825153	177.28496191

Chi-Square = 24.0402 With 10 Degrees Of Freedom  
Probability > Chi-Square = 0.0075


The above covariance matrices have been multiplied by the heterogeneity factor  
Check that large chi-square value is not from systematic variation  
A t value of 2.228092 will be used to compute 95 fiducial limits

Acute Toxicity Bioassays - Mysids  
 Paper Pulp- 02MAY95


By > hour=96

Probability	Dose	95% Fiducial Limits	
		Lower	Upper
0.01	136.30540501	353.36728662	90.33992967
0.02	124.11779830	317.46209422	82.69455563
0.03	116.38515245	294.72500787	77.80020338
0.04	110.56818408	277.64961139	74.08954453
0.05	105.83652897	263.78232972	71.04897342
0.06	101.80914725	251.99766711	68.44239753
0.07	98.27792131	241.68111860	66.14064505
0.08	95.11612963	232.45866672	64.06491331
0.09	92.24060659	224.08496697	62.16336706
0.10	89.59368292	216.38999232	60.39996091
0.15	78.63471188	184.70079552	52.92893474
0.20	69.92486956	159.79560725	46.71082084
0.25	62.45260079	138.74550203	41.05990551
0.30	55.74227258	120.23019582	35.59687034
0.35	49.52415311	103.57891064	30.02864751
0.40	43.62376515	88.46445861	24.05895696
0.45	37.91507125	74.78527420	17.33898846
0.50	32.29688791	62.59163362	9.45690191
0.55	26.67870458	51.97407849	-0.00127007
0.60	20.97001067	42.90299739	-11.32934188
0.65	15.06962272	35.13626997	-24.64675705
0.70	8.85150325	28.28395750	-40.01395259
0.75	2.14117504	21.92418460	-57.53252107
0.80	-5.33109373	15.64285024	-78.05220725
0.85	-14.04093605	8.97044287	-102.50310205
0.90	-24.99990709	1.15855432	-133.85143647
0.91	-27.64683077	-0.66367864	-141.48758431
0.92	-30.52235381	-2.62177039	-149.80473856
0.93	-33.68414548	-4.75215434	-158.97253823
0.94	-37.21537143	-7.10710827	-169.23588530
0.95	-41.24275314	-9.76598129	-180.96825077
0.96	-45.97440825	-12.85869397	-194.78339088
0.97	-51.79137662	-16.62251674	-211.80562344
0.98	-59.52402247	-21.57330935	-234.48626943
0.99	-71.71162919	-29.28444588	-270.32569934

ACUTE TOXICITY BIOASSAYS - MYSIDS  
 PAPER PULP- 82MAY85  
 BY-> HOUR=96



ACUTE TOXICITY BIOASSAYS - MYSIDS  
 PAPER PULP- 82MAY85  
 BY-> HOUR=96



RAW DATA, REFER TO FIG. 5

---

TEST DATE	TEST NUMBER
Start: 2-May-95	0000004029

---

SPECIES: Mysidopsis bahia  
TEST MATERIAL: Paper Pulp  
SOURCE: David Taylor  
PROTOCOL: EPA Acute, 4th addition, 1991

---

TEST TYPE: (96) hr acute ( ) chronic  
REFERENCE TOXICANT TEST:

---

=====

MYSID TEST DATA

=====

Test Number: 0000004029 ( ) Chronic (x) Acute 96 hours  
Test Date: 2-May-95  
Source: Test Material: BPP (%)

---

Conc	Rep No.	Cont.	Start	Daily Survival						Prop Alive	Females w/eggs	Prop w/eggs	Weight /Mysid
				1	2	3	4	5	6 End				
0.000	1	11	10	10	8	8	8		.80				
0.000	2	1	10	10	10	10	10		1.00				
6.250	1	10	10	9	9	9	8		.80				
6.250	2	8	10	7	6	6	6		.60				
12.500	1	2	10	7	7	7	7		.70				
12.500	2	3	10	3	3	3	3		.30				
25.000	1	7	10	9	5	5	5		.50				
25.000	2	6	10	10	6	6	5		.50				
50.000	1	9	10	9	7	1	1		.10				
50.000	2	12	10	9	7	1	0		0.00				
100.000	1	4	10	7	0	0	0		0.00				
100.000	2	5	10	3	1	1	1		.10				

---

y > hour=96

Iteration	Intercept	Slope	Mu	Sigma
0	5.49634416	-0.02144678	23.14305860	-46.62703879
1	5.67416007	-0.02972672	22.67858761	-33.63976684
2	5.71033013	-0.03218082	22.07308637	-31.07440547
3	5.71269129	-0.03235435	22.02768199	-30.90774671
4	5.71270117	-0.03235509	22.02748026	-30.90703555
5	5.71270117	-0.03235509	22.02748025	-30.90703553

Covariance Matrix

	Intercept	Slope
Intercept	0.09009237	-0.00211632
Slope	-0.00211632	0.00010545

Covariance Matrix

	Mu	Sigma
Mu	45.87520930	-6.09755290
Sigma	-6.09755290	96.22445449

Chi-Square = 26.9908 With 10 Degrees Of Freedom  
Probability > Chi-Square = 0.0026


The above covariance matrices have been multiplied by the heterogeneity factor  
Check that large chi-square value is not from systematic variation  
 $\tau$  value of 2.228092 will be used to compute 95 fiducial limits

Acute Toxicity Bioassays - Mysids  
 Paper Pulp- 02MAY95


By > hour=96

Probability	Dose	95% Fiducial Limits	
		Lower	Upper
0.01	93.92799676	274.42628271	61.19330151
0.02	85.50277119	245.92433760	55.98915149
0.03	80.15723559	227.87828184	52.64977018
0.04	76.13599731	214.32785927	50.11276116
0.05	72.86503013	203.32493766	48.02979800
0.06	70.08092327	193.97590324	46.24068903
0.07	67.63980618	185.79287321	44.65774612
0.08	65.45407696	178.47891284	43.22744246
0.09	63.46624368	171.83925436	41.91454262
0.10	61.63643989	165.73893313	40.69452089
0.15	54.06056341	140.63301606	35.49226269
0.20	48.03949726	120.93127029	31.10603281
0.25	42.87395891	104.31673381	27.05523215
0.30	38.23514588	89.75424959	23.05960516
0.35	33.93659405	76.73112954	18.88586849
0.40	29.85768827	65.01565281	14.28320809
0.45	25.91129953	54.55928964	8.95158399
0.50	22.02748025	45.42006758	2.55313518
0.55	18.14366098	37.64520968	-5.20967781
0.60	14.19727224	31.14270566	-14.49516106
0.65	10.11836646	25.65663629	-25.32722881
0.70	5.81981463	20.85761398	-37.72506322
0.75	1.18100160	16.42749441	-51.85305486
0.80	-3.98453675	12.07181641	-68.16271401
0.85	-10.00560290	7.46593770	-87.64481094
0.90	-17.58147938	2.09878655	-112.58583505
0.91	-19.41128317	0.85029286	-118.65768432
0.92	-21.39911645	-0.48997984	-125.26996994
0.93	-23.58484567	-1.94674493	-132.55746888
0.94	-26.02596277	-3.55545202	-140.71473474
0.95	-28.81006962	-5.36989252	-150.03843764
0.96	-32.08103680	-7.47811743	-161.01609750
0.97	-36.10227508	-10.04088958	-174.54075693
0.98	-41.44781068	-13.40762889	-192.55945470
0.99	-49.87303625	-18.64366537	-221.02951334

ACUTE TOXICITY BIOASSAYS - MYSIDS  
PAPER PULP- 02MAY95  
BY-> HOUR=96



ACUTE TOXICITY BIOASSAYS - MYSIDS  
PAPER PULP- 02MAY95  
BY-> HOUR=96



RAW DATA, REFER TO FIG. 6

TEST DATE

TEST NUMBER

Start: 2-May-95

0000004030

SPECIES: Mysidopsis bahia

TEST MATERIAL: Paper Pulp

SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

TEST TYPE: (96) hr acute ( ) chronic

REFERENCE TOXICANT TEST:

MYSID TEST DATA

Test Number: 0000004030

( ) Chronic (x) Acute 96 hours

Test Date: 2-May-95

Source:

Test Material: BPP (%)

Conc	Rep No.	Cont.	Start	Daily Survival						Proc	Females	Prop	Weight	
				1	2	3	4	5	6	End	Alive	w/eggs	w/eggs	/Mysid
0.000	1	1	10	10	9	9	9				.90			
0.000	2	2	10	10	10	9	9				.90			
6.250	1	9	10	7	7	7	7				.70			
6.250	2	5	10	7	7	7	7				.70			
12.500	1	3	10	0	0	0	0				0.00			
12.500	2	7	10	0	0	0	0				0.00			
25.000	1	8	10	1	1	1	1				.10			
25.000	2	12	10	5	0	0	0				0.00			
50.000	1	11	10	4	0	0	0				0.00			
50.000	2	4	10	4	1	1	1				.10			
100.000	1	6	10	7	4	0	0				0.00			
100.000	2	10	10	6	0	0	0				0.00			

Acute Toxicity Bioassays - Mysids  
Paper Pulp - 02MAY95

by > hour=96

Iteration	Intercept	Slope	Mu	Sigma
0	5.96830788	-0.05441196	17.79586572	-18.37831337
1	5.49918111	-0.06142942	8.12609237	-16.27884590
2	5.59056440	-0.07346053	8.03920681	-13.61275225
3	5.60256892	-0.07519683	8.01322208	-13.29843242
4	5.60270301	-0.07521886	8.01265804	-13.29453797
5	5.60270303	-0.07521886	8.01265793	-13.29453735

Covariance Matrix

	Intercept	Slope
Intercept	0.68978369	-0.03324949
Slope	-0.03324949	0.00319034

Covariance Matrix

	Mu	Sigma
Mu	63.94239963	18.06106537
Sigma	18.06106537	99.66198435

Chi-Square = 141.6567 With 10 Degrees Of Freedom  
Probability > Chi-Square = 0.0000


The above covariance matrices have been multiplied by the heterogeneity factor  
Check that large chi-square value is not from systematic variation  
t value of 2.228092 will be used to compute 95 fiducial limits

Acute Toxicity Bioassays - Mysids  
 Paper Pulp - 02MAY95


By > hour=96

Probability	Dose	95% Fiducial Limits	
		Lower	Upper
0.01	38.94037668	-28.57590548	17.72027114
0.02	35.31629970	-22.60899271	15.78168029
0.03	33.01693901	-18.69003445	14.41856319
0.04	31.28721949	-15.63279992	13.28398843
0.05	29.88022608	-13.04111278	12.25623673
0.06	28.68265368	-10.72374405	11.27002270
0.07	27.63261696	-8.56274231	10.27618324
0.08	26.69243430	-6.46347563	9.22197119
0.09	25.83737579	-4.31659064	8.02552157
0.10	25.05029303	-1.90755802	6.49136626
0.15	21.79156015	4.10301701	4.10301701
0.20	19.20162270	5.54243205	5.54243205
0.25	16.97968710	6.77732192	6.77732192
0.30	14.98432028	7.88629117	7.88629117
0.35	13.13531546	8.91391651	8.91391651
0.40	11.38079050	9.88903257	9.88903257
0.45	9.68326721	10.83246869	10.83246869
0.50	8.01265793	11.76094676	11.76094676
0.55	6.34204865	12.68942483	12.68942483
0.60	4.64452537	13.63286095	13.63286095
0.65	2.89000041	14.60797701	14.60797701
0.70	1.04099558	15.63560235	15.63560235
0.75	-0.95437123	16.74457160	16.74457160
0.80	-3.17630683	17.97946147	17.97946147
0.85	-5.76624429	11.72484946	27.11290356
0.90	-9.02497716	8.59645034	33.86352846
0.91	-9.81205992	8.00685171	35.32800441
0.92	-10.66711844	7.40620330	36.87908819
0.93	-11.60730109	6.78389359	38.54645252
0.94	-12.65733782	6.12654858	40.37095981
0.95	-13.85491021	5.41550567	42.41315743
0.96	-15.26190363	4.62176537	44.77083317
0.97	-16.99162314	3.69402456	47.62123375
0.98	-19.29098383	2.52292754	51.34817193
0.99	-22.91506082	0.77973022	57.11969116

ACUTE TOXICITY BIOASSAYS - MYSIDS  
PAPER PULP - 02MAY95  
BY-> HOUR=96



ACUTE TOXICITY BIOASSAYS - MYSIDS  
PAPER PULP - 02MAY95  
BY-> HOUR=96



RAW DATA, REFER TO FILE 7

---

TEST DATE	TEST NUMBER
Start: 9-May-95	0000004031

---

SPECIES: Menidia beryllina  
TEST MATERIAL: Paper Pulp  
SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

---

TEST TYPE: (96) hr acute ( ) chronic  
REFERENCE TOXICANT TEST: 0000004034

---

=====

FISH TEST DATA

=====

Test Number: 0000004031 ( ) Chronic (x) Acute 96 hours  
Test Date: 9-May-95  
Source: Test Material: BPP (%)

---

Conc	Rep	Cont. No.	Start	Daily Survival						Prod Alive	Weight /Fish
				1	2	3	4	5	6 End		
0.00	D	1	11	10	10	10	10	9		.90	
0.00	D	2	12	10	10	10	10	10		1.00	
6.25	D	1	3	10	10	10	10	10		1.00	
6.25	D	2	7	10	10	10	10	10		1.00	
12.50	D	1	8	10	10	10	10	10		1.00	
12.50	D	2	6	10	9	9	9	9		.90	
25.00	D	1	10	10	10	10	10	10		1.00	
25.00	D	2	2	10	10	10	10	10		1.00	
50.00	D	1	5	10	10	10	10	10		1.00	
50.00	D	2	1	10	10	10	10	10		1.00	
100.00	D	1	9	10	10	10	10	10		1.00	
100.00	D	2	4	10	10	9	9	9		.90	

---

RAW DATA, REFER TO FIG. 8

TEST DATE

TEST NUMBER

Start: 9-May-95

0000004032

SPECIES: Menidia beryllina

TEST MATERIAL: Paper Pulp

SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

TEST TYPE: (96) hr acute ( ) chronic

REFERENCE TOXICANT TEST: 0000004034

FISH TEST DATA

Test Number: 0000004032

( ) Chronic (x) Acute 96 hours

Test Date: 9-May-95

Source:

Test Material: BPP (%)

Conc	Rep	Cont. No.	Start	Daily Survival						Prop Weight /Fish
				1	2	3	4	5	6 End	
0.00	D	1	6	10	10	10	10	10	1.00	
0.00	D	2	5	10	10	10	10	10	1.00	
6.25	D	1	9	10	10	10	10	10	1.00	
6.25	D	2	1	10	10	10	10	10	1.00	
12.50	D	1	2	10	10	10	10	10	1.00	
12.50	D	2	8	10	10	10	10	10	1.00	
25.00	D	1	12	10	10	10	10	10	1.00	
25.00	D	2	11	10	10	10	10	10	1.00	
50.00	D	1	3	10	10	9	7	6	.60	
50.00	D	2	4	10	10	10	10	10	1.00	
100.00	D	1	10	10	10	10	10	10	1.00	
100.00	D	2	7	10	10	9	9	9	.90	

RAW DATA, REFER TO FIG. 9

TEST DATE

TEST NUMBER

Start: 9-May-95

0000004033

SPECIES: Menidia beryllina

TEST MATERIAL: Paper Pulp

SOURCE:

PROTOCOL: EPA Acute, 4th addition, 1991

TEST TYPE: (96) hr acute ( ) chronic

REFERENCE TOXICANT TEST: 0000004034

FISH TEST DATA

Test Number: 0000004033

( ) Chronic (x) Acute 96 hours

Test Date: 9-May-95

Source:

Test Material: BPP (%)

Conc	Rep	Cont. No.	Start	Daily Survival						Prop Weight /Fish
				1	2	3	4	5	6 End	
0.90	D	1	10	10	10	10	10		1.00	
0.00	D	2	3	10	10	10	10		1.00	
6.25	D	1	2	10	9	9	9	8	.80	
6.25	D	2	11	10	10	10	10	9	.90	
12.50	D	1	8	10	10	10	10	10	1.00	
12.50	D	2	1	10	10	10	10	10	1.00	
25.00	D	1	6	10	10	10	10	10	1.00	
25.00	D	2	5	10	10	10	10	10	1.00	
50.00	D	1	4	10	10	10	9	9	.90	
50.00	D	2	12	10	10	10	10	10	1.00	
100.00	D	1	7	10	10	10	10	10	1.00	
100.00	D	2	9	10	10	10	10	10	1.00	







RELATIVE FLUORESCENCE DATA

test species Skeletonema costatum  
 toxicant: ~~4 Liters~~ Paper Pulp

cooperator(s) 6 2TEN

test date 30 MAY 95 - 3 JUN 95

rep	hour 01			hour 24			hour 48			hour 72			hour 96		
	coor used 10x			coor used 10x			coor used 3x			coor used 1x			coor used 1x		
	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C
1	23.6	23.3	23.9	65.7	66.6	66.7	67.0	64.9	67.3	58.0	56.1	56.6	26.9	27.7	27.2
2	23.9	23.9	24.1	65.9	65.0	67.5	64.7	62.8	64.3	46.5	44.3	44.9	17.9	17.8	18.9
3	24.5	25.3	25.0	66.2	67.0	66.5	69.0	70.0	69.2	53.1	52.6	53.1	20.6	20.9	21.1
4	23.7	23.8	25.0	69.5	72.3	66.8	76.0	76.0	76.0	58.3	59.3	58.6	22.5	23.6	24.4
5	25.4	24.0	23.9	68.4	71.6	65.0	70.6	71.7	70.2	60.1	58.7	57.9	24.6	27.6	25.2
6	27.8	24.4	25.7	73.2	74.5	73.9	89.2	86.9	86.7	74.2	74.3	74.7	30.3	30.4	30.2
7	23.3	25.3	24.6	69.6	69.4	69.6	75.4	73.4	76.5	64.7	65.9	65.1	22.4	22.7	22.0
8	23.3	23.4	24.6	68.4	69.4	69.0	82.5	80.3	78.0	63.0	63.5	62.5	26.4	25.6	27.2
9	24.4	23.0	24.6	65.7	68.9	68.2	83.3	82.9	82.9	77.5	76.5	76.6	35.8	35.2	34.6
10	25.2	24.4	25.5	68.1	69.5	67.6	77.1	74.3	74.9	72.1	72.1	67.5	32.6	31.8	31.7
11	24.5	27.0	26.1	74.0	77.8	69.1	78.5	78.0	76.6	68.9	68.4	68.3	28.3	29.5	29.6
12	26.3	24.5	25.9	69.3	67.3	71.2	78.4	75.9	78.7	71.7	70.9	71.4	30.4	31.2	32.2
13	24.3	26.8	28.4	69.7	70.3	70.5	78.0	71.9	95.3	82.0	80.1	80.6	35.8	34.2	33.7
14	27.8	23.3	24.1	65.7	68.4	67.0	82.1	82.6	83.5	76.4	76.3	76.0	34.9	30.9	35.8
15	26.6	25.3	23.5	66.6	64.4	61.3	66.4	68.7	66.6	62.6	61.6	61.4	29.6	28.3	29.3
16	23.7	25.3	24.7	66.7	67.7	66.1	72.0	78.3	79.4	68.7	68.4	68.4	31.9	30.9	30.8
17	24.4	26.5	26.4	64.4	68.0	65.5	83.0	83.1	80.5	74.3	71.4	73.9	30.7	32.0	32.2
18	25.4	27.0	27.1	70.6	66.7	68.4	86.7	88.0	84.1	76.6	74.7	75.7	38.7	32.5	30.5
19															
20															
21															
22															
23															
24															

Right (red)

RELATIVE FLUORESCENCE DATA

test species: Skeletonema costatum

observer(s) G. RUSSEN

toxicant: ~~2,4-D~~ Metal/Ascorb. (-)

test date 20 MAR 75 - 5 JUN 75

rep	hour 01			hour 24			hour 48			hour 72			hour 96			rep
	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	
1	22.5	23.3	22.3	54.5	51.1	52.0	18.6	18.6	18.9	10.9	10.6	10.2	5.7	6.2	6.2	1
2	23.8	23.5	23.0	71.5	68.6	70.7	82.4	83.2	81.6	80.4	79.6	79.7	35.5	36.4	36.4	2
3	23.4	21.7	23.5	62.1	63.1	67.0	69.4	68.1	68.1	67.0	66.0	65.9	31.3	31.3	31.1	3
4	23.5	23.2	24.7	66.2	65.6	62.9	<del>74.3</del> 70.3	<del>73.2</del> 72.3	<del>95.6</del> 89.9	99.2	96.7	98.3	44.0	43.7	41.9	4
5	22.5	24.4	24.8	62.2	62.0	58.8	49.6	47.9	46.0	56.4	55.8	55.1	27.5	28.5	28.5	5
6	22.4	23.3	24.4	72.3	68.1	69.5	103.4	100.5	103.6	108.6	107.7	108.6	46.4	47.2	46.2	6
7	24.6	24.1	23.5	65.9	66.4	66.0	97.1	94.5	95.2	94.5	92.6	95.5	37.9	41.4	41.7	7
8	22.8	23.2	23.0	68.5	57.4	64.3	98.3	88.6	89.7	104.1	90.5	96.7	41.5	43.9	45.1	8
9	24.1	23.0	23.1	69.9	66.3	67.8	104.1	94.2	102.1	105.9	94.7	97.3	40.2	40.8	40.8	9
10	26.1	31.4	24.6	42.9	42.7	37.5	73.9	73.3	70.3	73.9	73.3	72.2	36.1	35.0	34.6	10
11	25.3	25.0	25.5	53.1	46.0	42.3	78.4	70.1	76.1	79.3	76.5	78.7	30.2	31.5	29.5	11
12	24.8	25.1	24.5	46.9	43.4	45.0	78.1	67.1	74.5	77.9	74.8	76.6	28.6	28.0	30.4	12
13	25.0	23.6	27.2	34.2	34.2	33.7	22.3	23.6	22.6	11.3	11.4	11.1	3.5	3.9	3.8	13
14	28.0	27.4	27.3	21.0	17.1	17.3	13.4	14.9	13.3	10.1	10.6	9.8	3.3	3.5	3.4	14
15	29.0	27.5	26.2	21.3	19.0	18.4	17.4	15.1	16.1	10.2	10.4	10.9	4.4	5.0	4.9	15
16	32.6	33.2	<del>32.0</del>	15.7	15.4	15.5	3.2	3.1	2.7	0.8	1.0	0.7	0.4	0.1	0.5	16
17	32.9	33.8	32.6	11.5	12.5	10.7	2.5	2.3	2.7	0.5	0.7	0.5	0.0	0.0	0.0	17
18	35.2	34.5	33.7	17.9	21.7	17.5	3.6	3.3	3.2	0.9	0.9	0.8	0.1	0.2	0.1	18
19																19
20																20
21																21
22																22
23																23
24																24

Left (yellow)



DATE: 8 APRIL RUN: 964A FILTER TYPE: 100-2 STIRRER: 500 cc 7.30  
 EXPERIMENTER: L. ADOTA  
 TEST ARTICLE OR CHEMICAL: 7.20% PAPER PULP (5%) (achate)  
 DARK COUNTS: \_\_\_\_\_ AVG = 6037  
 TIME RUN STARTED 1635 TIME RUN ENDED \_\_\_\_\_ 7 OF 5 RUNS

CONCENTRATIONS OR DILUTIONS

REP#	CONTRAC	6.25% (0.28%)	12.5% (0.57%)	25% (1.14%)	50% (2.28%)	100% (4.55%)
1	220,886	3,453,725	823,774	225,936	457,685	13,432
2	2,678,115	25,024	2,592	948	22,494	97,352
3	512,455	60,987	2,199,922	2,241	88,622	80,560
4	1,359,221	2,724,236	65,856	31,326	172,244	32,525
5	139,004	241,019	179,571	14,456	212,322	16,227
MEAN:	1,394,952	1,210,870	657,344	23,021	243,213	36,019
S.D.	1,562,099	1,317,859	919,166	119,412	129,687	35,052
C.V.	112	109	141	164	57.43	97.32
% of Control:	86.80	96.84	5.22	17.44	2.58	
Net change:						

Fig. 16



DATE: 11 Nov 91 RUH: 9611C FILTER TYPE: MC 2 STIRRER: 30 sec  
7.31 V  
 EXPERIMENTER: U.V. V. V. A.  
 TEST METHOD OR CIRCULAR: METAL JAMS (25%)  
 DARK COUNTS: \_\_\_\_\_ AVG #: \_\_\_\_\_

TIME RUN BEG. \_\_\_\_\_ TIME RUN ENDED \_\_\_\_\_

CONCENTRATIONS OR DILUTIONS

REP#	CONTR	6.25	12.5	25	50	76	bulk
1	2405920	2543933	1792915	696349	4036	64593	297918
2	2195842	2114238	1909102	400650	3349	132499	371625
3	2245167	2018477	1580464	269994	3587	203711	558991
4	1781792	2096956	1143482	293638	3324	73067	323973
5	1833353	1964449	1586971	269863	3575	77197	558539
MEAN	2092735	2153615	1662598	446099	3514	73545	422209
S.D.			163362	255		25168	113426
C.V.							
% of Control:	1.03	7914	.21	0.00	05.27	.20	
Net change:							

CC50 ~~18.83%~~ 18.83%

DATE: 6/25/45 - RUN: 16 HOURS FILTER TYPE: N/P - 2 STIRRER: 23.22.525  
 30565

EXPERIMENT: L2P07A

TEST ARTICLE OR CHEMICAL: SHIPPED METAL 5%

DARK COURTS: 2107 3056 3315 AVG = 2526

TIME RUN STARTED 1500 TIME RUN ENDED 1547

CONCENTRATIONS OR DILUTIONS

REF#	CONTROL	6.25%	12.5%	25%	50%	100%
1	1,201,825	21,358	1,335,589	4014	281,845	72,420
2	1,327,161	1,237,997	853,004	158,385	6937	123,143
3	1,261,235	1,386,031	1,234,220	11,821	13,822	2904
4	1,951,295	1,651,240	1,670,122	135,713	65,992	287,089
5	1,181,642	717,448	1,650,309	408,769	25,333	120,040
MEAN	1,187,041	1,082,292	1,325,658	143,956	78,596	144,519
S.D.	243,798	714,350	330,968	163,794	115,960	109,063
C.V.	66.39	66.07	24.84	11.378	147.54	72.07

% of control: 23.68 87.62 9.68 5.29 9.52

Net change:

IC50 = 18.64%

Microtox Test  
Paper Pulp - 05/03/95

5% Leachate

5 minute readings	rep	Control	12.5%	25%	50%	100%
Trial 1	1	93.5	83.0	75.5	69.5	46.0
	2	68.0	73.0	71.0	62.0	48.0
	3	<u>52.0</u>	<u>83.0</u>	<u>62.5</u>	<u>30.5</u>	<u>51.0</u>
	Mean	<u>71.2</u>	<u>79.7</u>	<u>69.7</u>	<u>77.3</u>	<u>49.3</u>
Trial 2	1	79.0	90.1	82.0	92.5	62.5
	2	96.5	102.0	84.0	81.0	85.0
	3	<u>87.0</u>	<u>85.5</u>	<u>99.0</u>	<u>72.5</u>	<u>61.0</u>
	Mean	<u>87.5</u>	<u>94.8</u>	<u>88.3</u>	<u>82.0</u>	<u>69.5</u>
Trial 3	1	-	103.5	83.0	76.0	81.0
	2	92.0	92.0	85.0	103.5	78.5
	3	<u>87.0</u>	<u>86.0</u>	<u>95.5</u>	<u>92.5</u>	<u>97.0</u>
	Mean	<u>89.5</u>	<u>93.8</u>	<u>87.8</u>	<u>90.7</u>	<u>85.5</u>

15 minute readings

Trial 1	1	96.0	85.0	79.5	69.5	50.0
	2	63.0	58.5	70.0	62.5	43.0
	3	<u>50.0</u>	<u>83.5</u>	<u>61.5</u>	<u>61.5</u>	<u>53.0</u>
	Mean	<u>69.7</u>	<u>75.7</u>	<u>70.3</u>	<u>77.8</u>	<u>50.3</u>
Trial 2	1	75.0	90.0	83.0	91.0	63.5
	2	95.0	99.5	81.0	80.0	81.0
	3	<u>83.0</u>	<u>83.0</u>	<u>97.0</u>	<u>72.0</u>	<u>58.5</u>
	Mean	<u>84.3</u>	<u>90.8</u>	<u>87.0</u>	<u>81.0</u>	<u>67.7</u>
Trial 3	1	-	99.5	73.0	68.5	72.5
	2	81.0	86.5	76.0	97.0	68.0
	3	<u>77.0</u>	<u>74.5</u>	<u>88.5</u>	<u>61.5</u>	<u>92.5</u>
	Mean	<u>79.0</u>	<u>86.8</u>	<u>79.2</u>	<u>82.3</u>	<u>77.7</u>

Calculated/Graphed values

	5 minute	15 minute
Trial 1	EC20 = 76% EC50 > 100% reduction = 31%	Inconclusive  reduction = 28%
Trial 2	EC20 = 98% EC50 = 100% reduction = 21%	EC20 > 100% EC50 > 100% reduction = 20%
Trial 3	No toxicity reduction = 4%	No toxicity reduction = 2%

reduction = percent reduction in light output at 100% leachate.

Microtox Test  
Metal leachates - 06/01/95

Centrifuged	rep	Control	12.5%	25%	50%	100%
5 minute	1	69.5	60.0	63.0	58.5	53.0
	2	72.0	74.0	62.5	57.0	44.0
	3	<u>95.0</u>	<u>71.5</u>	<u>63.5</u>	<u>58.0</u>	<u>52.0</u>
	<u>Mean</u>	<u>78.8</u>	<u>68.5</u>	<u>63.0</u>	<u>57.8</u>	<u>49.7</u>
15 minute	1	61.0	64.5	60.5	55.0	50.5
	2	64.0	71.5	64.5	55.0	43.0
	3	<u>92.0</u>	<u>71.0</u>	<u>61.5</u>	<u>56.5</u>	<u>50.5</u>
	<u>Mean</u>	<u>72.3</u>	<u>69.0</u>	<u>62.2</u>	<u>55.5</u>	<u>48.0</u>
Uncentrifuged						
5 minute	1	96.0	103.0	95.0	87.5	73.5
	2	96.5	103.0	100.0	82.5	78.5
	3	<u>97.5</u>	<u>103.0</u>	<u>103.0</u>	<u>90.0</u>	<u>70.0</u>
	<u>Mean</u>	<u>96.7</u>	<u>103.0</u>	<u>99.5</u>	<u>86.7</u>	<u>74.0</u>
15 minute	1	97.0	99.5	92.0	85.5	71.0
	2	99.0	103.0	95.0	89.5	74.5
	3	<u>99.5</u>	<u>103.0</u>	<u>103.0</u>	<u>89.0</u>	<u>70.0</u>
	<u>Mean</u>	<u>96.5</u>	<u>101.8</u>	<u>96.7</u>	<u>88.0</u>	<u>71.8</u>

Calculated/Graphed Values

	5 minute	15 minute
Centrifuged	EC20 = 27.5% EC50 > 100% reduction = 37%	EC20 = 46% EC50 > 100% reduction = 37%
Uncentrifuged	EC20 = 89% EC50 > 100% reduction = 23%	EC20 = 80% EC50 > 100% reduction = 27%

reduction = percent reduction in light output at 100% leachate.

Microtox Tests  
Paper Pulp - 05/03/95  
Metal Leachate - 06/01/95

The tests conducted on 05/03/95 tested the first paper pulp leachate used. Three trials were performed with four dilutions and a control. Five minute and 15-minute readings were taken. Both the EC20 and EC50 were determined graphing the calculated Microtox statistic,  $\bar{y}$ , on log/log paper. Also, the percent reduction of light output at the 100% leachate dilution was calculated. All results are reported on the attached page.

Trial 1 and 2 showed a 5-minute EC20 of 76% and 98%, respectively. The five minute EC50 in both those trials were at or exceeded 100%, the maximum dilution tested. The third trial showed no toxicity as the control and the 100% leachate reading were essentially the same. The 15-minute readings on trial 1 were inconclusive for determining EC values because all mean readings for the dilutions except 100% exceeded the control mean. This yielded only one usable point, and a dose response curve could not be plotted. The 15-minute EC20 and EC50 for trial 2 both exceeded the 100% dilution.

The tests conducted on 06/01/95 tested the metal leachate both centrifuged and uncentrifuged. Four dilutions and a control were tested, and 5 and 15-minute readings were taken. Only one trial of each were performed. The same calculations described above were made and are reported on the attached page.

The centrifuged sample appeared more toxic than the uncentrifuged, but this may be some effect of interference from the poor clarity of the uncentrifuged sample. The 5 and 15 minute EC20 for the centrifuged sample are 27.5 and 46%, respectively. Both EC50 values exceeded 100% leachate. The 5 and 15-minute EC20 values for the uncentrifuged sample were 69 and 80%, respectively. The EC50 values for this sample both exceeded 100%.

Microtox Test  
Metal and paper pulp leachates - 06/27/95

5% Metal Leachate	rep	Control	12.5%	25%	50%	100%
5 minute	1	88.5	74.5	99.5	66.0	63.
	2	78.0	78.0	71.0	72.0	68.
	3	<u>87.0</u>	<u>75.5</u>	<u>68.0</u>	<u>67.0</u>	63.
	Mean	<u>84.5</u>	<u>76.0</u>	<u>68.5</u>	<u>68.3</u>	65.
15 minute	1	100.0	85.0	82.0	80.5	82.
	2	90.5	92.0	88.0	91.0	89.
	3	<u>100.0</u>	<u>89.0</u>	<u>82.0</u>	<u>84.0</u>	80.
	Mean	<u>96.8</u>	<u>88.9</u>	<u>84.0</u>	<u>85.2</u>	83.
0.01% Paper Pulp Leachate						
5 minute	1	94.0	87.0	76.5	72.0	75.
	2	86.5	81.0	78.0	74.0	70.
	3	<u>90.0</u>	<u>84.0</u>	<u>78.5</u>	<u>73.0</u>	71.
	Mean	<u>90.2</u>	<u>84.0</u>	<u>77.7</u>	<u>73.0</u>	72.
15 minute	1	92.5	86.0	73.0	63.5	71.
	2	84.5	78.00	75.0	69.5	63.
	3	<u>89.5</u>	<u>80.0</u>	<u>77.0</u>	<u>67.0</u>	67.
	Mean	<u>88.8</u>	<u>81.3</u>	<u>75.0</u>	<u>68.3</u>	68.

Calculated/Graphed Values

	5 minute	15 minute
5% Metal Leachate	EC20 = 44% EC50 > 100% reduction = 23%	EC20 > 100% EC50 > 100% reduction = 13%
0.01% Paper Pulp	EC20 = 90% EC50 > 100% reduction = 20%	EC20 = 60% EC50 > 100% reduction = 23%

reduction = percent reduction in light output at 100% leachate.

## **APPENDIX D**

### **SOLID PHASE BENTHIC ORGANISM TOXICITY SCREENING REPORT**

Source: Toxicity Testing of Paper Pulp Slurry on Benthic Organisms.  
A Report Submitted to: NCCOSC RDTE DIV CODE 522  
San Diego, California  
Coastal Resources Associates, Inc., 1995

A Report Submitted to:  
NCCOSC RD-T&E Div.  
Code 522  
53475 Stroth Road, Rm 258  
San Diego, CA 92152-6310

June 30, 1995

**Toxicity Testing of Paper Pulp Slurry  
on Benthic Organisms**

Submitted by:  
Coastal Resources Associates, Inc.  
1185 Park Center Dr., Suite A  
Vista, CA 92083

Study Director:  
Thomas A. Dean, Ph.D.

A handwritten signature in cursive script, appearing to read "Thomas A. Dean", written over a horizontal line.

## Table of Contents

Summary.....	1
1.0 Introduction.....	2
2.0 Methods.....	2
3.0 Results.....	4
4.0 Detailed Results - 10 day Amphipod Test.....	5
5.0 Detailed Results - 96 hour Polychaete Test.....	12
6.0 Detailed Results - Sediment Analysis.....	21

# **Toxicity Testing of Paper Pulp Slurry on Benthic Organisms**

## **Summary**

Laboratory toxicity tests were conducted using paper pulp slurry derived from mixed paper and cardboard waste from US ships. Tests were conducted during the months of May and June 1995, using amphipods and polychaetes. The tests were performed to investigate what effect, if any, the paper pulp slurry would have on the benthic community if disposed of into the ocean. Results of the toxicity tests showed no observed effect from the paper pulp slurry. The survival of both amphipods and polychaetes was similar in the control sediments, and in the sediments with paper slurry added.

# Toxicity Testing of Paper Pulp Slurry on Benthic Organisms

## 1.0 Introduction

US ships are seeking to dispose of paper pulp slurry, derived from on board waste of mixed paper and cardboard, into the Baltic Sea, North Sea, Mediterranean Sea, Caribbean Sea, and Antarctic Ocean. Disposal of the paper slurry is proposed at an offshore distance of at least 12 miles and at least 50 meters depth. The slurry would be diluted prior to disposal to obtain a maximum concentration of 2% slurry, and then discharged into the ocean off the ship's wake.

Laboratory tests were conducted to examine the impact the paper slurry would have on the benthic infaunal community under "worst case" conditions. To accomplish this, the concentrations of paper slurry tested were higher than that expected to be found at the ocean floor. The organisms chosen for the laboratory tests were amphipods and polychaetes, which are both important components of most marine benthic communities. The amphipod species used during the tests was *Grandidierella japonica* and the polychaete used was *Neanthes arenaceodentata*.

## 2.0 Methods

A sample of the paper pulp slurry was delivered to the laboratory of Coastal Resources Associates, Inc. by NCCOSC. Tests were performed according to standard protocols for the 10 day amphipod test and the 96 hour polychaete test. The Standard Operating Procedures for these tests have been previously delivered to NCCOSC. The tests were performed in two types of sediment, fine sand and silty sand. Grain size analysis and total organic carbon analysis were also conducted on the sediment types to determine the influence of these variables on toxicity.

Both species were exposed to the paper slurry in the same manner. Sediment and seawater were collected from an unpolluted source and placed in test containers to settle overnight. On test initiation day, the seawater was renewed and the organisms were added to the test containers. The organisms were allowed to burrow for 1 hour before adding the test substance (paper slurry). The test substance was then added and allowed to settle on top of the organisms, simulating what would occur in the real environment.

The concentrations of the paper slurry used in the tests were equivalent to the amount of paper slurry that would settle on top of the sediment. Dilutions were prepared to create layers of paper slurry that were 0.01mm, 0.1mm, and 1.0mm in height. This was accomplished by preparing an initial stock solution of 0.1% (by volume of solid wet material) paper slurry in filtered seawater. The 0.1% stock solution was prepared by making serial dilutions of a 10% solution of the paper slurry. This final stock solution of 0.1% was then allowed to mix well using a stir plate for 1 hour. From prior calculations, enough of the 0.1% stock solution was added to each test container to achieve the desired layers of 0.01mm, 0.1mm, and 1.0mm of settled paper slurry.

After the test was initiated, the amphipods were exposed to the test substance for 10 days and the polychaetes were exposed for 96 hours. During these periods, water quality was monitored daily, and the overlying seawater was renewed every 48 hours without disturbing the sediment or the layer of paper slurry. At the end of the exposure period, the organisms were removed from the sediment and examined for mortality. In the amphipod test, the organisms were also examined to determine whether they would rebury in a new container of sediment and seawater (no test substance), after they were removed from the test container. This additional procedure to the amphipod test is to help determine if there were any sublethal effects on the organisms from the test substance.

### 3.0 Results

Toxicity tests were conducted using paper slurry at concentrations greater than that expected to be found at the ocean floor. No toxicity was observed in the amphipod or polychaete tests.

Results of the 10 day amphipod test showed no significant difference in survival between the controls and the highest concentration tested, a 1mm layer of paper slurry ( $P < 0.05$ , Dunnett's test). A 90 - 100% survival rate was seen in all test concentrations. The paper slurry also showed no effect in the organisms ability to rebury in new sediment after completion of the test ( $P < 0.05$ , Dunnett's test).

Results of the 96 hour polychaete test showed no significant difference in survival between the controls and the highest concentration tested, a 1mm layer of paper slurry ( $P < 0.05$ , Dunnett's test). A 95 - 100% survival rate was seen in all test concentrations.

There was no apparent effect of grain size on the toxicity of the paper slurry. There were no toxic effects of the paper slurry in either sediment sample. Fine sand and silty sand were collected for testing based on a visual examination of sediments in the field. We did not use coarse sand in our tests, because amphipods cannot bury effectively in coarse sediments, and do not survive well. However, grain size analysis indicated that the two samples of sediment used in the polychaete test differed only slightly (see Section 6.0). In the polychaete test, both sediment samples were composed of 96-97% sand and 3-4% silt and clay, but the silty sand sample was somewhat higher in total organic carbon (861 vs 552 mg/kg). In the amphipod test, the fine sediment was composed of 96% sand and 4% silt and clay, while the silty sediment was composed of 92% sand and 8% silt and clay. The total organic carbon was slightly higher in the silty sediment (1190 vs 1040). There were no effects observed from the paper slurry in either sediment sample.

Details of specific test results, along with the physical/chemical data and individual test data are given in Sections 4.0, 5.0, and 6.0 that follow.

## **4.0 Results of the 10 day Amphipod Test**

Table 1. Summary of test information for the 10 day amphipod test using paper pulp slurry.

**Test Information:**

Date and Time of Test Initiation: 13 June 1995, 1700 hours

Concentrations Used:

Fine Sand Test: 0, 0.01, 0.1, and 1.0 mm layer of paper slurry  
Silty Sand Test: 0, 0.01, 0.1, and 1.0 mm layer of paper slurry

Test Material Sources:

Dilution Water: La Jolla, CA.  
Sediment: Newport Bay and Agua Hedionda Lagoon, CA.  
Organisms: Newport Bay, CA.  
Paper Slurry: NRAD

Dates of Collection:

Dilution Water: 6 June and 16 June 1995  
Sediment: 10 June and 12 June 1995  
Organisms: 10 June 1995  
Paper Slurry: 27 April 1995

Table 2. Summary of final test results for the 10 day amphipod test using paper pulp slurry during June 1995. For survival, the NOEC (no observed effect concentration) and ANOVA Mean Square Errors (MSE) are given for analyses of arcsin transformed data. All tabulated means are for untransformed data.

**Fine Sand Test**

CONCENTRATION (mm Paper Slurry)	% SURVIVAL		% REBURIAL	
	MEAN	S.D.	MEAN	S.D.
0	98.0	4.47	90.0	12.25
0.01	100.0	0.00	88.0	26.83
0.1	90.0	12.25	85.4	20.48
1	96.0	5.48	92.0	13.04

**SURVIVAL:**

-----  
 NOEC = 1  
 LOEC = .  
 EC50 = .  
 ANOVA MSE = 92.51

**REBURIAL:**

-----  
 NOEC = 1  
 LOEC = .  
 EC50 = .  
 ANOVA MSE = 340.62

**Silty Sand Test**

CONCENTRATION (mm Paper Slurry)	% SURVIVAL		% REBURIAL	
	MEAN	S.D.	MEAN	S.D.
0	94.0	8.94	68.6	27.40
0.01	98.0	4.47	84.0	35.78
0.1	94.0	8.94	75.5	30.23
1	100.0	0.00	90.0	22.36

**SURVIVAL:**

-----  
 NOEC = 1  
 LOEC = .  
 EC50 = .  
 ANOVA MSE = 97.06

**REBURIAL:**

-----  
 NOEC = 1  
 LOEC = .  
 EC50 = .  
 ANOVA MSE = 572.44

Table 3. Individual test data for survival in the 10 day amphipod test using paper pulp slurry.

		<u>Fine Sand Test</u>		
Conc. Rep			# Alive	# Dead
0	1	. . . . .	10	0
0	2	. . . . .	10	0
0	3	. . . . .	9	1
0	4	. . . . .	10	0
0	5	. . . . .	10	0
0.01	1	. . . . .	10	0
0.01	2	. . . . .	10	0
0.01	3	. . . . .	10	0
0.01	4	. . . . .	10	0
0.01	5	. . . . .	10	0
0.1	1	. . . . .	9	1
0.1	2	. . . . .	9	1
0.1	3	. . . . .	10	0
0.1	4	. . . . .	7	3
0.1	5	. . . . .	10	0
1	1	. . . . .	10	0
1	2	. . . . .	10	0
1	3	. . . . .	10	0
1	4	. . . . .	9	1
1	5	. . . . .	9	1

		<u>Silty Sand Test</u>		
Conc. Rep			# Alive	# Dead
0	1	. . . . .	10	0
0	2	. . . . .	9	1
0	3	. . . . .	10	0
0	4	. . . . .	8	2
0	5	. . . . .	10	0
0.01	1	. . . . .	9	1
0.01	2	. . . . .	10	0
0.01	3	. . . . .	10	0
0.01	4	. . . . .	10	0
0.01	5	. . . . .	10	0
0.1	1	. . . . .	9	1
0.1	2	. . . . .	8	2
0.1	3	. . . . .	10	0
0.1	4	. . . . .	10	0
0.1	5	. . . . .	10	0
1	1	. . . . .	10	0
1	2	. . . . .	10	0
1	3	. . . . .	10	0
1	4	. . . . .	10	0
1	5	. . . . .	10	0

Table 4. Individual test data for reburial in the 10 day amphipod test using paper pulp slurry.

		<u>Fine Sand Test</u>					# able to Rebury	# Not able to Rebury
Conc. Rep								
0	1	.	.	.	.	.	9	1
0	2	.	.	.	.	.	9	1
0	3	.	.	.	.	.	9	0
0	4	.	.	.	.	.	7	3
0	5	.	.	.	.	.	10	0
0.01	1	.	.	.	.	.	10	0
0.01	2	.	.	.	.	.	10	0
0.01	3	.	.	.	.	.	4	6
0.01	4	.	.	.	.	.	10	0
0.01	5	.	.	.	.	.	10	0
0.1	1	.	.	.	.	.	9	0
0.1	2	.	.	.	.	.	9	0
0.1	3	.	.	.	.	.	10	0
0.1	4	.	.	.	.	.	4	3
0.1	5	.	.	.	.	.	7	3
1	1	.	.	.	.	.	10	0
1	2	.	.	.	.	.	9	1
1	3	.	.	.	.	.	7	3
1	4	.	.	.	.	.	9	0
1	5	.	.	.	.	.	9	0

		<u>Silty Sand Test</u>					# able to Rebury	# Not able to Rebury
Conc. Rep								
0	1	.	.	.	.	.	10	0
0	2	.	.	.	.	.	5	4
0	3	.	.	.	.	.	7	3
0	4	.	.	.	.	.	7	1
0	5	.	.	.	.	.	3	7
0.01	1	.	.	.	.	.	9	0
0.01	2	.	.	.	.	.	10	0
0.01	3	.	.	.	.	.	2	8
0.01	4	.	.	.	.	.	10	0
0.01	5	.	.	.	.	.	10	0
0.1	1	.	.	.	.	.	9	0
0.1	2	.	.	.	.	.	7	1
0.1	3	.	.	.	.	.	10	0
0.1	4	.	.	.	.	.	3	7
0.1	5	.	.	.	.	.	6	4
1	1	.	.	.	.	.	10	0
1	2	.	.	.	.	.	5	5
1	3	.	.	.	.	.	10	0
1	4	.	.	.	.	.	10	0
1	5	.	.	.	.	.	10	0

Table 5. Physical/chemical measurements taken every 24 hours for the fine sand test and the silty sand test in the 10 day amphipod test using paper pulp slurry. Water quality parameters include temperature (°C), pH, dissolved oxygen (mg/l), and salinity (ppt).

**Fine Sand Test**

Parameter	Conc.	Time (hours)										
		0	24	48	72	96	120	144	168	192	216	240
Temp.	0 mm	18.5	19.3	18.0	18.7	18.2	.	18.2	18.2	18.1	18.5	18.6
	0.01 mm	18.6	19.5	18.2	18.8	18.3	.	18.2	18.3	18.2	18.6	18.7
	0.1 mm	18.9	19.3	18.0	18.3	18.3	.	18.2	18.3	18.2	18.6	18.6
	1.0 mm	18.8	19.2	17.8	17.6	18.1	.	18.1	18.1	18.1	18.4	18.4
Salinity	0 mm	36	35	35	35	35	.	35	35	35	35	35
	0.01 mm	36	35	35	35	35	.	35	35	35	35	35
	0.1 mm	36	35	35	35	35	.	35	35	35	35	35
	1.0 mm	36	35	35	35	35	.	35	35	35	35	35
pH	0 mm	7.9	7.7	7.9	8.1	8.1	.	8.1	8.0	8.0	7.9	7.9
	0.01 mm	7.9	7.8	7.9	8.2	8.0	.	8.2	8.0	8.0	8.0	8.0
	0.1 mm	7.9	7.8	7.9	8.1	8.0	.	8.1	8.0	8.0	8.0	8.0
	1.0 mm	7.9	7.7	7.9	8.0	8.0	.	8.0	7.9	7.9	7.9	7.9
D.O.	0 mm	6.6	6.6	6.8	8.7	8.5	.	7.3	7.4	7.0	6.9	6.6
	0.01 mm	6.7	6.5	6.7	8.9	8.3	.	7.9	7.6	6.9	6.9	6.9
	0.1 mm	6.6	6.4	6.7	8.7	7.9	.	7.4	7.3	7.1	7.1	7.0
	1.0 mm	6.3	6.0	6.8	7.4	7.6	.	6.9	6.7	6.4	6.6	6.5

Note: Water quality parameters were not measured at the 120th test hour.

Table 5 continued.

**Silty Sand Test**

Parameter	Conc.	Time (hours)										
		0	24	48	72	96	120	144	168	192	216	240
Temp.	0 mm	18.7	19.0	18.0	18.2	17.7	.	17.7	17.9	17.7	18.0	18.1
	0.01 mm	18.7	19.0	17.8	18.1	17.6	.	17.6	18.1	17.7	17.9	18.0
	0.1 mm	18.5	18.8	18.0	18.1	17.6	.	17.5	17.8	17.7	17.9	18.0
	1.0 mm	18.8	19.2	17.8	18.4	18.0	.	18.1	17.9	18.1	18.2	18.4
Salinity	0 mm	36	35	35	35	35	.	35	35	35	35	35
	0.01 mm	36	35	35	35	35	.	35	35	35	35	35
	0.1 mm	36	35	35	35	35	.	35	35	35	35	35
	1.0 mm	36	35	35	35	35	.	35	35	35	35	35
pH	0 mm	7.9	7.8	7.9	8.1	8.0	.	8.1	7.9	8.0	8.0	8.0
	0.01 mm	7.9	7.8	7.9	8.1	8.0	.	8.1	8.0	8.0	8.0	8.0
	0.1 mm	8.0	7.9	7.9	8.1	7.9	.	8.1	7.9	7.9	7.9	7.9
	1.0 mm	7.9	7.9	7.9	8.0	7.9	.	8.0	7.9	7.9	7.9	7.9
D.O.	0 mm	6.6	6.1	6.6	8.2	8.5	.	7.9	7.2	6.9	6.9	6.9
	0.01 mm	6.6	6.1	6.8	8.2	8.4	.	8.0	7.1	6.8	6.9	7.0
	0.1 mm	6.7	6.2	6.7	8.1	8.3	.	7.6	7.1	6.6	6.7	6.8
	1.0 mm	6.4	6.0	6.7	7.8	8.1	.	7.5	6.9	6.3	6.4	6.4

Note: Water quality parameters were not measured on the 120th test hour.

## **5.0 Results of the 96 hour Polychaete Test**

Table 6. Summary of test information for the 96 hour polychaete test using paper pulp slurry during May 1995.

**Test Information:**

Date and Time of Test Initiation: 2 May 1995, 1500 hours

Concentrations Used:

Fine Sand Test: 0, 0.01, 0.1, and 1.0 mm layer of paper slurry

Silty Sand Test: 0, 0.01, 0.1, and 1.0 mm layer of paper slurry

Test Material Sources:

Dilution Water: La Jolla, CA.

Sediment: Newport Bay, CA.

Organisms: Long Beach, CA.

Paper Slurry: NRAD

Dates of Collection:

Dilution Water: 26 April 1995 and 3 May 1995

Sediment: 28 April 1995

Organisms: 2 May 1995

Paper Slurry: 27 April 1995

Table 7. Summary of final test results for the 96 hour polychaete test using paper pulp slurry during May 1995. For percent survival, the NOEC (no observed effect concentration) and ANOVA Mean Square Errors (MSE) are given for analyses of arcsin transformed data. All tabulated means are for untransformed data.

**Fine Sand Test**

CONCENTRATION (mm Paper Slurry)	% SURVIVAL	
	MEAN	S.D.
0	100.0	0.00
0.01	100.0	0.00
0.1	100.0	0.00
1	95.0	22.36

SURVIVAL:

-----  
 NOEC = 1  
 LOEC = .  
 EC50 = .  
 ANOVA MSE = 101.25

**Silty Sand Test**

CONCENTRATION (mm Paper Slurry)	% SURVIVAL	
	MEAN	S.D.
0	95.0	22.36
0.01	95.0	22.36
0.1	100.0	0.00
1	95.0	22.36

SURVIVAL:

-----  
 NOEC = 1  
 LOEC = .  
 EC50 = .  
 ANOVA MSE = 303.75

Table 8. Individual test data for survival in the 96 hour polychaete test using paper pulp slurry during May 1995.

		<u>Fine Sand Test</u>						%	%
Conc.	Rep						Alive	Dead	
0	1	.	.	.	.	.	100	0	
0	2	.	.	.	.	.	100	0	
0	3	.	.	.	.	.	100	0	
0	4	.	.	.	.	.	100	0	
0	5	.	.	.	.	.	100	0	
0	6	.	.	.	.	.	100	0	
0	7	.	.	.	.	.	100	0	
0	8	.	.	.	.	.	100	0	
0	9	.	.	.	.	.	100	0	
0	10	.	.	.	.	.	100	0	
0	11	.	.	.	.	.	100	0	
0	12	.	.	.	.	.	100	0	
0	13	.	.	.	.	.	100	0	
0	14	.	.	.	.	.	100	0	
0	15	.	.	.	.	.	100	0	
0	16	.	.	.	.	.	100	0	
0	17	.	.	.	.	.	100	0	
0	18	.	.	.	.	.	100	0	
0	19	.	.	.	.	.	100	0	
0	20	.	.	.	.	.	100	0	
0.01	1	.	.	.	.	.	100	0	
0.01	2	.	.	.	.	.	100	0	
0.01	3	.	.	.	.	.	100	0	
0.01	4	.	.	.	.	.	100	0	
0.01	5	.	.	.	.	.	100	0	
0.01	6	.	.	.	.	.	100	0	
0.01	7	.	.	.	.	.	100	0	
0.01	8	.	.	.	.	.	100	0	
0.01	9	.	.	.	.	.	100	0	
0.01	10	.	.	.	.	.	100	0	
0.01	11	.	.	.	.	.	100	0	
0.01	12	.	.	.	.	.	100	0	
0.01	13	.	.	.	.	.	100	0	
0.01	14	.	.	.	.	.	100	0	
0.01	15	.	.	.	.	.	100	0	
0.01	16	.	.	.	.	.	100	0	
0.01	17	.	.	.	.	.	100	0	
0.01	18	.	.	.	.	.	100	0	
0.01	19	.	.	.	.	.	100	0	
0.01	20	.	.	.	.	.	100	0	

Table 8 continued.

		<b><u>Fine Sand Test (continued)</u></b>	
Conc.	Rep	% Alive	% Dead
0.1	1	100	0
0.1	2	100	0
0.1	3	100	0
0.1	4	100	0
0.1	5	100	0
0.1	6	100	0
0.1	7	100	0
0.1	8	100	0
0.1	9	100	0
0.1	10	100	0
0.1	11	100	0
0.1	12	100	0
0.1	13	100	0
0.1	14	100	0
0.1	15	100	0
0.1	16	100	0
0.1	17	100	0
0.1	18	100	0
0.1	19	100	0
0.1	20	100	0
1	1	100	0
1	2	100	0
1	3	0	100
1	4	100	0
1	5	100	0
1	6	100	0
1	7	100	0
1	8	100	0
1	9	100	0
1	10	100	0
1	11	100	0
1	12	100	0
1	13	100	0
1	14	100	0
1	15	100	0
1	16	100	0
1	17	100	0
1	18	100	0
1	19	100	0
1	20	100	0

Table 8 continued.

		<u>Silty Sand Test</u>						%	%
Conc.	Rep						Alive	Dead	
0	1	.	.	.	.	.	100	0	
0	2	.	.	.	.	.	0	100	
0	3	.	.	.	.	.	100	0	
0	4	.	.	.	.	.	100	0	
0	5	.	.	.	.	.	100	0	
0	6	.	.	.	.	.	100	0	
0	7	.	.	.	.	.	100	0	
0	8	.	.	.	.	.	100	0	
0	9	.	.	.	.	.	100	0	
0	10	.	.	.	.	.	100	0	
0	11	.	.	.	.	.	100	0	
0	12	.	.	.	.	.	100	0	
0	13	.	.	.	.	.	100	0	
0	14	.	.	.	.	.	100	0	
0	15	.	.	.	.	.	100	0	
0	16	.	.	.	.	.	100	0	
0	17	.	.	.	.	.	100	0	
0	18	.	.	.	.	.	100	0	
0	19	.	.	.	.	.	100	0	
0	20	.	.	.	.	.	100	0	
0.01	1	.	.	.	.	.	100	0	
0.01	2	.	.	.	.	.	100	0	
0.01	3	.	.	.	.	.	100	0	
0.01	4	.	.	.	.	.	100	0	
0.01	5	.	.	.	.	.	100	0	
0.01	6	.	.	.	.	.	100	0	
0.01	7	.	.	.	.	.	100	0	
0.01	8	.	.	.	.	.	100	0	
0.01	9	.	.	.	.	.	100	0	
0.01	10	.	.	.	.	.	100	0	
0.01	11	.	.	.	.	.	100	0	
0.01	12	.	.	.	.	.	100	0	
0.01	13	.	.	.	.	.	100	0	
0.01	14	.	.	.	.	.	100	0	
0.01	15	.	.	.	.	.	100	0	
0.01	16	.	.	.	.	.	100	0	
0.01	17	.	.	.	.	.	100	0	
0.01	18	.	.	.	.	.	100	0	
0.01	19	.	.	.	.	.	100	0	
0.01	20	.	.	.	.	.	0	100	

Table 8 continued.

		<u>Silty Sand Test (continued)</u>						%	%
Conc. Rep								Alive	Dead
0.1	1	.	.	.	.	.	100	0	
0.1	2	.	.	.	.	.	100	0	
0.1	3	.	.	.	.	.	100	0	
0.1	4	.	.	.	.	.	100	0	
0.1	5	.	.	.	.	.	100	0	
0.1	6	.	.	.	.	.	100	0	
0.1	7	.	.	.	.	.	100	0	
0.1	8	.	.	.	.	.	100	0	
0.1	9	.	.	.	.	.	100	0	
0.1	10	.	.	.	.	.	100	0	
0.1	11	.	.	.	.	.	100	0	
0.1	12	.	.	.	.	.	100	0	
0.1	13	.	.	.	.	.	100	0	
0.1	14	.	.	.	.	.	100	0	
0.1	15	.	.	.	.	.	100	0	
0.1	16	.	.	.	.	.	100	0	
0.1	17	.	.	.	.	.	100	0	
0.1	18	.	.	.	.	.	100	0	
0.1	19	.	.	.	.	.	100	0	
0.1	20	.	.	.	.	.	100	0	
1	1	.	.	.	.	.	100	0	
1	2	.	.	.	.	.	100	0	
1	3	.	.	.	.	.	100	0	
1	4	.	.	.	.	.	100	0	
1	5	.	.	.	.	.	100	0	
1	6	.	.	.	.	.	0	100	
1	7	.	.	.	.	.	100	0	
1	8	.	.	.	.	.	100	0	
1	9	.	.	.	.	.	100	0	
1	10	.	.	.	.	.	100	0	
1	11	.	.	.	.	.	100	0	
1	12	.	.	.	.	.	100	0	
1	13	.	.	.	.	.	100	0	
1	14	.	.	.	.	.	100	0	
1	15	.	.	.	.	.	100	0	
1	16	.	.	.	.	.	100	0	
1	17	.	.	.	.	.	100	0	
1	18	.	.	.	.	.	100	0	
1	19	.	.	.	.	.	100	0	
1	20	.	.	.	.	.	100	0	

Table 9. Physical/chemical measurements taken every 24 hours for the fine sand test and the silty sand test in the 96 hour polychaete test using paper pulp slurry. Water quality parameters include temperature ( $^{\circ}\text{C}$ ), pH, dissolved oxygen (mg/l), and salinity (ppt).

**Fine Sand Test**

Parameter	Concentration	Time (hours)				
		0	24	48	72	96
Temp.	0 mm	15.4	15.4	15.6	15.0	15.5
	0.01 mm	15.3	15.1	15.2	14.7	15.2
	0.1 mm	16.0	15.5	15.7	15.0	15.6
	1.0 mm	15.3	15.9	16.8	16.2	16.9
Salinity	0 mm	36	36	36	36	36
	0.01 mm	36	36	36	36	36
	0.1 mm	36	36	36	36	36
	1.0 mm	36	36	36	36	36
pH	0 mm	7.8	7.9	8.0	8.1	8.1
	0.01 mm	7.8	7.9	8.0	8.1	8.1
	0.1 mm	7.8	7.9	8.0	8.0	8.0
	1.0 mm	7.8	8.0	8.1	8.1	8.1
D.O.	0 mm	7.8	8.2	8.2	8.3	8.3
	0.01 mm	7.4	8.0	7.9	8.2	8.3
	0.1 mm	7.6	8.0	8.1	8.1	8.2
	1.0 mm	7.6	8.3	8.1	8.3	8.2

Table 9 continued.

**Silty Sand Test**

Parameter	Concentration	Time (hours)				
		0	24	48	72	96
Temp.	0 mm	15.6	15.3	15.5	15.0	15.4
	0.01 mm	15.7	15.7	15.9	15.3	15.7
	0.1 mm	16.0	15.8	16.1	15.6	16.1
	1.0 mm	15.6	15.3	15.6	15.1	15.7
Salinity	0 mm	36	36	36	36	36
	0.01 mm	36	36	36	36	36
	0.1 mm	36	36	36	36	36
	1.0 mm	36	36	36	36	36
pH	0 mm	7.6	7.9	8.0	8.1	8.1
	0.01 mm	7.6	7.9	8.0	8.1	8.2
	0.1 mm	7.6	7.9	8.0	8.1	8.1
	1.0 mm	7.6	7.9	8.0	8.1	8.1
D.O.	0 mm	7.8	8.0	8.2	8.4	8.3
	0.01 mm	7.8	8.1	8.4	8.6	8.2
	0.1 mm	7.7	7.9	8.2	8.3	8.2
	1.0 mm	7.8	7.7	8.1	8.3	8.2

## **6.0 Results of the Sediment Analysis**

## Sediment Analysis

Table 10. Results of grain size analysis and Total Organic Carbon (TOC) content in sediments used in toxicity tests with amphipods and polychaetes.

### Amphipod Test

	<u>Fine Sand</u>	<u>Silty Sand</u>
Mean Grain Size:	166 microns	184 microns
% Sand:	95.5 %	91.7 %
% Silt:	2.1 %	3.4 %
% Clay:	2.4 %	4.9 %
TOC content:	1040 mg/kg	1190 mg/kg

### Polychaete Test

	<u>Fine Sand</u>	<u>Silty Sand</u>
Mean Grain Size:	187 microns	178 microns
% Sand:	97.3 %	96.1 %
% Silt:	1.2 %	1.5 %
% Clay:	1.5 %	2.4 %
TOC content:	552 mg/kg	861 mg/kg

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 1 of 9**

**SOP No.: TS 018.00**

**Effective Date: 03/23/95**

---

### **1.0 Scope**

This SOP describes general test methods for the 96 hour acute test for sediment toxicity with polychaetes.

### **2.0 Application**

This test is used as an acute marine sediment toxicity test for polychaetes.

### **3.0 Health & Safety**

Test substances used in the polychaete sediment toxicity test may be toxic and special care will be taken in handling these toxic substances. Health and safety procedures relevant to toxicity testing are described in SOP's H 001 through H 015.

### **4.0 Definitions**


NOEC - No observed effect concentration. The highest concentration of a test or reference substance that does not cause a statistically significant reduction in survival.

LOEC - Lowest observed effect concentration. The lowest concentration of a test or reference substance that causes a statistically significant reduction in survival.

LC<sub>50</sub> - A statistically or graphically derived estimate of the concentration of a test or reference substance that is lethal to 50% of the test systems exposed.

Control Substance - Any chemical substance or mixture, or any substance other than the test substance, feed, or water, that is administered to the test system in the course of the

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 2 of 9**

**SOP No.: TS 018.00**

**Effective Date: 03/23/95**

---

study for the purpose of establishing a basis for comparison with the test substance for known chemical or biological measurements.

**Dilution Water** - The water used to dilute test substances for use in toxicity tests.

**Reference Substance**- Any chemical substance or mixture or analytical standard other than the test substance, feed, or water, that is administered to the test system in the course of the study for the purpose of establishing a basis for comparison with the test substance for known chemical or biological measurements.

**Test Substance** - A substance or mixture administered or added to a test system in a study.


**Test System** - Any organism (animal or plant) to which a test, control, or reference substance is administered or added for study.


**Start time for the test** - The time of addition of the first test system to the test substance, reference substance, or control.

### **5.0 Equipment**

- 1 L glass beakers or jars (or equivalent disposable container)
- Sieve (0.5 mm mesh)
- Plastic sheeting
- Thermometer
- D.O. meter
- pH meter
- Refractometer
- Cool white fluorescent light
- Temperature controlled room

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.**  
**Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 3 of 9**

**SOP No.: TS 018.00**

**Effective Date: 03/23/95**

---

Continuous temperature monitor  
Air pump, air lines, and disposable pipet tip

## 6.0 Procedures

### 6.1 Test System

Species	<i>Neanthes arenaceodentata</i>
Source	Don Reish, California State University Long Beach, Long Beach, CA
Age/Life-stage from the parent's tube	2 to 3 months from time of emergence
Acclimation	Acclimate to 18°C by adjusting temperature at a rate of no more than 3°C per 24 hours (SOP SY )
Records	Maintain test system log sheet (SOP SY 001)
Feeding	None during conduct of test, 8 mg Tetramin per worm every other day during holding

### 6.2 Test Substance

Test substances will be supplied by the client or sampled by an employee of Coastal Resources Associates, Inc.

### 6.3 General Test Conditions

Temperature	17 to 20°C ± 3°C
Salinity	34 ppt
Photoperiod	None specified

---

Signed: Susan T. Rojano  
Susan T. Rojano, Laboratory Manager

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 4 of 9**

**SOP No.: TS 018.00**


**Effective Date: 03/23/95**

---


Test chamber	1 L beaker, jar, or equivalent disposable container
Dilution water source	Uncontaminated seawater
Number dilutions per sample	5 (unless otherwise specified in protocol)
Number of controls	1 (minimum) consists of a set of replicates using sediment from the location at which the organisms were collected and uncontaminated seawater, additional controls may be needed for testing other sediments
Number of replicates per test dilution	20 recommended (10 minimum)
Number of replicates per control	20 recommended (10 minimum)
Volume of dilutions	700 ml/replicate (> 175 ml for sediment)
Number of test systems per chamber	1
Renewal of test substances	At 48 hours
Renewal of reference substances	At 48 hours
Type of biological observations	Number of animals alive
Definition of death	Opaque white coloration, immobility, and lack of reaction to gentle prodding
Times of biological observations	Daily - number dead At termination - number alive
Type of physical/chemical measurements	Temperature, D.O., pH, salinity

---

Signed:

  
Susan T. Rojano, Laboratory Manager

Approved:

  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 5 of 9**

**SOP No.: TS 018.00**

**Effective Date: 03/23/95**

---

Times of physical/chemical measurements

Continuous - room temperature  
Daily - Chamber temperature, D.O., pH, salinity

Dilutions for physical/chemical measurements

One randomly selected replicate per treatment - Chamber temperature, D.O., pH, salinity  
Bath only - continuous temperature

Dilutions for test substance

Determined by range finding test or by purpose of the study

#### **6.4 Definitive Test with a Test Substance**

Steps for conducting the definitive test are as follows:

- Receive polychaetes (SOP SY 001) and acclimate (SOP SY 011).
- The day before test initiation, add homogenized sediment and seawater to the test chambers and allow the sediment to settle. Enough sediment must be added to create a 2 cm deep layer on the bottom of each test chamber and overlying water should be added to the 700 ml mark on the test chambers.
- Cover all test chambers with plastic sheeting to minimize evaporation and reduce the risk of contamination.
- On test initiation day, prepare dilutions of the test substance and equilibrate dilutions to appropriate test conditions (SOP SU 014)
- Siphon as much overlying seawater off as possible without disturbing the sediment.

---

Signed: Susan T. Rojano  
Susan T. Rojano, Laboratory Manager

---

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 6 of 9**


**SOP No.: TS 018.00**

**Effective Date: 03/23/95**

---

- Add the newly prepared dilutions of the test substance to the appropriate test chambers up to the 700 ml mark. Slowly pour the dilutions down the side of the test chamber or glass rod to reduce disturbance to the sediment.
- Measure physical/chemical parameters as described in section 6.3 above and record the results on the water quality data sheet (TS-18-1).
- Add 1 polychaete to each test chamber using a wide bore pipette with a fire polished tip (SOP SY 002).
- Confirm there is a polychaete in each test chamber. Record the results on the biological observations data sheet (TS-18-2).
- Approximately twenty-four hours after the start of the test, count the number of dead polychaetes in each test chamber. Remove any dead organisms from the test containers using a wide bore pipette with a fire polished tip. Record the results on the biological observations data sheet (TS-18-2).
- Measure physical/chemical parameters as indicated in section 6.3 above and record the results on the water quality data sheet (TS-18-1).
- Approximately forty-eight hours after the start of the test, count the number of dead polychaetes in each test chamber and record the results on the biological observations data sheet (TS-18-2).

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 7 of 9**


**SOP No.: TS 018.00**


**Effective Date: 03/23/95**

---

- Measure physical/chemical parameters as indicated in section 6.3 above and record the results on the water quality data sheet (TS-18-1).
- Siphon off 75% of the test water and any dead organisms. Siphon using an airline with a wide bore pipette with a fire polished tip attached. Use a separate clean tip for each dilution.
- Add newly prepared dilutions of the test substance to the appropriate test chambers up to the 700 ml mark. Slowly pour the dilutions down the side of the test chamber or a glass rod to reduce disturbance to the sediment.
- After renewal, measure physical/chemical parameters as indicated in section 6.3 above and record the results on the water quality data sheet (TS-18-1).
- Count the number of dead polychaetes daily and record on the biological observations data sheet (TS-18-2).
- Measure physical/chemical parameters as described in section 6.3 daily.
- Terminate the test after 96 hours of exposure.
- Sieve the contents of each test vessel individually through a 0.5 mm screen to remove the test organisms. Use dilution water with a salinity and temperature within two units for sieving.

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 8 of 9**

**SOP No.: TS 018.00**

**Effective Date: 03/23/95**

---

- Rinse material retained on the screen into a tray for closer examination.
- Count the number of live polychaetes and record the results on the biological observations data sheet (TS-18-2).
- Dispose of the polychaetes (SOP SY 003) and the test substance (SOP SU 006).

**6.5 Recording and analyzing data**

- Enter all data onto data sheets according to procedures given in SOP D 001. Enter these data into computer files (SOP D 002).
- Analyze the test data as described in SOP D 003 using SAS statistical software. Determine the  $LC_{50}$ , the NOEC, and the LOEC.

**6.6 Test Acceptability**

- Total survival in the controls must be 90% or greater.

**6.7 Documentation and Reports**

- Documents listed in SOP D 005 will be completed. Data sheets specific to this test procedure are attached. These data and any subsequent analysis of the data will be archived as indicated in SOP D 006.
- Reports will be prepared as per SOP D 007 and documents archived per SOP D 008.

---

Signed: Susan T. Rojano  
Susan T. Rojano, Laboratory Manager

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 96 Hour Acute Sediment Toxicity Test with Polychaete Worms**

**Pg. 9 of 9**

**SOP No.: TS 018.00**

**Effective Date: 03/23/95**

---

**7.0 Personnel**

All Coastal Resources Associates, Inc. technical staff trained in specific tasks related to this test will use this SOP.

---

Signed: *Susan T. Rojano*  
Susan T. Rojano, Laboratory Manager

Approved: *Thomas A. Dean*  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.**  
**Standard Operating Procedure**

**Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity**

**Pg. 1 of 9**

**SOP No.: TS 009.00**

**Effective Date: 03/23/95**

---

**1.0 Scope**

This SOP describes general test methods for the 10 day acute test for sediment toxicity with amphipods. The procedures are modified from ASTM E1367 (1993).

**2.0 Application**

This test is used as a marine sediment toxicity test for amphipods.

**3.0 Health & Safety**

Test substances used in the amphipod sediment toxicity test may be toxic and special care will be taken in handling these toxic substances. Health and safety procedures relevant to toxicity testing are described in SOP's H 001 through H 015.

**4.0 Definitions**

NOEC - No observed effect concentration. The highest concentration of a test or reference substance that does not cause a statistically significant reduction in survival.

LOEC - Lowest observed effect concentration. The lowest concentration of a test or reference substance that causes a statistically significant reduction in survival.

LC<sub>50</sub> - A statistically or graphically derived estimate of the concentration of a test or reference substance that is lethal to 50% of the test systems exposed.

Control Substance - Any chemical substance or mixture, or any substance other than the test substance, feed, or water, that is administered to the test system in the course of the

---

Signed: Susan T. Rojano  
Susan T. Rojano, Laboratory Manager

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity**

**Pg. 2 of 9**

**SOP No.: TS 009.00**

**Effective Date: 03/23/95**

---

study for the purpose of establishing a basis for comparison with the test substance for known chemical or biological measurements.

**Dilution Water** - The water used to dilute test substances for use in toxicity tests.

**Reference Substance**- Any chemical substance or mixture or analytical standard other than the test substance, feed, or water, that is administered to the test system in the course of the study for the purpose of establishing a basis for comparison with the test substance for known chemical or biological measurements.

**Test Substance** - A substance or mixture administered or added to a test system in a study.

**Test System** - Any organism (animal or plant) to which a test, control, or reference substance is administered or added for study.

**Start time for the test** - The time of addition of the first test system to the test substance, reference substance, or control.

## **5.0 Equipment**

- 1 L glass beakers or jars (or equivalent disposable container)
- Sieve (0.5 mm mesh)
- Plastic sheeting
- Thermometer
- D.O. meter
- pH meter
- Refractometer
- Cool white fluorescent light
- Temperature controlled room

---

Signed: Susan T. Rojano  
Susan T. Rojano, Laboratory Manager

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.**  
**Standard Operating Procedure**

**Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity**

**Pg. 3 of 9**

**SOP No.: TS 009.00**

**Effective Date: 03/23/95**

---

Continuous temperature monitor  
Air pump, air lines, and disposable pipet tip

**6.0 Procedures**


**6.1 Test System**


Species	<i>Grandidierella japonica</i> or <i>Rhepoxynius abronius</i>
Source	David Gutoff, San Diego, CA or Ken Brooks, Port Townsend, WA
Age/Life-stage	large immature and adult amphipods, 3 to 5 mm in length
Acclimation	Acclimate to 15°C for <i>Rhepoxynius</i> and 17°C for <i>Grandidierella</i> by adjusting temperature at a rate of no more than 3°C per 24 hours (SOP SY )
Identification	Source for ID is Environment Canada Report EPS 1/RM/26 (December 1992)
Records	Maintain test system log sheet (SOP SY 001)
Feeding	None

**6.2 Test and Reference Substance**

Test substances will be supplied by the client or sampled by an employee of Coastal Resources Associates, Inc.

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

Coastal Resources Associates, Inc.  
Standard Operating Procedure

Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity

Pg. 4 of 9

SOP No.: TS 009.00

Effective Date: 03/23/95


---

### 6.3 General Test Conditions

Temperature	15°C ± 3°C for <i>Rhepoxynius</i> 15 to 19°C ± 3°C for <i>Grandidierella</i>
Salinity	28 ppt for <i>Rhepoxynius</i> 30 to 35 ppt for <i>Grandidierella</i>
Photoperiod	Continuous throughout the test period
Test chamber	1 L beaker or jar, or equivalent disposable container
Dilution water source	Uncontaminated seawater
Number dilutions per sample	5 (unless otherwise specified in protocol)
Number of controls	1 (minimum) consists of a set of replicates using sediment from the location at which the organisms were collected and uncontaminated seawater, additional controls may be needed for testing other sediments
Number of replicates per test dilution	5
Number of replicates per control	5
Volume of dilutions	700 ml/replicate (> 175 ml for sediment)
Number of test systems per chamber	20
Renewal of test substances	At 48 hour intervals
Renewal of reference substances	At 48 hour intervals

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity**

**Pg. 5 of 9**

**SOP No.: TS 009.00**

**Effective Date: 03/23/95**

---

Type of biological observations	Emergence from sediment, number of animals alive, and ability to rebury
Definition of death	No movement when a pulse of water is applied through a disposable pipet to the test system
Times of biological observations	Daily - emergence At termination - number alive and ability to rebury
Type of physical/chemical measurements	Temperature, D.O., pH, salinity
Times of physical/chemical measurements	Continuous - room temperature Daily - Chamber temperature, D.O., pH, salinity
Dilutions for physical/chemical measurements	One randomly selected replicate per treatment - Chamber temperature, D.O., pH, salinity Bath only - continuous temperature
Dilutions for test substance	Determined by range finding test or by purpose of the study

#### **6.4 Definitive Test with a Test Substance**

Steps for conducting the definitive test are as follows:

- Receive amphipods (SOP SY 001 ) and acclimate (SOP SY 011).
  
- The day before test initiation, add homogenized sediment and seawater to the test chambers and allow the sediment to settle. Enough sediment must be added to create a 2 cm deep layer on the bottom of each test chamber and overlying water should be added to the 700 ml mark on the test chambers.

---

Signed: Susan F. Rojano  
Susan F. Rojano, Laboratory Manager

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity**

**Pg. 6 of 9**

**SOP No.: TS 009.00**


**Effective Date: 03/23/95**

---

- Cover each test chamber with plastic sheeting to minimize evaporation and reduce the risk of contamination.
- On test initiation day, prepare dilutions of the test substance and equilibrate dilutions to appropriate test conditions (SOP SU 014).
- Siphon as much overlying seawater off as possible without disturbing the sediment.
- Add the newly prepared dilutions of the test substance to the appropriate test chambers up to the 700 ml mark. Slowly pour the dilutions down the side of the test chamber or glass rod to reduce disturbance to the sediment.
- Measure physical/chemical parameters as described in section 6.3 above and record the results on the water quality data sheet (TS-9-1).
- Add 20 amphipods to each test chamber using a wide bore pipette with a fire polished tip (SOP SY 002).
- Carefully count as amphipods are added to confirm there are 20 in each test chamber. Record the results on the biological observations data sheet (TS-9-2).
- Approximately twenty-four hours after the start of the test, count the number of amphipods afloat in each test chamber. Record the results on the biological observations data sheet (TS-9-2).

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

Coastal Resources Associates, Inc.  
Standard Operating Procedure

Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity

Pg. 7 of 9

SOP No.: TS 009.00

Effective Date: 03/23/95

---

- Measure physical/chemical parameters as indicated in section 6.3 above and record the results on the water quality data sheet (TS-9-1).
- Approximately forty-eight hours after the start of the test, count the number of amphipods afloat in each test chamber and record the results on the biological observations data sheet (TS-9-2).
- Measure physical/chemical parameters as indicated in section 6.3 above and record the results on the water quality data sheet (TS-9-1).
- Siphon off 75% of the test water and any floating organisms. Siphon using an airline with a wide bore pipette with a fire polished tip attached. Use a separate clean tip for each dilution.
- Add newly prepared dilutions of the test substance to the appropriate test chambers up to the 700 ml mark. Slowly pour the dilutions down the side of the test chamber or a glass rod to reduce disturbance to the sediment.
- After renewal, measure physical/chemical parameters as indicated in section 6.3 above and record the results on the water quality data sheet (TS-9-1).
- Count the number of amphipods afloat daily and record on the biological observations data sheet (TS-9-2).

---

Signed: Susan T. Rojano  
Susan T. Rojano, Laboratory Manager

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity**

**Pg. 8 of 9**


**SOP No.: TS 009.00**


**Effective Date: 03/23/95**

---

- Measure physical/chemical parameters as described in section 6.3 daily. On renewal days measure the parameters before and after renewal.
- Renew the overlying water as described above at 48 hour intervals.
- Terminate the test after 10 days of exposure.
- Sieve the contents of each test vessel individually through a 0.5 mm screen to remove the test organisms. Use dilution water with a salinity and temperature within two units for sieving.
- Rinse material retained on the screen into a tray for closer examination.
- Count the number of live amphipods and record the results on the biological observations data sheet (TS-9-2).
- Place surviving amphipods from each dilution in a separate container with a 2 cm layer of control sediment.
- After 1 hour, count the number of surviving amphipods unable to rebury. Record the results on the biological observations data sheet (TS-9-2).
- Dispose of the amphipods (SOP SY 003) and the test substance (SOP SU 006).

---

Signed:   
Susan T. Rojano, Laboratory Manager

Approved:   
Thomas A. Dean, Ph.D., Laboratory Director

---

---

**Coastal Resources Associates, Inc.  
Standard Operating Procedure**

**Subject: 10 Day Amphipod Bioassay for Marine Sediment Toxicity**

**Pg. 9 of 9**

**SOP No.: TS 009.00**

**Effective Date: 03/23/95**

---

**6.5 Recording and analyzing data**

- Enter all data onto data sheets according to procedures given in SOP D 001. Enter these data into computer files (SOP D 002).
  
- Analyze the test data as described in SOP D 003 using SAS statistical software. Determine the  $LC_{50}$ , the NOEC, and the LOEC.

**6.6 Test Acceptability**

- Total survival in the controls must be 90% or greater.
  
- Each individual control replicate must have at least 80% survival.

**6.7 Documentation and Reports**

- Documents listed in SOP D 005 will be completed. Data sheets specific to this test procedure are attached. These data and any subsequent analysis of the data will be archived as indicated in SOP D 006.
  
- Reports will be prepared as per SOP D 007 and documents archived per SOP D 008.

**7.0 Personnel**

All Coastal Resources Associates, Inc. technical staff trained in specific tasks related to this test will use this SOP.

---

Signed: Susan T. Rojano  
Susan T. Rojano, Laboratory Manager

Approved: Thomas A. Dean  
Thomas A. Dean, Ph.D., Laboratory Director

---

## **APPENDIX E**

### **ZOOPLANKTON INTERACTION REPORT**

Source:       Effects of Paper Pulp Wastes on the Feeding of Copepods.  
              Marine Life Research Group 0218  
              San Diego, California  
              Scripps Institution of Oceanography, 1994-1995

Effects of paper pulp wastes on the feeding of copepods

Hae Jin Jeong

Marine Life Research Group 0218, Scripps Institution of  
Oceanography, University of California San Diego, La Jolla,  
California 92093-0218

## Introduction

The amount and types of anthropogenic products introduced into marine environments have continuously increased. Usually, when these products are introduced into estuaries and semi-enclosed embayments where water circulation is restricted, food webs in these ecosystems can be significantly affected by these products. However, this may not occur in open oceans because of their large water volume and active circulation.

It is planned to dump naval pulp wastes (pulverized paper products) into offshore waters, and it must be determined whether these wastes may affect the ecology of some major components of marine organisms. Copepods are one of the dominant macrozooplankton in most marine environments and play important roles in food webs as major consumers of phytoplankton and microzooplankton, an important food source for diverse carnivores, and as nutrient regenerators. Therefore, changes in their abundances or feeding rates can significantly affect the abundances of their prey and/or predators.

Pulp wastes themselves, and/or leached chemicals, may significantly reduce ingestion rates of copepods on suitable prey by clogging the predators' feeding apparatus or by poisoning them (H<sub>01</sub> and H<sub>03</sub> below). If copepods can survive in dense pulp wastes and then recover their feeding rates on suitable prey after pulp waste has sunk or been dispersed, the

wastes will not significantly affect the ecology of copepods (H02).

To investigate these topics, the following hypotheses will be tested:

H01: The ingestion rate of phytoplankton by copepods is independent of the presence of slurry of Pulp wastes.

H02: There is no effect on ingestion rates in slurry-free water of previous exposure to slurry.

H03: there is no difference in ingestion rates in slurry-free sea water in which slurry had been soaked for 24 hour and then removed by filtration, relative to sea water never contacting slurry.

## MATERIALS AND METHODS

### Preparation of experimental organisms and conditions

The dinoflagellates Gymnodinium sanguineum and Gonyaulax polyedra and common copepods Acartia spp. and Calanus pacificus were chosen for these experiments. G. sanguineum and G. polyedra are common red-tide dinoflagellates and known as prey for Acartia spp. and C. pacificus. They were grown in enriched f/4 seawater media (Guillard & Ryther 1962) without silicate, at room temperature (20-23°C) with continuous illumination of 100

$\mu\text{E m}^{-2}\text{s}^{-1}$  of cool white fluorescent lights. Cultures in exponential growth phase were used for feeding experiments.

Adult female C. pacificus were collected from the coastal waters off La Jolla Bay, CA using a 303  $\mu\text{m}$  mesh net, and adult female Acartia spp. from the waters of Mission Bay, CA using a 54  $\mu\text{m}$  mesh net. Copepods were maintained at 15 °C room in 1 gallon jars with G. sanguineum or G. polyedra in filtered sea water for at least two days before experiments.

### **Experimental designs**

The initial densities of the predator and prey, and slurry are given in Table 1. Experiments 1 and 2 was designed to test H<sub>01</sub> (ingestion rate of C. pacificus or Acartia spp. is independent of the presence of slurry) stated previously. Experiment 3 was designed to test H<sub>01</sub> and H<sub>02</sub> (no difference in ingestion rates between copepods previously incubated with and without slurry). Experiment 4 was designed to test H<sub>03</sub> (there is no difference in ingestion rates in slurry-free sea water in which slurry had been soaked for 24 hour and then removed by filtration, relative to sea water never contacting slurry).

To set up an experiment, three 1 ml aliquots from a G. sanguineum or G. polyedra culture were counted to determine density. The concentrations of G. sanguineum or G. polyedra were obtained by volume dilution with an autopipette. The wet weight of slurry was measured on a microbalance, and each concentration (ratio of wet weight of slurry to weight of sea water) of slurry

was obtained by adding a known weight of slurry into Polycarbonate (PC) bottles. Slurry inside bottles was not homogeneously distributed, even though bottles were rotated. Such an aggregation of slurry may be also true in nature.

Copepods maintained in a 15 °C room were rinsed with filtered sea water in a Petri-dish, and 5 healthy female Calanus (in experiments 1, 3, and 4) or 8 female Acartia spp. (in experiment 2) were transferred into each 500 or 270 ml PC bottle, respectively. Duplicate experiment bottles were set up, as were duplicate control bottles containing only G. sanguineum or G. polyedra and slurry at all slurry concentrations. Actual initial concentrations of G. sanguineum or G. polyedra were measured in one extra control bottle by counting and removing more than 200 individual cells with a Pasteur micropipette. Experimental and control bottles were placed on rotating wheels at 0.9 RPM under dim light at 15°C for 16 - 20 h. After incubation, 2 ml aliquots from each bottle were transferred into multiwell chambers for counting G. sanguineum or G. polyedra cells (after serial dilution where necessary), and C. pacificus or Acartia spp. were sieved onto a 101 µm net and counted. Ingestion rates (prey ingested copepod<sup>-1</sup> hour<sup>-1</sup>) of copepod on G. sanguineum or G. polyedra were calculated, using the equations of Frost (1972), from final concentrations of prey in bottles with and without Calanus or Acartia.

The slurry concentration of 0.6 % was used in experiment 3 because this concentration caused a large reduction in feeding

in experiment 1. Two different predator-prey combinations were initially set up in duplicate: (1) 5 female C. pacificus (10 C. pacificus l<sup>-1</sup>) and G. sanguineum (2) 5 female C. pacificus, G. sanguineum, and slurry. Duplicate control bottles were similarly set up without copepods. Bottles were incubated for 24 h as described above (in Table 1, t=0). After counting cells, all C. pacificus were sieved onto a 101  $\mu$ m net, counted, and transferred into new bottles containing only new G. sanguineum cells without slurry (in Table 1, t=24h). New duplicate control bottles containing only G. sanguineum were set up. Bottles were incubated again for 24 h as described above, and cells and Calanus were counted.

In experiment 4, 0.6% slurry in filtered sea water was placed in a 15°C room. Twenty-four hours later, the slurry was screened out onto a GF/C millipore filter, and the filtrate sea water was transferred into four PC bottles. G. sanguineum was added to all four, and 5 female C. pacificus to two of these. Controls were similarly set up using sea water which had not been exposed to slurry. Bottles were incubated for 24 h as described above, and cells and Calanus were counted.

### **Test of hypotheses**

In experiments 1 and 2, the initial concentration of G. sanguineum or G. polyedra was fixed, while that of slurry varied (Table 1). An Analysis of Variance (ANOVA, Zar 1984) was used to test whether ingestion rates of G. sanguineum or G.

polyedra by C. pacificus or Acartia spp., respectively, at one slurry concentration were significantly different from those at other slurry concentrations ( $H_01$ ).

$H_02$  can be rejected if ingestion rates of C. pacificus previously incubated with slurry are significantly different (by two-tailed, two-sample t test) from those never exposed to slurry.

$H_03$  can be rejected if ingestion rates in sea water in which slurry had been soaked for 24 hour and then removed by filtration are significantly different (by two-tailed, two-sample t test) from those in sea water never contacting slurry.

## RESULTS

**Test of  $H_01$**  (*ingestion rate of phytoplankton by copepods is independent of the presence of slurry*)

With increasing slurry concentration, the ingestion rates of Gymnodinium sanguineum by Calanus pacificus exponentially decreased from 205 to 12 prey Calanus<sup>-1</sup> h<sup>-1</sup> (Fig. 1).

Ingestion rates of G. sanguineum by C. pacificus were significantly reduced by slurry (ANOVA,  $p < 0.005$ ; Zar 1984). Therefore,  $H_01$  can be rejected when G. sanguineum and C. pacificus were prey and predator. Ingestion rates at slurry concentrations of 0.05 and 0.1% were not significantly different

from that without added slurry ( $p > 0.05$ ), but they were significantly depressed at slurry concentrations  $\geq 0.3\%$  ( $p < 0.05$ ).

With increasing slurry concentration, the ingestion rates of Gonyaulax polyedra by Acartia spp. also decreased from 22 to 5 prey Acartia<sup>-1</sup> h<sup>-1</sup> (Fig. 2).

Ingestion rates of G. polyedra by Acartia spp. were significantly reduced by slurry (ANOVA,  $p < 0.05$ ). Therefore,  $H_01$  can also be rejected when G. polyedra and Acartia spp. were prey and predator. The ingestion rate at a slurry concentration of 0.1% was not significantly different from that without added slurry ( $p > 0.05$ ), but was significantly depressed at 0.6% ( $p < 0.05$ ).

**Test of  $H_02$**  (*no effect on ingestion rates in slurry-free water of previous exposure to slurry*)

In experiment 3, after first day incubation, the ingestion rate of Calanus on Gymnodinium sanguineum incubated with the slurry concentration of 0.6% was significantly different from that without slurry (Fig. 3, two tailed-t test,  $p < 0.05$ ), similar to the result in experiment 1. However, the ingestion rate of the Calanus, originally incubated with 0.6% slurry for 24 hour and then transferred into new bottles containing G. sanguineum without slurry, was not significantly different from that of the Calanus, continuously incubated without slurry.

Therefore, H<sub>02</sub> cannot be rejected. The results show that Calanus recovers its feeding rate when slurry disappears.

**Test of H<sub>03</sub>** (*there is no difference in ingestion rates in slurry-free sea water in which slurry had been soaked for 24 hour and then removed by filtration, relative to sea water never contacting slurry*)

The ingestion rate of Calanus in slurry-free sea water in which slurry had been soaked for 24 hour and then removed by filtration was not significantly different from that in sea water never contacting slurry ( $p > 0.05$ , grey bars in Fig. 3).

### Discussion

The presence of slurry significantly significantly reduced the ingestion rates of Calanus pacificus on G. sanguineum at the slurry concentrations  $\geq 0.3\%$ . However, Calanus, originally exposed to 0.6 % slurry for 24 hour, can recover its feeding rates when slurry disappears (Fig. 3). Therefore, if slurry is diluted quickly due to water movement after being dumped at 0.6% concentration, its presence may not affect the abundance of Calanus. The presence of slurry also significantly reduced the ingestion rates of Acartia spp. on Gonyaulax polyedra, however, the magnitude of the reduction of ingestion rates by Acartia spp. (4 times) was smaller than that by Calanus (17 times). The habitat of Acartia spp. (i.e. estuaries or coastal waters) is in

habitat of Acartia spp. (i.e. estuaries or coastal waters) is in general more turbid and polluted than that of Calanus (i.e. offshore). The adaptation of Acartia spp. to turbid environments may be partially responsible for its greater tolerance of slurry.

Chemicals leached from slurry did not affect the feeding rate of Calanus (Fig. 3). Mechanical interferences of slurry on the feeding and/or swimming behavior of copepods may be mainly responsible for the reduction of the ingestion rates.

#### LITERATURE CITED

- Frost, B.W. (1972). Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod Calanus pacificus. *Limnol. Oceanogr.* 17:805-815
- Heinbokel, J.F. (1978). Studies on the functional role of tintinnids in the Southern California Bight. I. Grazing and growth rates in laboratory cultures. *Mar. Biol.* 47:177-189
- Zar, J.H. (1984). *Biostatistical analysis*. Prentice Hall, Englewood Cliffs

Table 1. Design of experiments.

Experiment No.	time <sup>1</sup>	Slurry (%) <sup>2</sup>	Prey <sup>3</sup> (cells. ml <sup>-1</sup> )	Predator <sup>4</sup> (inds. l <sup>-1</sup> )
1	t=0	0, 0.05, 0.1, 0.3, 0.6	123	10*
2	t=0	0, 0.1, 0.6	190	30
3	t=0	0, 0.6	183	10
	t=24h	0, 0	117	10
4	t=0	0, 0 <sup>5</sup>	117	10

1: time exposed to 0.6% slurry before measurement of ingestion

2: ratio of wet weight of slurry to that of filtered sea water

3 and 4: The initial densities of prey and predator (*Gymnodinium sanguineum* and *Calanus pacificus* in experiments 1, 3, and 4, and *Gonyaulax polyedra* and *Acartia* spp. in experiment 2)

5: water in which slurry had been soaked for 24 hours

\*: Incubation bottle size (500 ml in experiments 1, 3, and 4, and 270 ml in experiment 2)

## FIGURE CAPTIONS

Fig. 1. Ingestion rates of Gymnodinium sanguineum by Calanus pacificus as a function of the slurry concentration. Symbols represent treatment means  $\pm$  1 S.E. Relations are fitted by the curve linear regression.  $IR$  (prey eaten Calanus<sup>-1</sup> h<sup>-1</sup>) =  $183 \times e^{(-5.42 \times SC)}$  ( $R^2 = 0.831$ ); where SC = slurry concentration.

Fig. 2. Ingestion rates of Gonyaulx polyedra by Acartia spp. as a function of the slurry concentration. Symbols represent treatment means  $\pm$  1 S.E.

Fig. 3. Ingestion rates of Gymnodinium sanguineum by Calanus pacificus. Symbols represent treatment means  $\pm$  1 S.E. Black bars - Incubated without slurry in both Day 1 (t=0 in Table 1) and 2 (the initial G. sanguineum concentrations in Day 1 and 2 were 183 and 117 cells ml<sup>-1</sup>, respectively). Open bars - with 0.6% slurry (wet weight:wet weight) in Day 1 and without slurry in Day 2. Gray bar - in slurry-free sea water in which slurry had been soaked for 24 hour and then removed by filtration.




Fig 1





Fig 2



1  
2

## **APPENDIX F**

### **FISH INTERACTION REPORT**

Source: Fish Interaction Studies.  
San Diego, California  
National Marine Fisheries Service, 1995

**Effects of Dispersed Paper Effluent on the Filter-Feeding Capacity of Pacific sardine, *Sardinops sagax* : a preliminary study.**

**Russ Vetter**

**Genetics and Physiology Group**

**Southwest Fisheries Science Center**

**National Oceanic and Atmospheric Administration**

**I. Background**

Small, pelagic, clupeoid fishes (sardine, anchovy, herring, menhaden) are an important link in most coastal pelagic ecosystems. These organisms harvest the carbon found in phytoplankton and zooplankton and convert it to a large standing stock of small migratory, schooling fishes. In turn, these species are the forage base for larger predacious fishes (e.g. tuna, salmon, rockfish), seabird populations (pelicans, gulls, terns), and marine mammals (dolphin, seal, sea lion). On a weight basis, clupeoid fisheries are the most important fisheries world-wide. Off the west coast of North America northern anchovy, *Engraulis mordax* and Pacific sardine, *Sardinops sagax*, vary in abundance during different climate regimes but together represent the major stocks of clupeoid fishes. Presently the sardine biomass is estimated to be about 340,000 metric tons.

Fish such as Pacific sardine can reach such high biomass because they feed lower on the food chain than larger predacious fishes. They typically subsist on a diet of phytoplankton and zooplankton. Sardine can feed by two different methods. If the prey organisms are large enough, sardine will strike at and ingest individual particles. If the organisms are smaller but abundant, they will filter-feed. They use cartilaginous extensions on their gill rakers to filter water as it passes through the gills, and collect the organism trapped on the gill rakers. The size of the particles trapped depends on the size of the seive created by the gill rakers. This varies with species, and the size of the individual fish (Blaxter and Hunter 1982). Compared to invertebrate filter-feeders (clams, oysters, mussels), little is known about the filtration process in clupeoid fishes.

We have been asked to design experiments to test for possible effects of finely dispersed paper fibers. These fibers will potentially be released from Naval ships as a method of routine disposal of paper waste generated at sea. We have begun by examining the potential for different concentrations of these fibers to interfere with normal filter-feeding in Pacific sardine, *Sardinops sagax*.

## II. Experimental Design

To test for possible effects on filter-feeding we devised a series of experiments where groups of sardine were exposed to small prey (*Artemia nauplii*) that stimulate filter-feeding. We then monitored the disappearance of prey in four tanks containing schools of equal sized sardine. The four tanks were a control tank receiving prey only, and three tanks receiving prey plus three different concentrations of paper effluent. The experiment ran 14 days. Fish were exposed to the effluent every other day. We measured the rate of disappearance of prey, weight gain or loss, and presence of paper in the stomach and intestines of the fish.

The paper concentrations tested in this initial experiment were 30, 15 and 3 mg dry wt of paper / l of seawater. These concentrations were chosen as a reasonable range to bracket expected concentrations from the point of release from the ship, down to the expected concentration where dilution driven by the turbulence of the ship's wake would dissipate. Beyond that point, the rate of further dilution is dependent on ambient oceanographic conditions (these lower dilutions will be tested in the next series of experiments).

To determine the rate of prey disappearance we needed to maintain the tanks in a closed, recirculating mode. A feeding trial was done over an 8 hour period. Prey or prey plus effluent was added and samples taken throughout the 8 hour period to monitor the disappearance of prey under the various conditions. In the present experiment the two upper concentrations appeared to cause respiratory distress and it appeared that the fish would not survive two weeks if exposed to the paper effluent every day. We elected to subject the fish to the effluent every other day but to feed them every day. When the fish were not being tested or fed (the remaining 16 h each

day), the tanks were maintained in a flow-through mode.

We generated 8 feeding trials for the 14 day period. At the end we measured and weighed the fish, recorded growth information, and examined the condition of the gills and the digestive system.

### **III. Methods and Materials**

#### **1. Collection, Maintenance, and Measurement of Sardine**

Sardines were transported from a San Diego bait dealer in February of 1995 to the laboratory and held in circular, vinyl-lined tanks measuring 5 m in diameter with 0.7 m of water. Sardine were maintained on a diet of Oregon Moist pellets.

Using an electrical top-loading balance and a standard measuring board, 240 Pacific sardines (*Sardinops sagax*) were weighed (wet wt.) and measured (SL,FL,TL). To minimize bias, we alternated tanks for each addition of ten fish. In other words, ten fish went into tank 1A, ten fish into 2A, etc., until all four tanks contained 60 fish. Fish were anesthetized in MS-222 using 30 mg/liter.

At the end of the last 8 hour exposure period, fish were anaesthetized and frozen to retain stomach contents. Fish were weighed whole while frozen. Later groups of 20 of each group were thawed and the gastro-intestinal tracts weighed, dissected, examined for stomach contents, and preserved. The relative amount of paper in the stomach and intestine was noted. Whenever a sardine was found dead in a tank, it was promptly removed, weighed and measured. Fish were then frozen for future examination.

#### **2. Preparation, Addition, and Calculation of Artemia Concentrations**

Anchovies consume 1.7% to 5.1% of their body wt. per day (Leong and O'Connell, 1969). We assumed the same approximate metabolic demand for sardine. A 12 cm

sardine filters 270 liters per day (Yoneda and Yoshida, 1955). These basic measurements were used to calculate the number of Artemia nauplii needed to sustain a 16 gm sardine stocked at a density of 60 sardine per tank. Fish were fed daily with 24-48 hour old Artemia nauplii at a rate of 1.7% body weight. We presented the fish with the minimum (233 nauplii/l) needed to maintain weight. The calculation is presented in Table 1. The nauplii were added to tanks at 0800 on all days.

### **3. Preparation, Addition, and Calculation of Paper Concentrations**

To determine the paper effluent dosage protocol we first determined a dry wt to wet wt conversion. The results of this determination for the first shipment of paper (which was used for all experiments described here) is presented in Table 2a. Based on a dry to wet conversion of 6.55, 24 hours prior to each sampling day the frozen paper waste was weighed, thawed and sea-water was added to make a 12-liter mixture. A mechanical stirrer was used periodically throughout the day to help "fluff-up" paper.

Paper was added to each tank (except 4A) and allowed to mix for 10-15 minutes before adding artemia at time 0. The calculations for the different paper concentrations are presented in Table 2b.

Tank 1A - 30 mg dry wt. /liter

Tank 2A - 15 mg dry wt. /liter

Tank 3A - 03 mg dry wt. liter

The tanks were monitored to determine if the paper effluent remained suspended throughout the exposure period (Fig. 2 a-h, bottom panels).

### **4. Exposure Tanks**

Four identical fiberglass tanks, 2 m in diameter, were plumbed to a depth of 0.64 m each, creating a water volume per tank of 1700 liters. Tank bottoms sloped

slightly to a center outlet covered with a PVC "cap" perforated with holes for draining. Drain leads to a standpipe adjacent to tank such that the height of the standpipe determines the water level in tank (Figure 1). All four tanks were rigged with the same materials in order to be as identical as possible. In-line water filters used were 5-micron Cuno filters, changed every other morning.

When fish were not being exposed to paper or food (16h each day), the tanks were kept on a flow-through regime of fresh, ambient seawater. Flow rates were four liters/minute.

When fish were being exposed to paper and/or food the tanks were kept in a closed but recirculating mode. Submersible pumps (1/30 HP, epoxy-coated, 500 gph) were used on the tank bottom to assist in keeping paper waste suspended in water column (Fig. 1). In addition air lift in the center of the tank lifted material from the bottom of the tank and resuspended it at the top (Fig. 1). The airlift also provided full oxygen saturation to the tanks. In either closed or open mode the tank temperatures remained between 15.2 and 17.3 degrees C throughout experiment.

At the end of each sampling day, tanks were drained down to approximately 3 inches in depth to assure a clean tank before beginning the next sample. At the beginning of each sampling day, with the tanks clean and re-filled, the incoming water was shut off, the submersible pumps turned on, and an air-line with airstone was put inside the standing drain-pipe to lift and circulate any paper waste tending to accumulate on tank bottom. By using pumps and airstones we were able to keep the paper waste suspended and evenly distributed throughout the tanks.

## **5. Sampling**

Using a 250 ml plastic beaker, four 250 ml samples were collected from 4 places in each tank (total sample = 1000 ml). In each quadrant of each tank and midway between the center point and the inside wall, an inverted 250 ml beaker was lowered to approximately three inches below the surface. The beaker was then turned upright and removed. The four 250 ml samples were poured into a 1000 ml plastic graduated cylinder labelled specifically for that tank.

Samples were taken at six time points during the 8-hour duration of each experiment. Time points were 1,2,3,4,6 and 8 h from the start. A subjective assessment of feeding behavior was recorded at time of each sample.

## **6. Filtering and Counting Samples**

Prior to beginning an experiment, filter papers were dried, weighed and given an i.d. number before being placed in a desiccator. After collecting a 1000 ml sample from each of four tanks, a labelled and pre-weighed filter was placed in a clean Buchner funnel in vacuum filtering manifold. Upon prewetting filter using seawater from rinse bottle, sample was slowly poured into funnel, keeping an inch or two of sample water in funnel at all times to help distribute the sample evenly. After sample has been completely filtered, vacuum pump was turned off and the filter paper carefully removed. The filter paper was then placed in a petri dish under a dissecting microscope and the number of nauplii counted and recorded.

Once counted, the filters were then placed on a clean container and placed in a drying oven at 60 °C. Upon having dried for a minimum of 72 hours, the filters were removed from the oven, placed in a desiccator to cool, and weighed again. By subtracting the initial filter paper weight from the final filter paper weight gave us the weight of paper waste plus brine shrimp contained in our 1000 ml sample.

## **IV. Results**

### **1. Behavior and Appearance**

There is an obvious difference in the way the gills are flared during normal breathing and when filter-feeding. We used three subjective criteria to assess visual signs of filtering behavior: 1. actively filtering, 2. passively filtering (occasional gulping) and 3. little or no feeding activity. At the two highest paper concentrations the filtering activity was a 1 when the food was first introduced in the first hour but dropped to a two in the second hour and then 3 for the remainder of the 8 hours even though

abundant prey was available. In the control tank feeding was always active (1) until the prey were gone. At the two highest paper concentrations fish did not maintain tight schooling behavior. Occasionally an individual would display lethargic or disoriented swimming behavior.

At all three paper concentrations there were fish with lumpy abdomens. At the end of experiment when fish with this outward appearance were dissected there were lumps of accumulated paper in the intestine that corresponded to the protrusions on the skin.

## **2. Feeding**

The results from the eight feeding trials are presented in Fig. 2. All three paper concentrations had a readily observable negative effect on the ability of Pacific sardine to filter-feed on *Artemia nauplii*. There was a dose-dependent effect on filtering success with the highest concentration (30mg dw/l) almost completely inhibiting successful feeding on *Artemia nauplii*.

All treatments ingested the paper along with the prey, Fig. 3. The highest ingestion rate was in the group receiving the lowest levels of paper (3 mg dry wt/l).

## **3. Growth and Mortality**

All groups lost weight during the experiment. The average initial wts for the 4 groups were 16.19,16.32,16.10,and 16.12g. At the conclusion of the experiment the average wts were respectively 14.0,14.4,14.3, and 13.9 g. There were no significant differences between the four groups.

There were significant differences in mortality. All three treatment groups had higher mortality rates than did the control, (Fig. 4).

## **V. Discussion and Preliminary Conclusions**

At the three concentrations tested in these preliminary experiments. The paper

effluent had a strong effect on the filter-feeding of Pacific sardine. The effects at the two highest concentrations were not unexpected and represent a worst case scenario not likely to be encountered at sea except in the immediate area around a ship. The effects noted at the lowest concentration (3 mg/l) may be of greater concern if they represent potential environmental concentrations.

The greater concentration of paper in the stomachs of fish exposed to the lowest concentration is due to the behavior of the fish under the different treatments. At the two highest concentrations the fish are visibly stressed by the concentration and spent less time filter feeding. Thus they received less food and also less paper. At the lowest concentration the fish were measurably affected, but they did eventually remove all of the prey ( Fig. 2). However, they consumed large amounts of the paper along with the prey. The harmful, or even beneficial, effects of consuming the paper fibers is not known, but the protrusions of the gut in fishes with paper plugs in their intestines is an alarming result. We found no evidence for digestion of the paper fibers. Fecal pellets that looked exactly like undigested but condensed pellets of paper were commonly observed at the bottoms of the three exposed tanks.

Differences in mortality were observed (Fig. 4), but the causes of mortality are not known. Fish from the three paper treatments were not significantly lighter, and starvation is not considered a likely cause of the mortality. All fish received a full ration in the absence of paper on alternate days and there were no differences in total weight. Even if we can account for some of this weight as paper in the gut, the weight differences between groups were not significant.

Our following experiments will attempt to: 1. find the no-effect level, 2. investigate differences in weight loss in longer term experiments, 3. gain biochemical and microscopical insights into the causes of the observed mortality.

## VI. References and Background Literature

Blaxter, J.H.S., Hunter, J.R. 1982. The Biology of the Clupeoid Fishes, offprint from "Advances in Marine Biology", Vol. 20, pp 1-223.

Hunter, J.R., Dorr, H. 1982, Thresholds for filter feeding in northern anchovy, *Engraulis mordax*, CalCOFI Rep., /vol. XXIII.

Yoneda, Y., Yoshida, Y. 1955. The relation between the sardine and the food plankton - I. On the food intake by *Sardinops melanosticta*, Bulletin of the Japanese Society of Scientific Fisheries, Vol. 21, No. 2

Koslow, J. A., 1981. Feeding selectivity of schools of northern anchovy, *Engraulis mordax*, in the southern California bight, Fishery Bulletin: Vol. 79:131-229

Leong, R.J. H., O'Connell, C.P. 1969. A laboratory study of particulate and filter feeding of the northern anchovy(*Engraulis mordax*). J. Fish. Res. Bd. Canada 26: 557-582.

## VI. Figure Legends

Fig. 1 This is a the basic configuration of tanks showing the water filtration, air lifts, etc.


Fig. 2 This is the basic data for the 8 days of feeding trials (the first day and every other day thereafter). Squares are 30 mg/l, circles are 20mg/l, triangles are 3 mg/l and diamonds are the control. Feeding rates are plotted two ways, the second nauplii/fish corrects for mortality during the experimental period. The paper conc traces are best used to verify that there were no long term changes in concentration throughout the 8h of the experiment. This is a measure of how well the air lift and pumps kept the paper effluent resuspended and evenly distributed.

Fig. 3. This is a chart of the stomach weights for the four treatments. All treatments contained paper in the stomach, but the highest concentrations were in the low (3 mg/l) treatment.







Fig. 4. All four tanks began with 60 fish. Mortality was lowest in the control. There were no differences between the three paper treatments.







## METHODS & MATERIALS







SARDINES WERE TRANSPORTED FROM A SAN DIEGO BAIT DEALER IN FEBRUARY OF 1995 TO THE LABORATORY AND HELD IN CIRCULAR VINYL-LINED TANKS, 5 METERS IN DIAMETER WITH 0.7 METERS OF WATER. ~~FOR THE~~ FOUR IDENTICAL FIBERGLASS TANKS, 2 M. IN DIAMETER, WERE PLUMBED TO A DEPTH OF 0.64 M. EACH, CREATING A WATER CAPACITY OF 1700 LITERS. TANK BOTTOMS SLOPED SLIGHTLY TO A CENTER OUTLET COVERED WITH A PVC "CAP" PERFORATED WITH HOLES FOR DRAINING. DRAIN LEADS TO A STAND-PIPE ADJACENT TO TANK (HEIGHT OF STANDPIPE DETERMINES WATER LEVEL IN TANK.) (SEE FIG. 1)









- ALL 4 TANKS WERE RIGGED WITH THE SAME MATERIALS IN ORDER TO BE AS IDENTICAL IN ALL ASPECTS AS POSSIBLE.
- IN-LINE FILTERS USED WERE 5-MICRON CUNO-FILTERS, CHANGED EVERY OTHER MORNING.
- AMBIENT SEAWATER FLOW RATES INTO TANKS AT 4 LITERS/MINUTE.
- TANK TEMPS. REMAINED BETWEEN 15.2°C - 17.3°C THROUGHOUT EXPERIMENT.
- SUBMERSIBLE PUMP SPECS. = 1/30 H.P., EPOXY-COATED, 500 GPH











**Fig. 3 Stomach weights for the four treatments**



**Fig. 4 Percent mortality at the end of 14 days**





**Table 2a. Measurements and calculations used to determine the conversion of paper...dry wt. to wet wt.**

		(1st lot of paper material received)									
Total drying time: 950322(0930)-950411(1400)[20days, 4.5 hrs.]											
		Date: 950328(0845)		Date: 950329(1005)		Date: 950411(1400)					
		# 1		# 2		# 3					
Number	Filter	Tare wt. (g)	Paper	Wet Wt. (g)	Tare + Dry Wt. (g)	Tare + Dry Wt. (g)	Tare + Dry Wt. (g)	Dry Wt. (g)	% water	% Dry Wt.	Dry to Wet Conv.
1		1.5965		7.0690			2.6490	1.0525	85.11	14.89	6.72
2		1.5893		7.1378			2.7342	1.1449	83.96	16.04	6.23
3		1.6231		7.1278			2.6799	1.0568	85.17	14.83	6.74
4		1.5807		7.1649	2.6759	2.6729	2.6748	1.0941	84.73	15.27	6.55
5		1.5955		7.1642			2.6282	1.0327	85.59	14.41	6.94
6		1.5806		7.1914	2.6792	2.6777	2.6822	1.1016	84.68	15.32	6.53
7		1.6231		7.2478			2.7533	1.1302	84.41	15.59	6.41
8		1.6137		7.1217			2.7075	1.0938	84.64	15.36	6.51
9		1.5937		7.1266			2.7139	1.1202	84.28	15.72	6.36
10		1.5809		7.1514			2.5998	1.0189	85.75	14.25	7.02
11		1.6262		7.1747			2.7793	1.1531	83.93	16.07	6.22
12		1.5844		6.8342			2.5659	0.9815	85.64	14.36	6.96
13		1.6252		6.8652			2.6755	1.0503	84.70	15.30	6.54
14		1.5739		7.1573	2.6879	2.6845	2.6847	1.1108	84.48	15.52	6.44
15		1.5810		7.1907			2.7362	1.1552	83.93	16.07	6.22
16		1.6136		7.9458			2.8332	1.2196	84.65	15.35	6.52
17		1.5953		7.1571			2.6327	1.0374	85.51	14.49	6.90
18		1.6023		7.0713			2.6557	1.0534	85.10	14.90	6.71
19		1.5969		7.2978			2.7893	1.1924	83.66	16.34	6.12
20		1.5981		7.3858	2.7785	2.7776	2.7800	1.1819	84.00	16.00	6.25
								Mean	84.70	15.30	6.55
								Stdev	0.63	0.63	0.27


**Table 2b. Calculations used to determine the wet wt. concentrations of paper needed for the three treatments.**

Tank No.	Treatment	% dose	Concentration mg dry paper/l	Conv. dry to wet	Tank capacity (liters)	Concentration mg wet paper/l	Concentration gm wet paper/l	Wet wt. paper per tank (gms)	liters per tank of 12 l mixture	
1A	High	100	30	6.3	1700	189	0.189	321.3	7.5	
2A	Medium	50	15	6.3	1700	94.5	0.0945	160.65	3.75	
3A	Low	10	3	6.3	1700	18.9	0.0189	32.13	0.75	
4A	Control	0	0	****	1700	0	0	0	0	
<b>Total:</b>							<b>48</b>		<b>514.08</b>	<b>gms in 12 liters</b>

Vetter New Data

.01 - 1mg

National Oceanic and Atmospheric Administration  
Southwest Fisheries Science Center  
8604 La Jolla Shores Drive  
La Jolla, CA 92093



Genetics and Physiology Program

FAX (619) 546-5656  
Info (619) 546-7000

Date: 7-10-95  
Number of Pages: 5  
Addressee: Stacey Cumbie  
Addressee's FAX: 553-6305  
Originator: Russ Vetter  
Originator's Phone: 619-546-7125

**Comments:**

Stacy, here are the results from the low-level experiment. It is pretty much the same experimental design as before except with higher initial artemia levels. Only the highest concentration (1.0 mg/l) showed an effect. This does not surprise me since 0.1 mg and .01 mg/l are very low concentrations for some thing like paper. The time scale is different than for the first experiment but the results for the 1 mg treatment do suggest that the previous results for the lowest concentration in expt 1 (3.0 mg/l) are real. Let me know what you think.

Regards

## **Lower Threshold Filtration Experiments.**

### **I. Experimental Design**

1. # of fish : 60 per treatment

#### **2. treatments**

1. control
2. 1.0 mg/l dry wt
3. 0.1 mg/l
4. 0.01 mg/l

all treatments received a food ration of 863 nauplii per liter.

#### **3. exposure conditons**


Fish were exposed to the different conditions begining at 9:00 in the morning and were sampled every 15 minutes for two hours until food was gone from all treatments. Fish were exposed every day for 14 days. Filtering efficiency was measured every other day

### **II. Results**


1. no mortality under any treatment condition
2. a measurable sublethal effect on filtering efficiency at 1 mg/l, no effect at .1 and .01mg/l
3. There were no lasting effects on filtering efficiency. Fish were tested for two additional days and all groups were like the controls.

### **III. Conclusion**


The no-effect level for this experiment was between 1 and 0.1 mg/l




DAY 1 time (hr)




DAY 3 time (hr)




DAY 5 time (hr)




DAY 7 time (hr)



DAY 9





DAY 11



DAY 13

### Recovery Trials: After 14 Days All Groups Tested With Artemia Only.



## **APPENDIX G**

### **SHIPBOARD METAL WASTE DISCHARGE CORROSION REPORT**

Source: Shipboard Solid Waste Discharge Corrosion Study.  
San Diego, California  
Naval Command, Control & Ocean Surveillance  
Center, RDTE Division,  
Code 522, 1995

5216  
Ser 815/7  
21 Jun 95

MEMORANDUM

From: W. E. Glad, Code 02T  
To: S. L. Curtis, Code 522  
Via: Head, Materials Science Branch, Code 815 *WEG*

Subj: SHIPBOARD SOLID WASTE DISCHARGE CORROSION STUDY

Encl: (1) Solid Waste Discharge Corrosion Study of 21 Jun 95

1. Enclosure (1) is a report on the characterization and corrosion rates of metal food preparation waste that is commonly discharged into the ocean by navy ships. This report was prepared by the Code 815 materials laboratory at the request of Code 522. The investigators who took part in this study were Gordon Chase (Code 02T), Wayne Glad (Code 02T) and Tom Knoebel (Code 815).

2. Enclosure (1) contains the results of laboratory characterization of waste materials that were supplied by Code 522 and a summary of the literature concerning the corrosion rates of these materials. The waste consisted of tin plated steel food cans and lids and standard aluminum beverage cans. The lifetimes of this waste in the ocean environment were estimated from the materials properties (composition, coatings, dimensions) that were measured in the laboratory and the corrosion rates that were found in the literature.

*W. E. Glad*  
W. E. GLAD

21 JUN 95

SHIPBOARD SOLID WASTE DISCHARGE  
CORROSION STUDY

Enclosure (1)

# Solid Waste Discharge Corrosion Study

## Introduction and background

The purpose of this study is to quantitatively assess the impact on the marine environment of metal food preparation waste that is thrown over the side of Navy ships . This metal waste is rinsed, shredded, packed in burlap sacks, and thrown overboard. Since this waste will corrode and release its constituent elements to the environment it is important to know the composition of the waste and the rate which with these materials corrode. This will also give information about the lifetime of the waste on the ocean floor. Because conditions in the marine environment may differ from location to location, it is also important to know how the corrosion rates depend on these varying conditions.

## **Corrosion in Sea Water**

### **Types of Corrosion**

Fontana<sup>1</sup> describes eight types of related but somewhat different types of corrosion. Four of these types are important for this study. **Uniform attack** occurs as direct oxidation over a wide area of metallic surface. Uniform attack allows for corrosion at rates that can be measured relatively reliably. The corrosion rate (expressed as mass of metal oxidized per unit time) is proportional to the surface area of the corroding metal. To a large extent, mild steel undergoes uniform attack when it is immersed in sea water. **Pitting** is a kind of localized attack that often occurs on metals, such as aluminum, that naturally resist corrosion due to the formation of passive oxide films. While the eventual rate of corrosion in a pit is determined by the rate of reduction of oxygen (usually at a surface away from the pit), a pit requires some kind of initiation process to start the pit formation. Thus the overall rate of corrosion on a pit forming metal like aluminum may depend on factors other than the dissolved oxygen concentration. Pitting is assisted by the presence of chloride ion, so salinity may also be a factor in pitting corrosion rates. **Crevice Corrosion** (and related **filiform corrosion**) occurs in small openings such as joints and under defects in non-metallic coatings. Corrosion on coated metals will begin with crevice or filiform corrosion that progresses under the coating and eventually results in the rupture of the coating due to the build up of corrosion products. Filiform corrosion is common on coated food and beverage cans that are exposed to the atmosphere. In sea water this type of corrosion would be expected to lead to the destruction of coatings and the exposure of the metal underneath to direct attack.

### **Corrosion measurement**

The most reliable information about corrosion in sea water comes from the direct exposure of samples in the ocean environment. Well characterized samples (for chemistry, metallurgy and surface condition) are carefully measured and weighed before exposure. Good studies also record the exposure conditions (O<sub>2</sub> concentration, pH, temperature, flow rates) and their variation in time. After exposure the samples are cleaned to remove the corrosion products and any marine fouling, then weighed. Cleaning methods can be mechanical, chemical, or both, but standard methods are used to assure reproducibility in the measurements.<sup>2</sup> Corrosion rates are usually reported in mils (thousandths of an inch) per year (mpy) via the formula:

$$\text{mpy} = \frac{534 W}{\rho A T}, \quad (1)$$

where  $W$  is the weight loss in milligrams,  $\rho$  is the density of the sample in grams/cm<sup>2</sup>,  $A$  is the area of the specimen in square inches and  $T$  is the exposure time in hours. Reporting corrosion rates in this manner is

useful for the assessment of corrosion damage for structural materials. Since we are interested in the total amount of material corroded we must take into account the surface areas and densities of our samples.

Because direct exposure testing can be expensive and limited in the number of different conditions that can be realistically experienced, testing is also conducted in the laboratory. In addition to simple immersion tests, electrochemical studies that measure corrosion potentials and currents are also performed. These studies can be useful in elucidating corrosion mechanisms, as conditions such as temperature, oxygen concentration, and pH can be varied over wide ranges in the laboratory.

### **Sea water corrosion studies**

The most widely cited studies of corrosion in sea water were undertaken by the Civil Engineering Laboratory, Naval Construction Battalion, Port Hueneme, California. The studies involved the exposure of about 20,000 samples of 475 different metal alloys at three depths in the Pacific Ocean. An exhaustive summary of the results was published.<sup>3</sup> Some of these results, and the results of many other studies are summarized (with references) in *Corrosion of Metals in Marine Environments*,<sup>4</sup> report compiled by the Metals and Ceramics Information Center. A useful study that compared the corrosion of mild steel in polluted and unpolluted sea water was performed by Shimada *et. al.*<sup>5</sup> A study that examined the effect of water flow velocity on the corrosion of steel was produced by Peterson and Lemox.<sup>6</sup> The consensus of these studies is that the main factor influencing the corrosion rate for carbon steel is the dissolved oxygen concentration. This is subject to the caveats that excessive flow rates will increase the corrosion rate substantially, and that in polluted waters with high sulfide concentration (and a correspondingly low oxygen concentration) the corrosion rates will be significantly higher than normally predicted from the oxygen concentration.

In addition to Reinhart<sup>3</sup>, studies on the corrosion of aluminum include a five year field study by Ailor<sup>7</sup>, and laboratory study by Dexter.<sup>8</sup> There appears to be an increase in the corrosion rate of aluminum with depth, but the reasons are poorly understood.

### **Containers**

Food containers are designed to keep the outside environment away from the food products. These containers must themselves prevent a breach of the container from corrosion that is initiated either from the outside environment or from the inside by the food product. Tin plated steel cans have been used for food preservation since the heyday of the British empire. Under most conditions tin is more noble than iron, and provides a barrier coating to prevent corrosion. The steel sheet used in tin plated containers is usually a mild steel with very low concentrations of additional elements. A mild steel has a carbon content of less than 0.2 % (in the case of steel for cans, less than 0.14%) and about 0.5 % manganese.

In some cases, the tin plating is not sufficient to prevent the corrosive action of some food products. In addition to the tin plating, organic coatings, called enamels in the industry, are also used. The enamels are often oleoresins (natural products) or epoxies (synthetic products), but other coatings can be used. Coatings can be used on the interior surface of the can only, but are sometimes present on both the interior and exterior surface. An interesting but slightly dated discussion about tin container technology is given in the *Metals Handbook*.<sup>9</sup>

Many beverages are distributed in aluminum cans. Modern aluminum cans are deep drawn from 3004 alloy sheet. The lids on the cans are stamped from type 5182 sheet. The 3004 alloy contains about 1% manganese and 1% magnesium. The 5182 alloy contains about 4% magnesium and 0.3 % manganese. Because of the very corrosive nature of some of the beverages, these cans also have organic coatings on the inside. The outsides of the cans, however, are usually coated only with decorative paint that does not completely cover the metallic exterior.

## Chemistry of alloy components in sea water

The chemistry of the corrosion products from these containers, in the ocean environment, is influenced by the same factors that control the rates of corrosion themselves. While a detailed discussion of the marine chemistry of these corrosion products is beyond the scope of this study, a simplified discussion of the fates of these products follows:

**Iron:** Because of the presence of dissolved oxygen in the ocean, iron in this environment exists almost completely in the Fe(III) oxidation state. In spite of the high concentration of chloride in sea water, chloride complexes of Fe(III) are not a factor in the distribution of iron. The very small solubility product of Fe(OH)<sub>3</sub> (about 10<sup>-37</sup>) guarantees that at the near neutral pH levels in sea water most iron exists as solid or colloidal Fe(OH)<sub>3</sub>. The net stoichiometry for the oxidation of iron to Fe(OH)<sub>3</sub> is:



Fe(OH)<sub>3</sub> is metastable, and must undergo dehydration



to produce more stable forms such as Fe<sub>2</sub>O<sub>3</sub> that are common in ocean sediment.<sup>10</sup>

**Manganese:** Manganese can exist in sea water in either the Mn(II), Mn(III) or Mn(IV) oxidation states, depending on the oxidative potential of the sea water. Since manganese is only a trace element in sea water, yet is common in ocean sediments, most manganese must precipitate. The major precipitates of manganese are believed to be MnCO<sub>3</sub>, MnO(OH), and MnO<sub>2</sub>. The net reactions would be:



**Aluminum:** Aluminum is oxidized to Al(OH)<sub>3</sub> in neutral solution via:



Aluminum chemistry in the oceans can be quite complex. Aluminum is a major element in the earth's crust and in ocean sediments. To a large extent, at ocean pH levels, any aluminum produced by the corrosion of metallic aluminum probably ends up as insoluble Al(OH)<sub>3</sub>. This aluminum hydroxide may also eventually dehydrate to the oxide (Al<sub>2</sub>O<sub>3</sub>) as does iron.

**Magnesium:** Magnesium is oxidized via:



Magnesium is a major component of sea water, present at about the 1300 µg/ml level. A significant portion may be present as the ion pair  $\text{MgSO}_4$ .

**Tin:** Tin is easily oxidized from the Sn(II) state to the Sn(IV) state. At ocean pH levels the Sn(IV) probably exists as insoluble  $\text{SnO}_2$ .

With the exception of tin, these major corrosion products are common constituents of ocean sediments and the earth's crust in general. As average oxide percent of ocean sediment, aluminum is present at 12%, iron at 6.5%, magnesium at 2.3% and manganese at 0.9%. Tin is present at about the 11 ppm level.<sup>10</sup>

## **Experimental Methods**

### **General Analysis Methods**

A general discussion of the materials analysis methods used in this study is given here to provide background for the specific procedures that follow.

#### **ICP**

Inductively Coupled Plasma (ICP) spectroscopy is used in the materials laboratory to determine the composition of metal alloys. Metal samples are dissolved in mineral acids and diluted to known volumes. The solutions are nebulized into a continuously running plasma where the constituent metal atoms are excited to emit visible and ultraviolet light. The light is dispersed by a grating and detected by photo multiplier tubes. These atomic emission lines are usually well resolved. The intensities of the lines can be compared with the intensities from synthetic solution standards, allowing for a quantitative analysis of the dissolved material. The ICP provides measurement precision of about 1% relative, and is sensitive to concentrations in the low parts per billion for many elements. Metal alloys can be analyzed for major, minor, and trace elements using the ICP.

#### **Carbon/Sulfur Analysis**

Due to the insolubility of some metal carbides and relatively poor sensitivity for carbon, the ICP is not used to determine the carbon content of steels. Instead, a dedicated carbon/sulfur analyzer is used. Samples of steel are combusted in an induction furnace in a stream of oxygen. The carbon dioxide that is produced is measured using infrared absorption spectroscopy. Sulfur is similarly determined from the sulfur dioxide that is produced. Analysis times are usually less than a minute, and experimental precision is better than 1 % relative.

#### **Metallography**

Metallography is the examination of polished and sometimes etched metal samples under the microscope. Samples are mounted in plastics for support during grinding and polishing. The polished samples are examined on an inverted stage, incident-light, metallurgical microscope. The microscope is equipped with either bright or dark field illumination and has a polarized light illuminator and a Numarski prism attachment for enhanced phase contrast. The image may be viewed through binocular eyepieces and photographed. Metallographic methods are useful for examining the microstructure of metals and making microscopic measurements.

## Electron Microscopy and Energy Dispersive X-ray Analysis

The scanning electron microscope (SEM) uses a focused electron beam to image specimens. The SEM is capable of greater depth of field, higher magnification, and better resolution than optical microscopes. Non-conductive specimens must be made conductive to be imaged in the SEM. In our laboratory non-conductive specimens are sputter coated with a thin layer of gold. Samples in the electron beam of the SEM are induced by the beam to emit x-rays at energies that are characteristic of their elemental composition. Detection and analysis of these x-rays allow nondestructive qualitative and semi-quantitative elemental analysis of the sample. The spatial resolution of x-ray analysis in the SEM is on the order of 1 micron. Elements at the 0.2 percent level and above can be detected using this technique.

**Table 1**  
**X-ray Fluorescence Energies For Selected Elements**

Element	Energy (keV)								
C	0.277	S	2.31	Fe	6.4	Kr	1.59	Ag	2.98
N	0.392	Cl	2.62	Co	6.92	Rb	1.69	Au	2.123
O	0.525	Ar	2.96	Ni	7.47	Sr	1.81	Cu	0.93
F	0.677	K	3.31	Cu	8.03	Y	1.92	Fe	0.705
Ne	0.852	Ca	3.69	Zn	8.62	Zr	2.04	Pb	2.346
Na	1.04	Sc	4.09	Ga	9.24	Nb	2.16	Sn	3.414
Mg	1.25	Ti	4.51	Ge	9.88	Mo	2.29	Zn	1.022
Al	1.48	V	4.95	As	10.52	Ru	2.55	Cd	3.538
Si	1.74	Cr	5.41	Se	1.38	Rh	2.7	B	0.183
P	2.01	Mn	5.89	Br	1.48	Pd	2.84	Ba	4.466

## Analysis of Shipboard Metallic Waste

The samples of shredded food container waste that were received for analysis in the materials laboratory are shown in figure 1. Sample descriptions are given in table 2.

**Table 2**  
**Description of Samples in figure 1**

Sample Label	Description
A	Tin plated can lid
B	Tin plated can lid
C	Tin plated can body with attached bottom; can interior is white
D	Tin plated can body
E	Pepsi can body and top lid
F	Coca Cola can body and top lid
G	Sunkist Orange can body and top lid
H	Pepsi can body
I	Tin plated can body



Figure 1. Shredded Food Preparation Waste.

The samples consist of some tin plated can bodies and lids and some aluminum beverage cans. An initial visual examination showed that both the tin plated cans and the aluminum cans had organic coatings at least on their interior surfaces, and maybe on the exterior surfaces as well. The aluminum beverage cans have decorative paint on their outer surfaces. Pieces from samples B, C, and D were attached to an aluminum SEM stub with conductive copper tape and sputter coated with a thin layer of gold before examination in the SEM.

Pieces from samples C, D, E, F, and G were mounted on edge in filled epoxy hot pressed mounts. The mounts were ground through 600 grit silicon carbide paper. The samples were examined with direct and polarized light and selective photographs were made of the overall can wall configuration and of the coatings on the materials. The container thicknesses for the samples were measured using a calibrated eyepiece reticle. Some of the coatings were pigmented and active to polarized light, making them easily seen; others were not. The coatings on the aluminum cans were difficult to see, and the actual coating boundaries could not be positively identified. The samples were further polished with 6 micron diamond slurry on a cotton cloth and re-examined. Finally the samples were etched to reveal the can metal microstructure. Representative photomicrographs were made. The mount was then given a thin gold coating and examined in the SEM.

The aluminum cans had organic coatings on the inside of the body and organic coatings on both the insides and outsides of the lids. Most of the paint was removed from the metal surface by light grinding with silicon carbide abrasive paper. The organic coatings and remaining paint were removed from the samples by heating them in the flame of a Meeker burner. Samples weighing about 0.2 g were then dissolved in 5 ml concentrated  $\text{HNO}_3$ , 5 ml of concentrated  $\text{HCl}$  and 5 ml of deionized water. The solutions were diluted to 100.0 ml with deionized water and analyzed using the ICP.

Rectangular pieces of the tin plated can material were cut from samples A, C, and D. The areas of the pieces were measured. The samples were immersed in concentrated hydrochloric acid to remove the organic coatings and tin plating. The samples were attacked by the acid at edges and defects in the organic coatings. After about an hour the organic coatings were undercut and floated free from the samples; the tin platings had been dissolved. The samples were removed from the acid solution and washed with deionized water. The hydrochloric acid solutions were analyzed for tin using the ICP spectrometer. Pieces of steel from what remained of the samples were used for chemical analysis. Samples for ICP analysis were weighed, then dissolved in a mixture of 2.5 ml of concentrated  $\text{HNO}_3$ , 2.5 ml of concentrated  $\text{HCl}$ , and 2.5 ml of deionized water. Solutions were diluted to 50.0 ml and then analyzed with the ICP. Samples for carbon/sulfur analysis were weighed into crucibles and combusted in the carbon/sulfur analyzer.

## **Results**

### **Aluminum Containers**

The results of the ICP analysis for the aluminum containers are given in table 3 below.

**Table 3**  
**ICP Analysis Results for Aluminum Containers**  
(Values in weight Percent; Balance is Aluminum)

<b>Sample</b>	<b>Cr</b>	<b>Cu</b>	<b>Mg</b>	<b>Mn</b>	<b>Si</b>	<b>Ti</b>	<b>Zn</b>	<b>Fe</b>
Sample E body	0.005	0.181	1.09	1.01	0.132	0.018	0.05	0.40
Sample F body	0.007	0.160	1.06	1.08	0.162	0.024	0.03	0.36
Sample E lid	0.019	0.028	4.01	0.288	0.030	0.009	0.01	0.19
Sample F lid	0.010	0.009	4.10	0.291	0.030	0.014	0.01	0.15
3004 Specification	-	0.25 max	0.8-1.3	1.0-1.5	0.30 max	-	0.25 max	0.7 max
5182 Specification	0.10 max	0.15 max	4.0-5.0	0.20-0.50	0.20 max	0.10 max	0.25 max	0.35 max

The chemistry of the body alloy is consistent with aluminum alloy type 3004. The lid is consistent with type 5182.

The dimensions of the aluminum containers as measured by metallography are given in table 4 below.

**Table 4**  
**Aluminum Container Dimensions**

<b>Sample</b>	<b>Metal Thickness (in.)</b>	<b>Coating thickness (in.)</b>
Sample G side wall	0.0044	
Sample F side wall	0.0043	0.0002
Sample F upper side wall	0.0067	0.0002
Sample H, bottom	0.0128	
Sample H, lower side	0.0101	
Sample F, top	0.0096	
Sample F, pop top	0.0090	
Sample F, pull tab	0.0133	

Figure 2 is a micrograph of the aluminum side wall of sample F.




Figure 2. 800 X.

The inner coating is just barely visible. It is presumed that this is an organic coating of some type, but its exact nature is not known. Also visible in the microstructure of the aluminum are inclusions of aluminum-manganese intermetallic particles that are typical of this alloy type.

Figure 3 is a micrograph of the top to side joint from sample F.




Figure 3 50 X.

Modern aluminum beverage cans are two piece cans that have joints only on the top lid. The bottom of the can is formed from the same piece of metal as the can side walls.

## Tin Plated Steel Containers

Figures 4-11 show energy dispersive x-ray spectra from surfaces of the tin plated steel material that was examined in the SEM. The outside, "tin," surface of sample B showed both light and dark regions in the SEM. The dark regions were areas where a thick organic coating was present, as indicated by figure 4. The bright areas must have been locations where the organic coating was either very thin or missing altogether. The x-ray spectrum from the bright area (figure 5) shows intense tin and iron lines. Most likely this is from a thin (less than 1  $\mu\text{m}$ ) coating of tin that is penetrated by the electron beam to cause x-ray emission from the iron below. The inside can surface of sample B, which had a "gold" appearance, was covered with an organic coating, as is shown from the x-ray spectrum in figure 6. The x-ray spectrum from an area on the same side of sample B where the golden coating had been scraped away is shown in figure 7. Tin is present beneath this coating as well. The tin to iron intensity ratios indicate that the thickness of the tin is probably larger on the "gold" side of sample B than the "tin" side. The x-ray spectra of sample D (figures 10 and 11) were very similar to sample B. Samples B and D appear to be tin plated on each side and have organic coatings on top of the tin plate. Sample C is tin plated on one side (figure 8) and has a paint-like coating on the other side (figure 9) that contains some titanium dioxide pigment.




Figure 5. X-ray spectrum showing tin plate over iron on "tin" side of sample B.




Figure 7. X-ray spectrum showing a thicker tin plate on "gold" side than "tin" side of Sample B.




Figure 4. X-ray spectrum showing organic coating on "tin" side of Sample B.




Figure 6. X-ray spectrum showing organic coating on "gold" side of Sample B.

12




Figure 9. X-ray spectrum showing paint-like coating on "white" side of Sample C.




Figure 8. X-ray spectrum showing tin plate over iron on "tin" side of Sample C.




Figure 11. X-ray spectrum from a scratch on the "gold" side of Sample showing tin plate.




Figure 10. X-ray spectrum showing tin plate under organic coating on Sample D.

11.11

The chemical analysis results for the tin plated steel containers are given in Table 5 below:

**Table 5**  
**Chemical Analysis of Can Body and Lid Steel**  
 (Values in weight percent; Balance is iron.)

Sample	C	Al	Cr	Ni	Mn	Cu	Si	S	P	Mo	V	Co
D	0.110	0.052	0.016	0.013	0.51	0.014	0.010	0.007	0.013	0.000	0.003	0.001
A	0.109	0.022	0.041	0.021	0.43	0.014	0.003	0.009	0.004	0.001	0.002	0.002
Type L specification (maximum values)	0.13	-	0.06	0.04	0.60	0.06	0.020	0.05	0.015	0.05	0.02	0.02

The steel meets the specifications for Type L steel (see ASTM A 623), a type commonly used for tin plate food containers. This steel is a mild carbon steel that is very low in residual elements. Tables 6 gives results of the thickness measurements for tin plated steel containers.

**Table 6**  
**Tin Plated Steel Container Dimensions**

Sample	Metal Thickness (in.)	Coating thickness (in.)
Sample C side wall	0.0096	<0.005
Sample D side wall	0.0113	0.003
Sample D welded Seam	0.0131	0.0019
Sample D near seam	0.0067	<0.005
Sample C, bottom lid	0.0087	0.0002

Figure 12 is an optical micrograph of the side wall seam from sample D.




Figure 12. 100 X

The seam appears to be a resistance weld. It does not have the bent geometry one would expect from a solder joint. Neither is there evidence of lead in the x-ray spectrum from the joint. A fair amount of an organic coating had been applied around the seam. The x-ray spectrum from the coating is shown in figure 13 below.




Figure 13. X-ray spectrum from sample D Seam Coating

The coating appears to be a chlorinated organic material that is filled with barium sulfate. The barium sulfate may be present as an aid to automated x-ray inspection of the coating integrity.

Figure 14 is an optical micrograph of the bottom to side joint from sample C.




Figure 14. 20 X.  
Optical micrograph of Bottom to Side Joint of Sample C

This joint has typical geometry for a side to end joint. Figure 15 is the x-ray spectrum from the joint sealant.




Figure 15. X-ray spectrum from Joint Sealant in figure 14.

The sealant is chlorinated organic compound, that in this case is filled with aluminum and silicon compounds, probably oxides. (It is also possible that silicon is present as embedded silicon carbide from the grinding media).

Figure 16 is a micrograph of the sample C side wall that shows the white coating in cross-section.




Figure 16. 800 X.  
Optical Micrograph of Sample C side wall.

The coating is actually two layers. The layer closest to the steel appears to contain the most pigment. The x-ray spectrum of the layer closest to the steel is shown in figure 17 below.




Figure 17. X-ray spectrum from inner coating layer in figure 16

The layer actually looks like a paint that contains a significant amount of titanium dioxide pigment.

Figure 18 barely shows the presence of the tin plating on the etched surface of sample D.




Figure 18. 1250 X  
Optical micrograph of Sample D showing the tin plating.

The plating is too thin for its thickness to be measured optically. The measurement of tin plating thickness is usually done indirectly by stripping the tin from the sheet and using bulk chemical analysis to determine how much tin was present on a given surface area.

Tin plating thicknesses were estimated from the ICP analysis of the solutions containing tin stripped from samples A, C, and D by:

$$t = \frac{m_{Sn}}{\rho_{Sn}A} \quad (9)$$

where  $t$  is the total plating thickness,  $m_{Sn}$  is the mass of the tin from the analysis,  $\rho_{Sn}$  is the density of tin and  $A$  is the area of the sample. The results for three samples were:

**Table 7**  
**Tin Plate Total thickness from ICP Measurement**

Sample	Total thickness
A	45 $\mu$ inch
C	10 $\mu$ inch
D	73 $\mu$ inch

In the case of the sample A and D, the total thickness includes coatings on both sides. The measurement for sample C is for only one side, as the acid did not attack the paint like coating on the other side. Examinations in the SEM confirm the presence of a tin plating of around 30  $\mu$  inch adjacent to the paint surface on Sample C.

## Discussion

### **Aluminum Containers**

The aluminum container bodies are constructed of type 3004 aluminum. The lid materials are of type 5182 aluminum. The containers have an organic coating on the inside of the body. The lids appear to have organic coatings on both the inside and outside.

Reinhart<sup>3</sup> has no corrosion data for type 3004 aluminum. He does give data for type 3003-H14 (shown in Table 8), which does not have the 1% magnesium that is present in type 3004. Data is given for samples that were directly exposed to sea water and samples that were buried in bottom mud. Since magnesium additions have been shown to not affect the pitting potential of aluminum alloys<sup>4</sup>, the performance of type 3003 would be similar to that of type 3004. The H14 in the designation refers to some strain hardening to the alloy. The aluminum in beverage can bodies is severely strain hardened.

**Table 8**  
Corrosion rate data for 3003-H14 aluminum<sup>3</sup>

Depth (ft)	O <sub>2</sub> Concentration (ml/l)	pH	Salinity (ppt)	Temperature °C	Exposure time (days)	Rate Exposed (mpy)	Rate Buried (mpy)
5	5.9	8.1	33.51	15	181	1.1	-
5	5.9	8.1	33.51	15	366	1.0	-
5	5.5	8.1	33.31	15	588	1.2	-
2340	0.4	7.5	34.36	5	197	1.2	1.6
2370	0.4	7.5	34.36	5	402	1.4	1.7
5640	1.3	7.6	34.51	2.3	123	0.5	1.9
5640	1.3	7.6	34.51	2.3	751	2.3	2.5
5300	1.2	7.5	34.51	2.6	1064	2.0	1.9
6780	1.6	7.7	34.40	2.2	403	3.9	3.7

These data appear somewhat scattered, but the corrosion rates are higher at the greatest depths.

Similarly, there is no data for type 5182, but the data for most 5000 series aluminum alloys are very similar. The 5000 alloys tend to be more corrosion resistant than the 3000 series alloys. The composition of type 5086 (4 % Mg, 0.4% Mn ) is not too different from the lid alloy 5182.

**Table 9**  
Corrosion rate Data for 5056 Aluminum<sup>3</sup>

Depth (ft)	O <sub>2</sub> Concentration (ml/l)	pH	Salinity (ppt)	Temperature °C	Exposure time (days)	Rate Exposed (mpy)	Rate Buried (mpy)
5	5.9	8.1	33.51	15	181	1.2	-
5	5.9	8.1	33.51	15	366	0.8	-
5	5.5	8.1	33.31	15	588	1.6	-
2340	0.4	7.5	34.36	5	197	0.7	1.1
2370	0.4	7.5	34.36	5	402	0.6	1.3
5640	1.3	7.6	34.51	2.3	123	0.1	1.4
5640	1.3	7.6	34.51	2.3	751	2.0	-
5300	1.2	7.5	34.51	2.6	1064	0.9	1.2
6780	1.6	7.7	34.40	2.2	403	0.6	0.8

Electrochemical measurements of corrosion potentials and currents on aluminum by Dexter<sup>8</sup> indicate that the apparent increase in corrosion rates of aluminum with depth is probably due to the effect of reduced pH. He found that when oxygen concentration and pH are varied together, the effect of pH dominates the corrosion rate. Lower pH increases both the pit initiation rate and the pit growth rate.

The above data show very little effect of temperature on the corrosion of aluminum. While one study does indicate that the corrosion rate of 3004 aluminum is a factor of two higher at 25 °C than 10 °C, the corrosion rate of type 6061 alloy aluminum in tropical waters near the Panama canal zone is not significantly greater than the rate near Port Hueneme, California.<sup>4</sup> The passivity of the oxide films on aluminum may be diminished at higher temperatures, as the corrosion mechanism has been seen to change from pitting to uniform attack at higher temperatures.<sup>4</sup>

Corrosion data for shallow water immersion tests of 5086-H112 by Ailor give the following rates:

**Table 10**  
**Corrosion rates for 5086 aluminum from a five year study<sup>7</sup>**

Exposure time	Rate mpy
1 year	0.25
2 year	0.17
5 years	0.15

These rates are about an order of magnitude smaller than those observed by Reinhart. Since exposure conditions are not given in Ailor's study it is difficult to comment critically on the reasons for the difference. It may be that some experimental factor such as the cleaning method is involved.

## Tin Plated Steel Containers

While the samples that were examined are somewhat limited, some conclusions about the materials can be drawn. Food containers are made from a mild steel that has a low concentration of residual elements. The cans may or may not have tin platings. The cans may or may not have organic coatings. Some foods, such as tomatoes, are known to taste better if there is no organic coating on the inside of the can. Joints are sealed with organic sealants that can contain a variety of inorganic fillers. We observed no lead containing solders, but our sample was quite limited. The more corrosion resistant organic coatings (and the thicker tin plate) will usually be on the inside of the can. The measured thicknesses of the cans (with the exception of seam areas) range from about 0.009 inch to 0.013 inch. In the tin industry this is commonly referred to as 80-112 lb "plate". The samples that we examined had varying thickness of tin plating. Table 11 lists common tin plating thickness.

The tin plating thickness measurement of Sample D in table 7 is consistent with a D 100/25 differential coating. Sample A is consistent with a D 50/25 differential coating. Sample C is probably also from a D 50/25 differential coating. The thickness of the thin platings thus can vary from can to can (or between bodies and lids) and between the inside and outside. The thicker tin platings are usually on the inside of the can. A "base box" of tin plated sheet, for the thicknesses common in food containers, will weigh around 100 pounds. Thus, less than 1 percent of the mass of a tin plated steel food container is tin.

Corrosion rates for pure tin in seawater have been measured both at the surface and at depth.<sup>3</sup> The rates range from 8 mpy at the surface to 0.5 mpy at 5640 ft. The rates are somewhat correlated with the dissolved oxygen content. In seawater tin is cathodic (more noble than) to iron, so one might expect iron in contact with tin to corrode preferentially. However, with the very thin tin platings present on our samples (0.015 to 0.060 mil), it seems likely that the tin plating (where not protected by organic coatings) would be rapidly undercut and spalled away from the can surface by iron oxide corrosion products. The tin plating should corrode away quite rapidly in seawater.

**Table 11**  
**Common tin plating thicknesses<sup>11</sup>**

Designation	Tin Coating weight each surface (lb/base box)	Tin Thickness each side ( $\mu$ inch)
10	0.05/0.05	.06/0.06
20	0.10/0.10	12/12
25	0.125/0.125	15/15
35	0.175/0.175	22/22
50	0.25/0.25	30/30
75	0.375/0.375	45/45
100	0.50/0.50	60/60
D 50/25	0.25/0.125	30/15
D 75/25	0.375/0.125	45/15
D 100/25	0.50/0.125	60/15
D 100/50	0.50/0.25	60/30
D 135/25	0.675/0.125	82/15

Table 12 below shows corrosion rate data for AISI type 1010 steel, a mild steel very similar in composition to the Type L steel used in tin plated food cans.

**Table 12**  
**Corrosion Rates for AISI 1010 Steel (from Reinhart<sup>3</sup>)**

Depth	O <sub>2</sub> Concentration (ml/l)	pH	Salinity (ppt)	Temperature °C	Exposure time (days)	Rate Exposed (mpy)	Rate Buried (mpy)
5	5.9	8.1	33.51	15	181	9.1	-
5	5.9	8.1	33.51	15	366	8.0	-
5	5.5	8.1	33.31	15	588	8.9	-
2340	0.4	7.5	34.36	5	197	1.6	1.2
2370	0.4	7.5	34.36	5	402	1.1	1.1
5640	1.3	7.6	34.51	2.3	123	2.7	1.9
5640	1.3	7.6	34.51	2.3	751	0.8	0.6
5300	1.2	7.5	34.51	2.6	1064	1.0	0.7
6780	1.6	7.7	34.40	2.2	403	1.9	1.1

Table 13 shows corrosion rate data for JIS SS41steel (equivalent to AISI 1020), a mild steel with around 0.2% carbon.

**Table 13**  
**Corrosion rate data for JIS SS41 steel (from Shimada<sup>5</sup>)**

Depth (ft)	O <sub>2</sub> (ml/l)	pH	Exposure time (days)	Rate (mpy) Exposed
6.6	6.9	8.1	720	11
98	6.4		720	9
197	3.3		720	4
295	6.2		720	8




Figure 19. Corrosion rate of mild steel vs. oxygen concentration.

The data from the Reinhart and Shimada studies show remarkably good correlation between oxygen concentration and corrosion rate, as is shown in figure 19. However the Shimada study also examined corrosion rates in a polluted bay, where the dissolved oxygen concentration was only 0.09 ml/l. The corrosion rate under those conditions was about 7.9 mpy, much larger than would be expected from the oxygen level. This may be due to the presence of sulfate reducing bacteria and the sulfide that they produce. The presence of sulfide generally increases the corrosion rate of steel in seawater. This may be due to catalytic effects that increase the rates of both the oxidation and reduction reactions at the steel surface, as well as stabilization of the corrosion products as insoluble sulfides.<sup>12</sup>

Peterson and Lennox<sup>5</sup> measured the corrosion rates of mild steel where the mean temperature was 25 °C, the mean dissolved oxygen content was 8.6 ml/l and the mean pH was 8.07. Samples were exposed in a laboratory cell at a low flow rate, from a pier with normal tidal flow, and in a flume at flow of 0.23 meters/s. The corrosion rates were 2.3, 4.2, and 7.9 mpy respectively. While these data show the effect of

flow, the corrosion rates are somewhat less than might be expected from the oxygen concentration when compared with the data in figure 19. This is indicative of the variability in results that might be expected between different studies that use perhaps slightly different materials and methodologies at different locations. Other near-surface immersion studies<sup>4</sup> (exposure conditions unknown) give initial corrosion rates for carbon steel of about 4-5 mpy, which stabilize over the long term (many years) to around 3 mpy.

Controlled laboratory studies have shown that the corrosion of steel should be accelerated by increasing temperature.<sup>5</sup> However, in natural waters, the temperature has a significant effect on other factors such as the dissolved oxygen concentration and level of biological activity. For example, Shimada et. al.<sup>5</sup> found that in unpolluted seawater the corrosion rates were highest in winter months when oxygen concentrations were highest even though temperatures were lower. In polluted seawater (with very low oxygen levels) the corrosion rates were highest in summer months when biological activity was at its highest. Reinhart and Jenkins<sup>13</sup> used linear multivariate regression analysis to give the following formula for the dependence of the corrosion rate of steel on oxygen concentration and temperature:

$$\text{Corrosion Rate}(\mu\text{m/yr}) = 21.3 + 25.4 [\text{O}_2 \text{ (ml/l)}] + 0.356 [\text{T (}^\circ\text{C)}] \quad (10)$$

At ocean oxygen concentrations that typically range from 1-7 ml/l, it is clear from equation 10 that oxygen concentration dominates the corrosion rate for steel in seawater.

The effect of biofouling on the corrosion rates of steel is open to some question. In theory, marine organism growth should slow uniform attack corrosion by restricting access of oxygen to the steel surface. Pitting and crevice corrosion should be increased by the creation of differential aeration cells at marine organism attachment sites. Comparison of corrosion rates between samples immersed in filtered and unfiltered sea water show no significant difference, either in the general corrosion rate or the depth of pits.<sup>14</sup> Other studies<sup>15</sup> indicate that under fouling conditions initial corrosion rates can be quite large (>13 mpy) until specimens are covered by marine organisms (after about 1.5 years). At that point corrosion rates decrease until oxygen is excluded from the specimen surface and corrosion is controlled by sulfate reducing bacteria. The corrosion rates then stabilize at about 2-3 mpy.

## **Use of Corrosion rates to predict lifetimes and mass loading**

The simplest use of the corrosion rate information is in the prediction of the lifetime of a particular container in seawater. A complicating factor is the presence of the organic coatings on the container surfaces. These coatings will tend to protect the metal from corrosion, but not indefinitely. The shredding process provides ample edges and defects for the origination of crevice corrosion. The crevice corrosion will eventually destroy the usefulness of the coating. The inside of the tin plated steel cans are generally much better protected with tin plating and coatings than the outside. Similarly, the aluminum can bodies have a protective coating on the inside, and only decorative paint on the outside. The outsides of the cans will begin to corrode first. The can may completely corrode from the outside in before the interior surfaces are significantly attacked. Experience from the exposure of some shipboard waste in San Diego bay for ten months seems to confirm this hypothesis. The coated interiors of tin plated steel cans remained in relatively good condition after ten months exposure (the tin plating was still intact under the organic coating) while the exteriors of the cans showed severe corrosion. Consequently, container lifetimes can be estimated by simply dividing the container wall thicknesses by the corrosion rates. For a steel can that is nominally 0.010 inch thick, the lifetime at a corrosion rate of 4 mpy would be 2.5 years. We measured aluminum can wall thicknesses that varied from as little as 0.004 inch near the middle to 0.013 inch on the bottom. For an aluminum can body to completely corrode at 2 mpy would take 6.5 years. The aluminum

can lids would be expected to corrode more slowly, as they are coated on both sides and are made of the more corrosion resistant 5000 series alloy.

Additional information can be gained by calculating what we call the mass removal rate fraction,  $m_f$ . We define this as fraction of mass removed by corrosion per year for a given container thickness, or

$$m_f = r\rho \left( \frac{\text{effective container surface area}}{\text{container mass}} \right) \quad (11)$$

where  $r$  is the corrosion rate in inches per year and  $\rho$  is the density of the alloy. A "base box" of tin plated steel has a total area of 31360 in<sup>2</sup>. If we assume on the average 100 lbs per base box, and a corrosion rate of 4 mpy,  $m_f$  for tin plated steel cans would be 0.355 pounds of corroded metal per pound of cans per year. An aluminum can weighs about 0.48 ounce<sup>16</sup> and has a surface area of about 25 in<sup>2</sup> of 3004 aluminum alloy and about 3.6 in<sup>2</sup> of 5182 aluminum alloy. Because the nature of the corrosion of aluminum makes it difficult to predict corrosion rates as a function of ocean conditions, and because the rates do not vary so greatly between the two can alloys it seems prudent to just make an order of magnitude estimate independent of alloy or conditions. If one takes 2 mpy as the corrosion rate (this should probably be considered an upper bound), then  $m_f$  for aluminum can waste would be about 0.18 pounds corroded per pound of waste per year.

In a situation where waste is thrown overboard at a constant rate,  $W$ , the amount of material remaining uncorroded on the ocean floor,  $M$ , can be modeled by the differential equation:

$$\frac{dM}{dt} = W - m_f M \quad (12)$$

where  $t$  is the time. This simply says that the rate of waste accumulation on the ocean floor equals the difference between the rate of addition and the rate of loss due to corrosion. Under steady state conditions, where  $dM/dt$  is zero, we have

$$M = \frac{W}{m_f} \quad (13)$$

and by definition of steady state, the amount of material being lost as corrosion equals the amount of material that is being added by being thrown overboard. Using the  $m_f$  calculated above for tin plated steel cans, if 100 pounds per year of waste is thrown overboard, at steady state 282 pounds of waste would be always present on the ocean floor. A complete solution of equation 12 leads to:

$$M = \frac{W}{m_f} (1 - e^{-m_f t}) \quad (14)$$

This indicates that steady state is only achieved at infinite time. Steady state is approached with time constant  $1/m_f$ . Given the  $m_f$  calculated above for tin plated steel, 50% of steady state is reached in 2.0 years, 90% in 6.5 years, and 99% in 13 years.

## References

1. Fontana, M. G., *Corrosion Engineering*, McGraw Hill, New York,(1986).
2. ASTM Standard Practice G1-81, ASTM, Philadelphia (1981).
3. Reinhart, F. M. "Corrosion of Metals and Alloys in the Deep Ocean", Technical Report R 834, Civil Engineering Laboratory, Naval Construction Battalion Center, Port Hueneme, California 93043, February 1976.
4. Beavers, J. A. , Koch, G. H., and Berry, W. E., *Corrosion of Metals in Marine Environments*, MCIC Report 86-50, Battelle Columbus Division, Columbus, Ohio 43201-2693, July 1986.
5. Shimada, H. Miida, K., and Yokooji, T, "The Relation Between the Corrosion Behavior of Steels and the Characteristics of Various Ocean Environments", *International Ocean Development Conference, vol 2*, 165-83 (1975).
6. Peterson, M. H. and Lennox T. J. Jr., " Effects Of Exposure Conditions On The Corrosion Of Mild Steel, Copper, And Zinc In Seawater", *Materials Performance*, 23, 15-18, (1984).
7. Ailor, W. H. Jr., "Aluminum Alloys After Five Years in Seawater", *Materials Performance and the Deep Sea, ASTM STP 445*, 115-130 ,American Society for Testing and Materials, )1969).
8. Dexter, S. C. ,"Effect of Variation in Sea Water Upon the Corrosion of Aluminum", *Corrosion*, 36, 423-432, (1980).
9. Brick, R. M. ,Daly, J. J., Koehler, E. L. and Skibbe, A. G., "The Selection of Tin Coatings for Steel Containers", *Metals Handbook, vol 1*, pp1133-1141, Taylor Lyman Editor, Metals Park, Ohio, 1961.
10. Horne, R. A., *Marine Chemistry*, John Wiley and Sons, New York, 1969.
11. ASTM Standard A 624 (1984).
12. West, J. M. *Basic Corrosion and Oxidation*, p122, John Wiley and Sons, New York, 1980.
13. Reinhart, F. M. and Jenkins, J. F., "The Relationship Between the concentratio of Oxygen in Seawater and the Corrosion of Metals", U. S. Naval Civil Eng. Lab. Report, pp. 562-577.
14. Traverso, E. et al., "Effects of Rolling Direction and Tensile Stress on the Corrosion of Naval Carbon Steel in Natural Marine Environments-I: Natural Seawater", *British Corrosion Journal*, 14 (2), pp97-102 (1979).
15. Shumacher, M. *Seawater Corrosion HandBook*, p. 383,Noyes Data Corporation, Park Ridge, New Jersey (1979).
16. Hosford, W. F., and Duncan, J. L. ,"The Aluminum Beverage Can", *Scientific American*,271(3), 49,(1994).

## **APPENDIX H**

### **WAKE DISPERSION MODELING RESULTS**

Source: Wake Dispersion Modeling.  
San Diego, California  
Naval Command, Control & Ocean Surveillance  
Center, RDTE Division, Code 522,  
Naval Coastal Systems Station, 1995

**DIANA::HYMAN**


**JOB 224**

**CVN20.LAS;1**


**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN20.LAS;1  
**Last Modified:** 25-MAY-1995 12:52  
**Owner UIC:** [HYMAN]

**Length:** 1707 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LASER\_B1102C  
**Submitted:** 25-MAY-1995 12:52  
**Printer queue:** LASER\_B1102C  
**Printer device:** LPS17A


Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified X = 0.56 km = 1.69 L




CONTOUR FROM 0.00000E+00 TO 1.00000E+01 CONTOUR INTERVAL OF 0.20000E+01



CONTOUR FROM 0.00000E+00 TO 0.30000E+01 CONTOUR INTERVAL OF 0.05000E+01




CONTOUR FROM 0.00000E+00 TO 1.00000E+01 CONTOUR INTERVAL OF 0.20000E+01




CONTOUR FROM 0.00000E+00 TO 0.30000E+01 CONTOUR INTERVAL OF 0.05000E+01


Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified X = 1.06 km = 3.19 L




CONTOUR FROM 0.0000E+00 TO 0.0000E+00 DEGREE  
 CENTER INTERVAL OF 0.1000E+02 PT(3,3)= 066600E-02



CONTOUR FROM 0.0000E+00 TO 0.2000E-01 CENTUR INTERVAL OF 0.0300E-02 PT(3,3)= 0.2666E-02 LABELS SCALED BY 10000




CONTOUR FROM 0.0000E+00 TO 0.0000E+00 DEGREE  
 CENTER INTERVAL OF 0.1000E+02 PT(3,3)= 066600E-02




CONTOUR FROM 0.0000E+00 TO 0.2000E-01 CENTUR INTERVAL OF 0.0300E-02 PT(3,3)= 0.2666E-02 LABELS SCALED BY 10000


Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified  
 $X = 2.06 \text{ km} = 6.20 \text{ L}$




CONT. FROM 0.0000E+00 TO 0.3449E3    CONTOUR INTERVAL OF 0.8095E99    PT(3,3) = 0.5E9E7E-02



CONT. FROM 0.0000E+00 TO 0.3449E3    CONTOUR INTERVAL OF 0.8095E99    PT(3,3) = 0.5E9E7E-02





CONT. FROM 0.0000E+00 TO 0.7388E-01    CONTOUR INTERVAL OF 0.2988E-02    PT(3,3) = 0.8499E-03    LABELS SCALED BY 10000.





CONT. FROM 0.0000E+00 TO 0.7388E-01    CONTOUR INTERVAL OF 0.2988E-02    PT(3,3) = 0.8499E-03    LABELS SCALED BY 10000.



Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified X = 3.06 km = 9.22 L





CONTOUR FROM 0.0000E+00 TO 0.028339E+01 CENTER INTERVAL OF 0.043338E-01 PT(13,31) = 0.0424252E-02



CONTOUR FROM 0.0000E+00 TO 0.000002E-01 CENTER INTERVAL OF 0.000002E-02 PT(13,31) = 0.000000E-03 LABELS SCALED BY 10000.



CONTOUR FROM 0.00000E+00 TO 0.22899E+03 CONTOUR INTERVAL OF 0.65708E+01 P1(3,31)= 0.22899E+02



CONTOUR FROM 0.88888E+08 TO 0.10555E+01 CONTOUR INTERVAL OF 0.81888E+02 P1(3,31)= 0.80888E+03 LABELS SCALED BY 10000.

**DIANA::HYMAN**

**JOB 1104**


**FFG10.LAS;1**

**File:** \_\$40SDUA29:[HYMAN.GRID.DISPERSION]FFG10.LAS;1  
**Last Modified:** 26-MAY-1995 10:26  
**Owner UIC:** [HYMAN]


**Length:** 1209 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LASER\_B1102C  
**Submitted:** 26-MAY-1995 10:26  
**Printer queue:** LASER\_B1102C  
**Printer device:** LPS17A

Unstratified



X = 0.56 km = 4.26 L



CONTOUR FROM 0.0000E+00 TO 0573539 CONTOUR INTERVAL OF 0114788 PT(3,31)= 0573539E-02




CONTOUR FROM 0.58000E+00 TO 0.82288E-01 CONTOUR INTERVAL OF 0.46899E-02 PT(3,31)= 0.22869E-02 LABELS SCALED BY 10000.




Frigate - 10 kts (5.15 m/s)


Unstratified X = 1.06 km = 8.07 L




CONTOUR FROM 0.0000E+00 TO 0.9999E+00 CONTOUR INTERVAL OF 0.9999E+00 PT(3,3) = 0.9999E+00



CONTOUR FROM 0.0000E+00 TO 0.9999E+00 CONTOUR INTERVAL OF 0.9999E+00 PT(3,3) = 0.9999E+00






CONTOUR FROM 0.0000E+00 TO 0.9999E+00 CONTOUR INTERVAL OF 0.9999E+00 PT(3,3) = 0.9999E+00




CONTOUR FROM 0.0000E+00 TO 0.9999E+00 CONTOUR INTERVAL OF 0.9999E+00 PT(3,3) = 0.9999E+00


Unstratified X = 1.56 km = 11.87 L



Frigate - 10 kts (5.15 m/s)  
 Unstratified  
 X = 2.06 km = 15.68 L







CONTOUR FROM 0.0000E+00 TO 0.22E29 CONTOUR INTERVAL OF 0.22E29  
 10 MICRON PART. POPULATION AT 2.06 KM  
 0239396E-02







CONTOUR FROM 0.6680E+08 TO 0.2684E-02 CONTOUR INTERVAL OF 0.2684E-02  
 3000 MICRON PART. POPULATION AT 2.06 KM  
 0.66821E-03 LABELS SCALED BY 10000.

Frigate - 10 kts (5.15 m/s)  
 Unstratified  
 $X = 0.66 \text{ km} = 5.01 \text{ L}$




Frigate - 10 kts (5.15 m/s)  
 Unstratified X = 1.32 km = 10.01 L




Unstratified


X = 2.63 km = 20.01 L




CONTOUR FROM 10.101 TO 10.593 CONTOUR INTERVAL OF 0.24616E-01 P113.31= 10.293



MAXIMUM VECTOR







CONTOUR FROM 0.10000E-03 TO 0.15594E-01 CONTOUR INTERVAL OF 0.76222E-03 P113.31= 0.24191E-03 LABELS SCALED BY 10000



CONTOUR FROM -0.68174E-06 TO 0.39922E-01 CONTOUR INTERVAL OF 0.16961E-02 P113.31= 0.00000E+00 LABELS SCALED BY 10000.

Frigate - 10 kts (5.15 m/s)  
 Unstratified X = 3.94 km = 30.01 L





**DIANA::HYMAN**



**JOB 577**

**CVN20-STRAT.LAS;2**



**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN20-STRAT.LAS;2  
**Last Modified:** 7-JUN-1995 10:20  
**Owner UIC:** [HYMAN]

**Length:** 2049 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LASER\_B1102C  
**Submitted:** 7-JUN-1995 10:20  
**Printer queue:** LASER\_B1102C  
**Printer device:** LPS17A

Stratified  
X = 0.56 km = 1.69 L







CONTOUR FROM 0.0000E+00 TO 281868 CONTOUR INTERVAL OF 056258 FIG. 31- 02Z019E-01



CONTOUR FROM 000000E+00 TO 800000E+00 CONTOUR INTERVAL OF 100000 FIG. 300000E+00 300118RBL 91508E028Y 10000.



Aircraft Carrier - 20 kts (10.3 m/s)  
 Stratified X = 1.06 km = 3.19 L





CONTOUR FROM 0.00000E+00 TO 1.47588 CONTOUR INTERVAL OF 0.283515 P1(3,3)= 0114188E-01

CONTOUR FROM 0.00000E+00 TO 1.47588 CONTOUR INTERVAL OF 0.283515 P1(3,3)= 0114188E-01

Stratified X = 2.06 km = 6.20 L







CONTOUR FROM 0.00000E+00 TO 0.785587 CONTOUR INTERVAL OF 0.152369 PT(3,3)= 0.85688E-02







CONTOUR FROM 0.00000E+00 TO 0.46688E-01 CONTOUR INTERVAL OF 0.56688E-02 PT(3,3)= 0.18688E-02 LABELS SCALED BY 10000.

Aircraft Carrier - 20 kts (10.3 m/s)  
 Stratified  
 $X = 3.06 \text{ km} = 9.22 L$







CONTOUR FROM 0.00000E+00 TO 0.00000E+01 CONTOUR INTERVAL OF 0.00000E-02 PLOT(S) IS 0.00000E-03 LABELS SCALED BY 10000.


Aircraft Carrier - 20 kts (10.3 m/s)  
 Stratified X = 5.06 km = 15.24 L



Aircraft Carrier - 20 kts (10.3 m/s)  
 Stratified X = 0.55 km = 1.67 L




Aircraft Carrier - 20 kts (10.3 m/s)  
Stratified X = 1.11 km = 3.34 L




CONTOUR FROM 10 080 TO 10 753 CONTOUR INTERVAL OF 0.33685E-01 P113.31= 10 293

KINETIC ENERGY AT 3.34 L




CONTOUR FROM 0.10000E-03 TO 0.88145E-01 CONTOUR INTERVAL OF 0.36539E-02 P113.31= 0.97852E-03 LABELS SCORED BY 1000.







0 103E-00  
MAXIMUM VECTOR

SCALAR FIELD AT 3.34 L







CONTOUR FROM -0.82785E-03 TO 0.12786 CONTOUR INTERVAL OF 0.61345E-02 P113.31= -0.15878E-35 LABELS SCORED BY 1000.0





Aircraft Carrier - 20 kts (10.3 m/s)  
 Stratified X = 2.21 km = 6.67 L



Aircraft Carrier - 20 kts (10.3 m/s)  
 Stratified  
 X = 3.32 km = 10 L



Aircraft Carrier - 20 kts (10.3 m/s)  
 Stratified  
 X = 5.53 km = 16.67 L



**DIANA::HYMAN**

**JOB 1792**


**FFG10-STRAT.LAS;2**

**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]FFG10-STRAT.LAS;2  
**Last Modified:** 12-JUN-1995 13:48  
**Owner UIC:** [HYMAN]

**Length:** 5091 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 12-JUN-1995 13:48  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER


Stratified  
 X = 0.66 km = 5.01 L

velocity - 1000 m/s




CONTOR FROM 4.9581 TO 5.4259 CONTOUR INTERVAL OF 0.22850E-01 P(13,31)= 5.1450

AXIAL VELOCITY AT 5.01 L




0.150E+00  
 MIXING VECTOR



CONTOR FROM 0.10000E-03 TO 0.22825E-01 CONTOUR INTERVAL OF 0.11362E-02 P(13,31)= 0.30376E-03 LABELS SCHEED BY 10000.





KINETIC ENERGY AT 5.01 L



CONTOR FROM -0.27425E-02 TO 0.11914 CONTOUR INTERVAL OF 0.60941E-02 P(13,31)= 0.00000E+00 LABELS SCHEED BY 1000.0

SCALAR FIELD AT 5.01 L


Frigate - 10 kts (5.15 m/s)  
 Stratified  
 X = 1.32 km = 10.01 L




Frigate - 10 kts (5.15 m/s)  
 Stratified  
 X = 1.97 km = 15.01 L




CONTOUR FROM 5.0036 TO 5.3109 CONTOUR INTERVAL OF 0.11361E-01 PT(3,3)= 5.1450  
 AXIAL VELOCITY AT 15.01 L



0.183E+00  
 MAXIMUM VECTOR




CONTOUR FROM 0.15537E-04 TO 0.22357E-02 CONTOUR INTERVAL OF 0.11101E-03 PT(3,3)= 0.67451E-03 LABELS SCALED BY 0.10000E+06  
 KINETIC ENERGY AT 15.01 L




CONTOUR FROM -0.20154E-02 TO 0.79910E-01 CONTOUR INTERVAL OF 0.41363E-02 PT(3,3)= -0.15108E-35 LABELS SCALED BY 10000.  
 SCALAR FIELD AT 15.01 L


Frigate - 10 kts (5.15 m/s)  
 Stratified  
 X = 2.63 km = 20.01 L




CONTOUR FROM 5 0910 TO 5.2789 CONTOUR INTERVAL OF 0.9368E-02 PT(3,31)= 5.1450  
 AXIAL VELOCITY AT 20.01 L



CONTOUR FROM 0.80722E-05 TO 0.12195E-02 CONTOUR INTERVAL OF 0.63746E-04 PT(3,31)= 0.35475E-03 LABELS SCALED BY 0.10000E+08  
 KINETIC ENERGY AT 20.01 L



0.252E+00  
 VELOCITY VECTOR




CONTOUR FROM -0.40247E-02 TO 0.66691E-01 CONTOUR INTERVAL OF 0.36759E-02 PT(3,31)= 0.13562E-31 LABELS SCALED BY 10000.  
 SCALAR FIELD AT 20.01 L


Stratified - 0.001 K/s (0.000 100/s)

Stratified


X = 3.94 km = 30.01 L




CONTINUUM FROM 5.0651 TO 5.1450 CONTINUUM INTERVAL OF 0.024575-02 P113.31= 5.1450



0.422E+00  
MAXIMUM VECTOR



CONTINUUM FROM 0.6031E-01 TO 0.94950E-05 CONTINUUM INTERVAL OF 0.6031E-01 P113.31= 0.94950E-05



CONTINUUM FROM -0.19528E-20 TO 0.28195E-02 CONTINUUM INTERVAL OF 0.28195E-02 P113.31= -0.19528E-20 LABELS SCALED BY 10000.

**DIANA::HYMAN**





**JOB 430**

**CVN20.LAS;1**





**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN20.LAS;1  
**Last Modified:** 7-JUN-1995 08:25  
**Owner UIC:** [HYMAN]

**Length:** 6542 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 7-JUN-1995 08:25  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER





Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified  
 $X = 0.55 \text{ km} = 1.67 \text{ L}$







Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified  
 $X = 1.11 \text{ km} = 3.34 \text{ L}$







Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified X = 2.21 km = 6.67 L



Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified X = 3.32 km = 10 L



Aircraft Carrier - 20 kts (10.3 m/s)  
 Unstratified  
 $X = 5.53 \text{ km} = 16.67 \text{ L}$



**DIANA::HYMAN**





**JOB 1672**

**FFG-20.LAS;1**

**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]FFG-20.LAS;1  
**Last Modified:** 1-JUN-1995 15:29  
**Owner UIC:** [HYMAN]

**Length:** 1604 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 1-JUN-1995 15:29  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER

# Unstratified X = 0.56 km = 4.26 L





CONTOUR FREQ 0.0000E+00 18 097588 CONTOUR INTERVAL 0.07477E-01 1713.31= 0.9938E-02

CONTOUR FREQ 0.0000E+00 18 0.9886E-02 CONTOUR INTERVAL 0.3660E-02 1713.31= 0.6888E-08 LABELS SCALED BY 010000E+06



Frigate - 20 kts (10.3 m/s)

Unstratified

X = 1.06 km = 8.07 L







CONTR FREQ 0.0000E-00 18 0395877 CONTOUR INTERVAL OF 075888E-01 P113.31= 0.88388E-02







CONTR FREQ 0.0000E-00 18 0.37808E-02 CONTOUR INTERVAL OF 0.28888E-02 P113.31= 0.88388E-02 LABELS SCALED BY 0.10000E+06

Trigate - 20 kts (10.3 m/s)  
 Unstratified  
 X = 2.06 km = 15.68 L







CONTOUR FROM 0.0000E+00 TO 0.2502E-02 CONTOUR INTERVAL OF 0.2502E-02  
 CONTOUR FROM 0.0000E+00 TO 0.5808E-08 CONTOUR INTERVAL OF 0.5808E-08

Frigate - 20 kts (10.3 m/s)  
 Unstratified  
 $X = 3.06 \text{ km} = 23.29 \text{ L}$



Frigate - 20 kts (10.3 m/s)

Unstratified X = 5.06 km = 38.52 L




CONTOUR FROM 0.00000E+00 TO 0.00000E+00 BY 1.0000E+00


CONTOUR FROM 0.00000E+00 TO 0.00000E+00 BY 0.10000E+06




Unstratified X = 1.32 km = 10.01 L




CONTour FROM 10.048 TO 10.794 CONTOUR INTERVAL OF 0.37298E-01 PT(13,31)= 10.293  
AXIAL VELOCITY AT 10.01 L



0.605E-01  
MAXIMUM VECTOR




CONTour FROM 0.10000E-03 TO 0.41008E-01 CONTOUR INTERVAL OF 0.20454E-02 PT(13,31)= 0.41363E-03 LABELS SCALED BY 10000.  
KINETIC ENERGY AT 10.01 L





CONTour FROM -0.40261E-06 TO 0.49650E-01 CONTOUR INTERVAL OF 0.24825E-02 PT(13,31)= 0.00000E+00 LABELS SCALED BY 10000.  
SCALAR FIELD AT 10.01 L


Frigate - 20 kts (10.3 m/s)  
 Unstratified  
 $X = 2.63 \text{ km} = 20.01 \text{ L}$



CONTour FROM 0.10000E-03 TO 0.15344E-01 CONTOUR INTERVAL OF 0.76222E-03 P(13,31)= 0.24131E-03 LABELS SCALED BY 10000.




CONTour FROM -0.69174E-06 TO 0.35922E-01 CONTOUR INTERVAL OF 0.16991E-02 P(13,31)= 0.00000E-00 LABELS SCALED BY 10000.





MAXIMUM VECTOR


Frigate - 20 kts (10.3 m/s)  
 Unstratified  
 $X = 3.94 \text{ km} = 30.01 \text{ L}$







CONTOUR FROM 0.10000E-03 TO 0.88552E-02 CONTOUR INTERVAL OF 0.45796E-03 P113.31= 0.16397E-03 LABELS SCALED BY 0.10000E+06  
 KINETIC ENERGY AT 30.01 L



CONTOUR FROM -0.27591E-06 TO 0.24167E-01 CONTOUR INTERVAL OF 0.12084E-02 P113.31= 0.00000E-00 LABELS SCALED BY 10000.  
 SCALAR FIELD AT 30.01 L



Frigate - 20 kts (10.3 m/s)  
 Unstratified X = 6.57 km = 50.01 L



**DIANA::HYMAN**

**JOB 668**



**FFG20-STRAT.LAS;1**

**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]FFG20-STRAT.LAS;1  
**Last Modified:** 13-JUN-1995 08:08  
**Owner UIC:** [HYMAN]



**Length:** 1793 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 13-JUN-1995 08:08  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER

Frigate - 20 kts (10.3 m/s)

Stratified X = 3.06 km = 23.29 L





CONTOUR FROM 0.00000E+00 TO 0.14093E-01 PARTICLE INTERVAL OF 0.26548E-01 PARTICLES AT 3.06 KM





CONTOUR FROM 0.00000E+00 TO 0.12490E-02 PARTICLE INTERVAL OF 0.26548E-03 PARTICLES AT 3.06 KM



Stratified  
X = 0.56 km = 4.26 L







CONTOUR FROM 0.00000E+00 TO 0.20000E+02 CONTOUR INTERVAL OF 0.57250E+01 PLOT 3.31= 023000E-02



CONTOUR FROM 0.00000E+00 TO 0.30000E+02 CONTOUR INTERVAL OF 0.30000E+01 PLOT 3.31= 0.60000E-03 LURELS SCALED BY 010000E+00

Frigate - 20 kts (10.3 m/s)





Stratified X = 1.06 km = 8.07 L



Frigate - 20 kts (10.3 m/s)

Stratified

X = 2.06 km = 15.68 L



**DIANA::HYMAN**

**JOB 674**





**FFG20-STRAT.LAS;2**

**File:** \_\$40SDUA29:[HYMAN.GRID.DISPERSION]FFG20-STRAT.LAS;2  
**Last Modified:** 13-JUN-1995 08:11  
**Owner UIC:** [HYMAN]

**Length:** 6156 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 13-JUN-1995 08:11  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER


Stratified

X = 1.32 km = 10.01 L




Frigate - 20 kts (10.3 m/s)

Stratified X = 0.66 km = 5.01 L




CONTOUR FROM 9.8010 TO 11.146 CONTOUR INTERVAL OF 0.62308E-01 PT(3,31)= 10.293

AXIAL VELOCITY AT 5.01 L




0.167E+00  
MAXIMUM VECTOR



CONTOUR FROM 0.10000E-03 TO 0.11784 CONTOUR INTERVAL OF 0.59770E-02 PT(3,31)= 0.62455E-03 LABELS SCALED BY 1000.0


KINETIC ENERGY AT 5.01 L




CONTOUR FROM -0.40343E-03 TO 0.63990E-01 CONTOUR INTERVAL OF 0.32197E-02 PT(3,31)= 0.00000E+00 LABELS SCALED BY 10000.0

SCALAR FIELD AT 5.01 L


Frigate - 200 Kts (100.0 m/s)  
 Stratified  
 X = 2.63 km = 20.01 L




CONTOUR FROM 10.190 TO 10.607 CONTOUR INTERVAL OF 0.20866E-01 P1(3,31)= 10.293  
 AXIAL VELOCITY AT 20.01 L



0.135E+00  
 MAXIMUM VECTOR







CONTOUR FROM 0.17148E-04 TO 0.14037E-01 CONTOUR INTERVAL OF 0.70100E-03 P1(3,31)= 0.20866E-03 LABELS SCALED BY 10000.  
 KINETIC ENERGY AT 20.01 L







CONTOUR FROM -0.20009E-03 TO 0.31089E-01 CONTOUR INTERVAL OF 0.19819E-02 P1(3,31)= 0.00000E+00 LABELS SCALED BY 10000.  
 SCALAR FIELD AT 20.01 L

Frigate - 20 kts (10.3 m/s)  
 Stratified  
 $X = 3.94 \text{ km} = 30.01 \text{ L}$



Frigate - 20 kts (10.3 m/s)  
 Stratified  
 X = 6.57 km = 50.01 L



**DIANA::HYMAN**


**JOB 232**

**CVN10.LAS;1**



**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN10.LAS;1  
**Last Modified:** 25-MAY-1995 13:02  
**Owner UIC:** [HYMAN]

**Length:** 1316 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LASER\_B1102C  
**Submitted:** 25-MAY-1995 13:02  
**Printer queue:** LASER\_B1102C  
**Printer device:** LPS17A


Aircraft Carrier - ... m/s)  
 Unstratified X = 0.56 km = 1.69 L







CONTOUR FROM 0.00000E+00 TO 241377 CONTOUR INTERVAL OF 0188755 PT(3,31)= 0281577E-01



CONTOUR FROM 0.80000E+00 TO 0.42000E+01 CONTOUR INTERVAL OF 0.85555E-01 PT(3,31)= 0.99898E-02 LABELS SCRLED BY 100000




Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified X = 1.06 km = 3.19 L




CONTOUR FROM 0.0000E+00 TO 136639 CONTOUR INTERVAL OF 0239559 PT(3,3)= 013789E-01

CONTOUR FROM 0.86098E+00 TO 0.49999E+01 CONTOUR INTERVAL OF 0.86998E-02 PT(3,3)= 0.86998E-02 LABELS SCALED BY 100000


Aircraft Carrier - 10 kts (5.3 m/s)  
 Unstratified  
 X = 1.56 km = 4.70 L




CONTOUR FROM 0.0000E+00 TO 781538 CONTOUR INTERVAL OF 0185035 P1(3,3)= 0912679E-01



CONTOUR FROM 0.0000E+00 TO 781538 CONTOUR INTERVAL OF 0185035 P1(3,3)= 0912679E-01







CONTOUR FROM 0.0000E+00 TO 0.35185E-01 CONTOUR INTERVAL OF 0.28888E-02 P1(3,3)= 0.28488E-03 LABELS SCALED BY 100000



CONTOUR FROM 0.0000E+00 TO 0.35185E-01 CONTOUR INTERVAL OF 0.28888E-02 P1(3,3)= 0.28488E-03 LABELS SCALED BY 100000

Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified X = 2.06 km = 6.20 L



CONTOUR FROM 0.00000E+00 TO 0.34853E-01 CONTOUR INTERVAL OF 0.00000E+00 PT(3,31)= 0.790169E-02

CONTOUR FROM 0.00000E+00 TO 0.78658 CONTOUR INTERVAL OF 0.18263 PT(3,31)= 0.790169E-02

**DIANA::HYMAN**

**JOB 415**

**CVN10.LAS;1**

*Aircraft Carrier  
10 2's (5 5 m s)  
Justified*





**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN10.LAS;1  
**Last Modified:** 7-JUN-1995 08:10  
**Owner UIC:** [HYMAN]

**Length:** 7709 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 7-JUN-1995 08:10  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER





$$L = 332 \text{ m}$$

$$X = 1.67 \times 332 = 554 \text{ m}$$


Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified X = 0.55 km = 1.67 L




Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified X = 1.11 km = 3.34 L




Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified  
 $X = 1.66 \text{ km} = 5.00 \text{ L}$




CONTour FROM 0.10000E-03 TO 0.13465E-01 CONTOUR INTERVAL OF 0.70345E-03 PT(3,31)= 0.26665E-03 U-RBELS SCALED BY 10000.



CONTour FROM 0.10000E-03 TO 0.13465E-01 CONTOUR INTERVAL OF 0.70345E-03 PT(3,31)= 0.26665E-03 U-RBELS SCALED BY 10000.







0.274E-01  
 MAXIMUM VECTOR







CONTour FROM -0.16883E-05 TO 0.21782 CONTOUR INTERVAL OF 0.16881E-01 PT(3,31)= 0.17624E-35 U-RBELS SCALED BY 10000.0





Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified  
 $X = 2.21 \text{ km} = 6.67 \text{ L}$



Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified  
 $X = 2.77 \text{ km} = 8.34 \text{ L}$



Aircraft Carrier - 10 kts (5.15 m/s)  
 Unstratified  
 X = 3.32 km = 10 L



**DIANA::HYMAN**

**JOB 412**





**CVN10-STRAT.LAS;2**

**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN10-STRAT.LAS;2  
**Last Modified:** 7-JUN-1995 08:07  
**Owner UIC:** [HYMAN]

**Length:** 8383 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 7-JUN-1995 08:07  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER







Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified  
 $X = 1.06 \text{ km} = 3.19 \text{ L}$



CONTOUR FROM 0.00000E+00 TO 1.00000E+03 INCREMENTS 0.00000E+00  
 CONTOUR INTERVAL OF 0.37753 FT(3.91) = 0.11789E-01





Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified  
 X = 1.56 km = 4.70 L



CONTOUR FROM 0.0000E+00 TO 1.2028E+01 CENTER INTERVAL OF 0.2000E+01

CONTOUR FROM 0.0000E+00 TO 0.0006E+01 CENTER INTERVAL OF 0.0001E+01





Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified X = 2.06 km = 6.20 L







CONTOUR FROM 0.0000E+00 TO 0.1727E+01 CONTOUR INTERVAL OF 0.267900 PLS:31= 0190799E-02

CONTOUR FROM 0.0000E+00 TO 0.9850E+03 CONTOUR INTERVAL OF 0.9850E+03 PLS:31= 0000E+00





Aircraft Carrier - ... m/s  
 Stratified X = 2.56 km = 7.71 L







Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified  
 $X = 0.55 \text{ km} = 1.67 \text{ L}$




Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified  
 X = 1.11 km = 3.34 L




Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified  
 X = 1.66 km = 5.00 L




Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified  
 X = 2.21 km = 6.67 L



CONTour FRSH 5.1221 10 5.2735 CONTOUR INTERVAL OF 0.7561E-02 PT(3,31)= 5.1450  
 AXIAL VELOCITY AT 6.67 L



0.283E+00  
 MAXIMUM VECTOR



CONTour FRSH 0.50608E-01 10 0.85551E-02 CONTOUR INTERVAL OF 0.42776E-03 PT(3,31)= 0.33676E-03 LABELS SCALED BY 0.10000E+06  
 KINETIC ENERGY AT 6.67 L




CONTour FRSH -0.31427E-02 10 0.12819 CONTOUR INTERVAL OF 0.65666E-02 PT(3,31)= 0.15583E-22 LABELS SCALED BY 1000.0  
 SCALAR FIELD AT 6.67 L


Aircraft Carrier - 10 kts (5.15 m/s)  
 Stratified  
 $X = 2.77 \text{ km} = 8.34 \text{ L}$




AXIAL VELOCITY AT 8.34 L  
 CONTOUR FROM 5 1283 TO 5 2619 CONTOUR INTERVAL OF 0.61965E-02 PT13.31 = 5 1419



MINIMUM VECTOR  
 CONTOUR FROM -0 11663E-01 TO 0 12072  
 CONTOUR INTERVAL OF 0.66282E-02 PT13.31 = 0 21554E-19



KINETIC ENERGY AT 8.34 L  
 CONTOUR FROM 0.52285E-04 TO 0.82378E-02 CONTOUR INTERVAL OF 0.40928E-03 PT13.31 = 0.31918E-03



SCALAR FIELD AT 8.34 L  
 CONTOUR FROM -0 11663E-01 TO 0 12072 CONTOUR INTERVAL OF 0.66282E-02 PT13.31 = 0 21554E-19

**DIANA::HYMAN**


**JOB 220**

**CVN25.LAS;1**


**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN25.LAS;1  
**Last Modified:** 25-MAY-1995 12:50  
**Owner UIC:** [HYMAN]

**Length:** 1737 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LASER\_B1102C  
**Submitted:** 25-MAY-1995 12:49  
**Printer queue:** LASER\_B1102C  
**Printer device:** LPS17A


Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified X = 0.56 km = 1.69 L




1000 MICRON PART. POPULATION AT 0.56 KM  
 CONTOUR FROM 0.0000E+00 TO 1.9999E+01 CONTOUR INTERVAL OF 0.201438 P1(3,31) = 0.18099E-01



CONCENTRATION FIELD(G/M\*\*3) AT 0.56 KM  
 5000 MICRON PART. POPULATION AT 0.56 KM  
 CONTOUR FROM 0.0000E+00 TO 0.2000E-01 CONTOUR INTERVAL OF 0.31709E-02 P1(3,31) = 0.20000E-09 LABELS SCALED BY 100000




5000 MICRON PART. POPULATION AT 0.56 KM  
 CONTOUR FROM 0.0000E+00 TO 1.9999E+01 CONTOUR INTERVAL OF 0.201438 P1(3,31) = 0.18099E-01




5000 MICRON PART. POPULATION AT 0.56 KM  
 CONTOUR FROM 0.0000E+00 TO 1.9999E+01 CONTOUR INTERVAL OF 0.201438 P1(3,31) = 0.18099E-01

Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified X = 1.06 km = 3.19 L





CONTOUR FROM 0.00000E+00 TO 0.777620 CONTOUR INTERVAL OF 0.155953 PLOT(3,3)= 0759675E-02





CONTOUR FROM 0.00000E+00 TO 0.28228E-01 CONTOUR INTERVAL OF 0.06469E-02 PLOT(3,3)= 0.28886E-02 LABELS SCALED BY 100000

Aircraft Carrier - 20 knots (2.8/3 m/s)  
 Unstratified  
 X = 2.06 km = 6.20 L







CONTOUR FROM 0.0000E+00 TO 0.0825E8 CONTOUR INTERVAL OF 0.0125E8 PT(3,31)= 0.068998E-02



CONTOUR FROM 0.0000E+00 TO 0.0800E+01 CONTOUR INTERVAL OF 0.0000E+00 PT(3,31)= 0.06197E-03 LABELS SCALED BY 10000.





Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified X = 3.06 km = 9.22 L



CONTOUR FROM 0.4688E-01 TO 0.2551E5 CONTOUR INTERVAL OF 0.2551E5

CONTOUR FROM 0.0000E+00 TO 0.2551E5 CONTOUR INTERVAL OF 0.2551E5

Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified X = 5.06 km = 15.24 L



**DIANA::HYMAN**


**JOB 72**

**CVN25.LAS;1**


**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN25.LAS;1  
**Last Modified:** 8-JUN-1995 15:09  
**Owner UIC:** [HYMAN]

**Length:** 7539 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 8-JUN-1995 15:09  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER


Aircraft Carrier - 25 kts (12.875 m/s)  
Unstratified  
X = 0.55 km = 1.67 L




CONTOUR FROM 12.395 TO 13.977 CONTOUR INTERVAL OF 0.79068E-01 PT(13,31)= 12.663  
AXIAL VELOCITY AT 1.67 L



0.50ZF(00)  
MAXIMUM VECTOR







CONTOUR FROM 0.10000E-03 TO 0.40284 CONTOUR INTERVAL OF 0.21075E-01 PT(13,31)= 0.97797E-03 LABELS SCALED BY 1000.0  
KINETIC ENERGY AT 1.67 L




CONTOUR FROM -0.15857E-03 TO 0.16824 CONTOUR INTERVAL OF 0.84188E-02 PT(13,31)= -0.71689E-37 LABELS SCALED BY 1000.0  
SCALAR FIELD AT 1.67 L


Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified  
 $X = 1.11 \text{ km} = 3.34 \text{ L}$




Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified  
 $X = 2.21 \text{ km} = 6.67 \text{ L}$




CONTour FROM 12.691 TO 13.332 CONTOUR INTERVAL OF 0.32408E-01 PT(3,31)= 12.689



0.255E+00  
 MAXIMUM VECTOR







CONTour FROM 0.1000E-03 TO 0.56818E-01 CONTOUR INTERVAL OF 0.29862E-02 PT(3,31)= 0.61273E-03 LABELS SCALED BY 10000.




CONTour FROM -0.34033E-03 TO 0.67081E-01 CONTOUR INTERVAL OF 0.33700E-02 PT(3,31)= 0.58078E-33 LABELS SCALED BY 10000.


Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified  
 $X = 3.32 \text{ km} = 10 \text{ L}$




Aircraft Carrier - 25 kts (12.875 m/s)  
 Unstratified  
 X = 5.53 km = 16.67 L




CONTOUR FROM 12.754 TO 13.229 CONTOUR INTERVAL OF 0.23775E-01 PT(3,31): 12.889  
 AXIAL VELOCITY AT 16.67 L



0.155E+00  
 HORIZONTAL VECTOR



CONTOUR FROM 0.10000E-03 TO 0.16781E-01 CONTOUR INTERVAL OF 0.83409E-03 PT(3,31): 0.25917E-03 LABELS SCALED BY 10000.  
 KINETIC ENERGY AT 16.67 L



CONTOUR FROM -0.36179E-01 TO 0.36953E-01 CONTOUR INTERVAL OF 0.18699E-02 PT(3,31): -0.1145E-25 LABELS SCALED BY 10000.  
 SCALAR FIELD AT 16.67 L

**DIANA::HYMAN**

**JOB 1221**





**CVN25-STRAT.LAS;1**

**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]CVN25-STRAT.LAS;1  
**Last Modified:** 9-JUN-1995 15:31  
**Owner UIC:** [HYMAN]

**Length:** 6533 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 9-JUN-1995 15:31  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER





Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified X = 1.06 km = 3.19 L





CONTOUR PLOT OF CONCENTRATION FIELD (G/M\*\*3) HT 1.06 KM  
 CONTOUR INTERVAL OF 0.0000E+00 TO 0.149500 PLOT OF 022705 010405E-02

Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified  
 X = 2.06 km = 6.20 L







CONTOUR FROM 0.00000E+00 TO 0.650550 CONTOUR INTERVAL OF 0.100000 PLOT LABELS SCALED BY 10000.



CONTOUR FROM 0.00000E+00 TO 0.72000E+01 CONTOUR INTERVAL OF 0.20000E+02 PLOT LABELS SCALED BY 10000.



Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified  
 X = 3.06 km = 9.22 L





CONTOUR FROM 0.00000E+00 TO 0.16088E-01 CONTOUR INTERVAL OF 0.28888E-02 PT(3,31)= 0.67998E-03 LABELS SCALED BY 10000.

CONTOUR FROM 0.00000E+00 TO 0.88888E-02 CONTOUR INTERVAL OF 0.18121E-01 PT(3,31)= 0.80888E-02

Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified X = 5.06 km = 15.24 L







CENTOUR FROM 0 00000E+00 TO 0380708 CWINTER INTERVAL OF 0757552E-01 P T I 3.31= 0.28875E-02







CENTOUR FROM 0.00000E+00 TO 0 98609E-02 CWINTER INTERVAL OF 0.28809E-02 P T I 3.31= 0.58639E-08 LABELS SCALED BY 0100000E+08





Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified  
 X = 0.55 km = 1.67 L




Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified  
 $X = 1.11 \text{ km} = 3.34 \text{ L}$




Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified  
 $X = 2.21 \text{ km} = 6.67 \text{ L}$




Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified  
 X = 3.32 km = 10 L




CONTour FROM 12.635 TO 13.272 CONTOUR INTERVAL OF 0.316205-01 P1(3.31)= 12.869  
 AXIAL VELOCITY AT 10.00 L



0.185E+00  
 MAXIMUM VELOCITY







CONTour FROM 0.10000E-03 TO 0.29100E-01 CONTOUR INTERVAL OF 0.14500E-02 P1(3.31)= 0.49257E-03 LABELS SCALED BY 10000.  
 KINETIC ENERGY AT 10.00 L



CONTour FROM -0.12120E-02 TO 0.45017E-01 CONTOUR INTERVAL OF 0.23115E-02 P1(3.31)= -0.69135E-28 LABELS SCALED BY 10000.  
 SCALAR FIELD AT 10.00 L

Aircraft Carrier - 25 kts (12.875 m/s)  
 Stratified  
 X = 5.53 km = 16.67 L



**DIANA::HYMAN**

**JOB 703**

**FFG25.LAS;1**


**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]FFG25.LAS;1  
**Last Modified:** 13-JUN-1995 08:39  
**Owner UIC:** [HYMAN]

**Length:** 1734 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 13-JUN-1995 08:39  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER


Frigate - 25 kts (12.875 m/s)

Unstratified



X = 0.56 km = 4.26 L



CONTOUR FROM 0.0000E+00 TO 0.248698E-01 P113.31= 0238999E-02



CONTOUR FROM 0.0000E+00 TO 0.42086E-02 P113.31= 0.8888E-08 LABELS SCALED BY 0100000E+06







CONTOUR FROM 0.0000E+00 TO 0.248698E-01 P113.31= 0238999E-02

CONTOUR FROM 0.0000E+00 TO 0.42086E-02 P113.31= 0.8888E-08 LABELS SCALED BY 0100000E+06

Frigate - 25 kts (12.875 m/s)





Unstratified X = 1.06 km = 8.07 L



CONTOUR FROM 0.00000E+00 TO 0.99999E-02 CENTER INTERVAL OF 0.10000E-02 P1(3,3) = 0.99999E-03 LABELS SCALED BY 0100000E+05

CONTOUR FROM 0.00000E+00 TO 0.99999E-01 P1(3,3) = 0.128128E-02

Frigate - 25 kts (12.875 m/s)  
 Unstratified  
 X = 2.06 km = 15.68 L




CONTOUR FROM 0.00000E+00 TO 0.47893E-02 CONTOUR INTERVAL OF 0.47893E-02 LABELS SCALED BY 0.10000E+06



CONTOUR FROM 0.00000E+00 TO 0.12500E+01 CONTOUR INTERVAL OF 0.25000E+01 LABELS SCALED BY 0.10000E+06




..rigate - 25 kts (12.875 m/s)  
 Unstratified  
 X = 5.06 km = 38.52 L







CONTOUR FROM 0.00000E+00 TO 0.00000E+02 INTERVAL 0.00000E+00 BY 10000.




CONTOUR FROM 0.00000E+00 TO 0.00000E+02 INTERVAL 0.00000E+00 BY 0.10000E+06




Frigate - 25 kts (12.875 m/s)  
 Unstratified X = 0.66 km = 5.01 L




Frigate - 25 kts (12.875 m/s)  
Unstratified  
X = 1.32 km = 10.01 L




CONTOUR FROM 12.565 TO 13.550 CONTOUR INTERVAL OF 0.9223E-01 P113.31= 12.869



0.105E+00  
MAXIMUM VECTOR







CONTOUR FROM 0.10000E+03 TO 0.6392E+01 CONTOUR INTERVAL OF 0.3161E-02 P113.31= 0.47615E-03 LABELS SCALED BY 10000.







CONTOUR FROM -0.77590E+06 TO 0.40266E+01 CONTOUR INTERVAL OF 0.20133E-02 P113.31= 0.00000E+00 LABELS SCALED BY 10000.

Frigate - 25 kts (12.875 m/s)

Unstratified X = 2.63 km = 20.01 L







Frigate - 25 kts (12.875 m/s)  
Unstratified  
X = 3.94 km = 30.01 L



Frigate - 25 kts (12.875 m/s)

Unstratified X = 6.57 km = 50.01 L



**DIANA::HYMAN**


**JOB 691**

**FFG25-STRAT.LAS;1**


**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]FFG25-STRAT.LAS;1  
**Last Modified:** 13-JUN-1995 08:28  
**Owner UIC:** [HYMAN]

**Length:** 1767 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 13-JUN-1995 08:28  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER


Frigate - 20 kts (2.8/3 m/s)  
 Stratified  
 $X = 0.66 \text{ km} = 5.01 \text{ L}$




CONTOUR FROM 12.392 TO 14.122 CONTOUR INTERVAL OF 0.06194E-01 PT(3,3)= 12.069  
 AXIAL VELOCITY AT 5.01 L



0.205E-00  
 MAXIMUM VECTOR







CONTOUR FROM 0.10000E-03 TO 0.19142 CONTOUR INTERVAL OF 0.5681E-02 PT(3,3)= 0.52787E-03 LABELS SCALED BY 1000.0  
 KINETIC ENERGY AT 5.01 L




CONTOUR FROM -0.5576E-04 TO 0.5172E-01 CONTOUR INTERVAL OF 0.25897E-02 PT(3,3)= 0.00000E-00 LABELS SCALED BY 10000.  
 SCALAR FIELD AT 5.01 L




Frigate - 25 kts (12.875 m/s)  
 Stratified  
 $X = 1.32 \text{ km} = 10.01 \text{ L}$



Frigate - 25 kts (12.875 m/s)  
 Stratified  
 X = 2.63 km = 20.01 L






~ 0 0









Frigate - 25 kts (12.875 m/s)  
Stratified  
X = 6.57 km = 50.01 L







Frigate - 25 kts (12.875 m/s)

Stratified  $X = 0.56 \text{ km} = 4.26 \text{ L}$



Frigate - 25 kts (12.875 m/s)



Stratified X = 1.06 km = 8.07 L





Frigate - 25 kts (12.875 m/s)

Stratified

X = 5.06 km = 38.52 L







CONTOUR FREQ 0.00000E+00 TO 0.00000E+02 CENTER INTERVAL OF 0.00000E+00







CONTOUR FREQ 0.00000E+00 TO 0.00000E+02 CENTER INTERVAL OF 0.00000E+00

Frigate - 25 kts (12.875 m/s)

Stratified  
X = 2.06 km = 15.68 L



Frigate - 25 kts (12.875 m/s)  
 Stratified X = 3.06 km = 23.29 L



**DIANA::HYMAN**



**JOB 1784**

**FFG10-STRAT.LAS;1**


**File:** \_\$40\$DUA29:[HYMAN.GRID.DISPERSION]FFG10-STRAT.LAS;1  
**Last Modified:** 12-JUN-1995 13:43  
**Owner UIC:** [HYMAN]

**Length:** 1826 blocks  
**Longest record:** 27 bytes  
**Priority:** 100  
**Submit queue:** LPS40\$LAZER  
**Submitted:** 12-JUN-1995 13:43  
**Printer queue:** LPS40\$LAZER  
**Printer device:** LAZER

Stratified  
 X = 0.56 km = 4.26 L







CONTOUR FROM 0.00000E+00 TO 0.7616598 CONTOUR INTERVAL OF 0.152338 P113.31= 0.81658E-02



CONTOUR FROM 0.00000E+00 TO 0.46809E-01 CONTOUR INTERVAL OF 0.30940E-02 P113.31= 0.45658E-03 LABELS SCALED BY 10000.

Frigate - 10 kts (5.15 m/s)  
 Stratified  
 $X = 1.06 \text{ km} = 8.07 \text{ L}$







CONTOUR FROM 0.00000E+00 TO 0.00000E+01 CENTER INTERVAL OF 0.00000E+00 P113.31= 0.0000E-03 LABELS SCALED BY 010000E+06

CONTOUR FROM 0.00000E+00 TO 0.00000E+01 CENTER INTERVAL OF 0.00000E+00 P113.31= 0.0000E-02

Frigate - 10 kts (5.15 m/s)

X = 1.56 km = 11.87 L

Stratified







CONTR. FIGH 0.0000E+00 TO 0.9999E-02 CONTOUR INTERVAL OF 0.0000E-02 PT(3,3)= 0.45168E-02

CONTR. FIGH 0.0000E+00 TO 0.9999E-02 CONTOUR INTERVAL OF 0.2000E-02 PT(3,3)= 0.99979E-03 LABELS SCALED BY 0.100000E+06



Frigate - 10 kts (5.15 m/s)

Stratified



X = 2.06 km = 15.68 L



Frigate - 10 kts (5.15 m/s)  
 Stratified  
 X = 3.06 km = 23.29 L



CONTOUR FROM 0.00000E+00 TO 0.28190E-02



CONTOUR FROM -0.00000E+00 TO 0.84889E-02

## **APPENDIX I**

### **AMBIENT DISPERSION MODELING RESULTS**

Source: Ambient Dispersion Modeling.  
San Diego, California

# **FAR FIELD DISPERSION OF PAPER PARTICULATES FROM SURFACE VESSEL DISCHARGES**

by **Scott A. Jenkins, Ph.D. and Joseph Wasyl**

## **I. INTRODUCTION**

This study effort performs numerical dilution modeling of paper particulate discharges in the far field of surface vessel wakes. These dilution simulations account for ambient advection and mixing effects at a sufficiently large distance downstream from the vessel that residual mixing from the wake turbulence is negligible. The ambient advection and mixing effects are associated with the action of wind stress throughout the depths of the mixed layer, surface wave boundary layers due to sea and swell conditions, interfacial shear from current variations across the pycnocline, hindered settling from density variations across the pycnocline, and bottom current boundary layers associated with seafloor roughness. The far field advection-mixing boundary value problem is specified for characteristic ambient conditions for the central Baltic Sea in winter and for the southern portions of the North Sea in summer. The particle distributions used to initialize the ambient advection-mixing code (SEDXPORT) were provided by the far field cross-wake data plane from the wake code, TBWAKE.

The SEDXPORT code used to calculate the far field particle dilution has evolved from a sediment transport model originally developed for non-interacting spherical particles in a field of gravity waves (Jenkins & Inman, 1985). It was subsequently expanded to include dispersion by both currents and waves, and was adapted to solve problems of sewage dispersion for the State of California Water Resources Control Board (Jenkins, Nichols & Skelly, 1989). The model was further refined to include cohesive particle dynamics in problems of scour and erosion of muddy seabeds (Jenkins & Wasyl, 1990), and hindered settling dynamics due to particle-to-particle stress transfer in high concentration suspensions (Aijaz & Jenkins, 1994). Recently the model has been expanded to include vertical stratification of the water column due to river plumes and mixed layer dynamics, calculating features of hindered settleline layers at the pycnocline interface and bottom turbid layers. In this most recent version, the model has been integrated into the Navy's Coastal Water Clarity Model (CWC) and the Littoral Remote Sensing Simulator (LRSS) (see Hammond et al, 1995). The SEDXPORT code has been validated for

relatively small (less than 30 microns) optically active particles in mid-to-inner shelf waters (see Hammond et al, 1995, and Schoonmaker et al, 1994). Validation of the SEDXPORT code was shown by three independent methods: 1) direct measurement of particle numbers and particle size distributions by means of a laser particle sizer, 2) measurements of water column optical properties, and 3) comparison of computed particle dispersion patterns with LANDSAT imagery.

The SEDXPORT code is a time stepped finite element model which solves the advection diffusion equations over a fully configurable 3-dimensional grid. The vertical dimension is treated as a two-layer ocean, with a homogeneous surface mixed layer and a homogeneous bottom layer separated by a pycnocline interface. The code accepts any arbitrary density and velocity contrast between the mixed layer and bottom layer that satisfies the Richardson number and Froude number stability criteria. The code does not time split advection and diffusion calculations, and will compute additional advective field effects arising from spatial gradients in eddy diffusivity, i.e., the so-called "gradient eddy diffusivity velocities" after Armi (1979). Eddy mass diffusivities are calculated from momentum diffusivities by means of a series of Peclet number corrections based upon particle size and mass and upon the mixing source. Peclet number corrections for the surface and bottom boundary layers are derived from the work of Nielsen (1979), Jenssen & Carlson (1976), and Jenkins & Wasyl (1990). Peclet number correction for the wind-induced mixed layer diffusivities are calculated from algorithms developed by Martin and Meiburg (1994), while Peclet number corrections to the interfacial shear at the pycnocline are derived from Lazaro and Lasheras (1992a and 1992b). The momentum diffusivities to which these Peclet number corrections are applied are due to Thoracle (1914), Schmidt (1917), Durst (1924) and Newman (1952) for the wind-induced mixed layer turbulence and to List et al (1990) for the current-induced turbulence.

## 2. GRIDDING AND INITIALIZATION

In the far field, the ship wake is treated as an infinite line source particles. Therefore, the ambient advection mixing problem was treated as 2-dimensional in the cross-wake plane. SEDXPORT was gridded in a 200 x 200 YZ-computational domain with 1.0 meter depth increments (Z-dimension) and 1.5 meter horizontal increments (Y-dimension). This allowed the cross-wake data plane of the initial particle distribution from the TBWAKE code to be nested

inside the far field grid using compatible grid cell dimensions. The TBWAKE YZ-data plane was 99 x 41 in 1.5 x 1.0 meter grid cells, and was centered inside the SEDXPORT grid with the sea surface at  $Z = 0$ . The remaining portions of the 200 x 200 SEDXPORT grid not occupied by the TBWAKE grid were initialized with zero particles at time  $t = 0$ .

Time step lengths varied depending upon the size of particulates. SEDXPORT computes advection-mixing dynamics independently for each of the size fractions which make up the particulate distribution, because each size fraction has a different Peclet number and corresponding diffusivity. Typically, time step lengths would also be controlled by the mean currents; but because no information is available on the spatial structure of the current field for the particular sites nor the orientation of the ship track relative to the currents, the currents are assumed to be normal to the computational plane, i.e., parallel to the axis of the wake. Similarly, the wave propagation is also assumed to be parallel to the ship track. Consequently, particle advection is exclusively due to settling and the only dynamic influence of the currents is to enhance diffusivities by interfacial shear at the pycnocline or by current boundary layer turbulence at the bottom. The particle size distribution was divided into 9 particle size bins according to the following mass fractions and corresponding settling velocities and time step lengths (see Table 1).

TABLE 1: PARTICLE SIZE PROPERTIES

Particle Size (microns)	Mass Fraction	Settling Velocity (cm/sec)	Time Step Length (sec)
200	0.088	0.048	7000
500	0.111	0.241	1120
800	0.105	0.296	437
1000	0.067	0.391	280
1200	0.066	0.391	194.4
1400	0.065	1.227	142.8
1600	0.064	2.095	109.4
1800	0.063	2.992	86.4
3000	0.367	7.713	31.1

The dry particle density was assumed to be the same for all size bins and was taken as  $1.54 \text{ gm/cm}^3$ . Note that the predominant mass fraction belonged to the largest size bin of 3000 microns, and that these large particles had significantly large settling velocities, 0 (7.7 cm/sec). The time step lengths indicated by Table 1 insured that particles moved only a few grid cells (less than 4) in a time step to maintain numerical stability of the resulting solutions. Because the larger particles advect vertically faster due to gravity, they require shorter time step intervals.

The environmental conditions for the Baltic and North Seas are specified according to characteristic "climate atlas" figures provided by NRaD. SEDXPORT boundary conditions and forcing function inputs derived from these climate atlas figures are shown in Table 2 below:

TABLE 2: BOUNDARY CONDITIONS AND INPUT FORCING

	Central Baltic (Winter)	Southern North Sea (Summer)
Depth	200 m	50 m
Swell Height	0 m	0 m
Swell Period	0 sec	0 sec
Wind Wave Height	2.0 m	0.5 m
Wind Wave Period	6.0 sec	6.0 sec
Winds	9.0 m/sec	5.0 m/sec
Mixed Layer Depth	50 m	50 m
Mixed Layer Density	6.0 sigma t	14.6 sigma t
Bottom Layer Density	12.0 sigma t	24.6 sigma t
Mixed Layer Current	0 cm/sec	10 cm/sec
Bottom Layer Current	0 cm/sec	0 cm/sec
Bottom Roughness	2.0 cm	10.0 cm

In both the Baltic and North Sea simulations, no specific bathymetry was evaluated for effects on wave shoaling or current boundary layers; and thus horizontal gradients in diffusivity were not possible. The absence of such gradients insures that the 2-dimensional YZ-computational plane remains adequate for representation of the problem. The bottom was treated as a flat plane boundary with random roughness of height indicated in Table 2.

### III. RESULTS

Cross wake particle dispersion simulations from SEDXPORT are plotted in Appendix A for the winter time Baltic Sea conditions, and in Appendix B for the summer time North Sea conditions. In each of these appendices, dispersion plots are given for three separate particle size bins spanning the complete particle size distribution shown in Table 1, namely 200 micron, 1200 micron and 3000 micron size particles, each with specific gravities of 1.54. Each particle size dispersion pattern begins from the initial far field distribution from TBWAKE at time  $t = 0$ . All

plots have depth on the vertical axis with the surface appearing at the top of the plot at  $Z = 0.0$ . For the Baltic cases in Appendix A, the full vertical scale is from 0 to 200 meters while for the North Sea cases in Appendix B, the full range vertical scale is from 0 to 50 meters. In both appendices, the horizontal scale is in terms of grid cell number across the wake from port to starboard. Thus the full range horizontal scale is from 0 meters on port to 300 meters on starboard, with the ship wake centered at grid cell #100, or at  $Y = 150$  meters. Thus all dispersion plots have a vertical exaggeration, which was 1.8 to 1 for the Baltic and 7.2 to 1 for the North Sea. All dispersion plots have been dynamically scaled in terms of particle number concentrations.

The Baltic cases in Appendix A give the worst-case scenarios in terms of arrival time at the seafloor, but the best-case scenarios for minimum dilution. This is due to the greater depths, higher sea states and stronger winds of the Baltic relative to the North Sea. All particle size fractions are released from separate port and starboard discharges which appear in the initial distributions as four distinct "blobs", two on the port side and two on the starboard side of the wake centerline. In all initial far field distributions, the port side blobs have greater particle abundance than the starboard side blobs. It is interesting how the 200 micron size particles in the Baltic conditions begin merging of the port and starboard blobs as they pass through the hindered settling regime at the 50-meter depth pycnocline. Below the pycnocline, the port and starboard blobs become fully merged, with an asymmetric center of mass displaced to the port side of the wake. The larger size particles which fall faster and have smaller diffusivities. They are less well-mixed, and the dispersion patterns continue to show distinct port and starboard blobs throughout the residence time in the water column.

In the Baltic, the 200 micron size particles begin with 0 (20,000 particles/m<sup>3</sup>) at the surface at time zero and dilute about 20 to 1 while passing through the mixed layer. They begin arriving at the bottom after about 78 hours while the center of mass of the dispersion pattern has reached depths of about 120 meters and has diluted to 0 (500 particles/m<sup>3</sup>) or 40:1 minimum dilution. The center of mass of the 200 micron size particles reaches the bottom after about 117 hours and has diluted to 0 (250 particles/m<sup>3</sup>) or a minimum dilution of 80 to 1. By contrast, the larger size fractions have much shorter bottom arrival times and less dilution. The center of mass of the 1200 micron particles reaches the bottom in 10.8 hours with a minimum dilution of 20 to

1. The 300 micron size particles which account for 37% of the total mass reach the bottom in 41 minutes with a minimum dilution of only 10 to 1.

In the calmer conditions and shallower depths of the North Sea in summer, there is much less ambient mixing and correspondingly less dilution. Inspection of the dispersion plots in Appendix B reveals much less merging of blobs and less lateral dispersion (although the vertical exaggeration is also greater in these plots). Here the center of mass of the 200 micron size particles reaches the bottom in about 29 hours with a minimum dilution of 20 to 1. The 1200 micron size particles reach the bottom in 97 minutes at 7 to 1 dilution, while the 3000 micron size particles reach the bottom in only 10 minutes at 5 to 1 dilution.

## REFERENCES

- Armi, L.A., 1979, "Effects of variations in eddy diffusivity on property distributions in the oceans," *Journal of Marine Res.*, v. 37, n. 3, pp. 515-530.
- Durst, C.S., "The relationship between current and wind," *Quart. J. R. Met. Soc.*, v. 50, p. 113 (London), 1924.
- Jenkins, S.A., & D.L. Inman, "On a submerged sphere in a viscous fluid excited by small-amplitude motions," *Jour. Fluid Mech.*, v. 157, pp. 199-224.
- Jenkins, S.A., J.A. Nichols & D.W. Skelly, 1989, "Coupled physical-biological dispersion model for the fate of suspended solids in sewage discharged into the ocean," SIO Reference Series 89-3, 53 pp.
- Jenkins, S.A., & J. Wasyl, 1990, "Resuspension of estuarial sediments by tethered wings," *Jour. of Coastal Res.*, v. 6, n. 4, pp. 961-980.
- Jenkins, S.A., & S. Aijaz, 1994, "On the electrokinetics of the dynamic shear stress in fluid mud suspensions," *Jour. Geophysical Res.*, v. 99, n. C6, pp. 12,697-12-706.
- Jenkins, S.A., & J. Wasyl, 1994, "Time stepped suspended transport model for the dispersion of optical particles in coastal waters," submitted to Office of Naval Research, Code 1153, 84 pp.
- Hammond, R.R., Jenkins, S.A., Cleveland, J.S., Talcott, J.C., Heath, A.L., Wasyl, J., Goosby, S.G., Schmitt, K.F., and Levin, L.A., 1995, "Coastal Water Clarity Modeling," SAIC, Technical Report 01-1349-03-4841-000, 491 pp.
- Lazaro, B.J. & Lasheras, J.C., 1992a, "Particle dispersion in a developing free shear layer, Part 1, Unforced flow," *J. Fluid Mech.* 235, pp. 143-178.
- Lazaro, B.J. & Lasheras, J.C., 1992b, "Particle dispersion in a developing free shear layer, Part 2, Forced flow," *J. Fluid Mech.* 235, pp. 179-221.
- List, E.J., Gartrell, G. & C.D. Winant, 1990, "Diffusion and dispersion in coastal waters," *Jour. Hydraulic Eng.*, v. 116, n. 10, pp. 1158-1179.
- Martin, J.E., & Meiburg, E., 1994, "The accumulation and dispersion of heavy particles in forced two-dimensional mixing layers, 1: The fundamental and subharmonic cases," *Phys. Fluids*, A-6, pp. 1116-1132.
- Neumann, G., "Ober die komplexe Natur des Seeganges, Teil 1 and 2," *Deut. Hydrogr. Zeit.*, v. 5, n. 2/3, pp. 95-110, n. 5/6, pp. 252-277, 1952.

- Nielsen, P., I.A. Svendsen, & C. Staub, "Onshore-offshore sediment movement on a beach," in Proceedings of 16th International Conference of Coastal Engineering, pp. 1475-1492, ASCE, New York, 1978.
- Nielsen, P., "Some basic concepts of wave sediment transport," Series Paper No. 20, Institute of Hydrodyn. and Hydr. Eng., Techn. Univ. of Denmark, 1979.
- Schoonmaker, J.S., R.R. Hammond, A.L. Heath & J.S. Cleveland, 1994, "A numerical model for prediction of sub-littoral optical visibility," *SPIE Ocean Optics XII*, pp. 18.
- Thorade, H., "Die Geschwindigkeit von Triftströmungen and die Ekman'sche Theorie," 1914, *Ann. d. Hydr. u. Marit. Meteorol.*, v. 42, p. 379.

**APPENDIX A:**


**CROSS-WAKE PARTICLE DISPERSION PLOTS FOR WINTER TIME CONDITIONS  
IN THE BALTIC SEA**

**VERTICAL EXAGGERATION = 1.8 TO 1.0**


FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.02 cm PARTICLES, t=0 hrs




Port-Starboard grid Cells, (1.5 m / grid cell)




Particle Numbers / Cubic Meter




Port-Starboard grid Cells, (1.5 m / grid cell )



Particle Numbers / Cubic Meter




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.02 cm PARTICLES, t=117 hrs




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.12 cm PARTICLES, t=0




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.12 cm PARTICLES, t=4.3 hrs




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.12 cm PARTICLES, t=1.8hr




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.12 cm PARTICLES, t=10.8 hrs




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.12 cm PARTICLES, t=6.5 hrs





Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.3 cm PARTICLES, t=0




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.3 cm PARTICLES, t=11.7 min




Port-Starboard grid Cells, ( 1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.3 cm PARTICLES, t=27.6 min




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: BALTIC; JANUARY, 0.3 cm PARTICLES, t=41.5 min



Port-Starboard grid Cells, (1.5 m / grid cell)




Particle Numbers / Cubic Meter

**APPENDIX B:**


**CROSS-WAKE PARTICLE DISPERSION PLOTS FOR SUMMER TIME CONDITIONS  
IN THE SOUTHERN NORTH SEA**

VERTICAL EXAGGERATION = 7.2 TO 1.0

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.02 cm PARTICLES, t=0




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.02 cm PARTICLES, t=9.7 hrs




Port-Starboard grid Cells, (1.5 m / grid cell)




Particle Numbers / Cubic Meter


FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.02 cm PARTICLES, t=19.4 hrs




Port-Starboard grid Cells, (1.5 m / grid cell )



Particle Numbers / Cubic Meter




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.12 cm PARTICLES, t=0




Port-Starboard grid Cells, (1.5 m / grid cell )



Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.12 cm PARTICLES, t=32.4 min




Port-Starboard grid Cells, (1.5 m / grid cell)




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.12 cm PARTICLES, t=64.8 min




Port-Starboard grid Cells, (1.5 m / grid cell )



Particle Numbers / Cubic Meter




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.3 cm PARTICLES, t=0




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.3 cm PARTICLES, t=3.4 min




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.3 cm PARTICLES, t=6.9 min




Port-Starboard grid Cells, (1.5 m / grid cell )




Particle Numbers / Cubic Meter

FAR FIELD PARTICLE DISPERSION: NORTH SEA; JULY, 0.3 cm PARTICLES, t=10.4 min



Port-Starboard grid Cells, (1.5 m / grid cell )



Particle Numbers / Cubic Meter

## **APPENDIX J**

### **FIELD STUDIES PLAN/ACQUISITION REPORT**

**Source:** Field Studies Plan.  
San Diego, California  
Naval Command, Control & Ocean Surveillance  
Center, RDTE Division, Code 522, 1995

## SOLID WASTE FIELD SAMPLING PLANS

### GENERAL DESCRIPTION

An at-sea test will be conducted late January 1995 approximately 2 miles offshore Coronado (approximate central position: 32° 37.09'N, 117° 10.90' W) to measure the dispersion of pulped cellulose and dye as part of the Shipboard Solid Waste Discharge Project. Three vessels, and one helicopter will be involved: The R/V ECOS, the R/V ACOUSTIC EXPLORER, a 21' skiff, and a helicopter hired through the photoshop, topside. The ACOUSTIC EXPLORER will serve as a discharge vessel, discharging pulped cellulose material and rhodamine fluorescent dye in a liquid slurry while transiting at approximately 10 knots along a 1 km line. The ECOS, will serve as a measurement vessel, taking a suite of measurements in the wake of the ACOUSTIC EXPLORER after the material has been discharged. The helicopter will be used to obtain aerial photography of the wake and nearby area for a period of time following the discharge in order to visually record the dispersion of the material. The skiff will be used for transferring personnel, and for making some measurements such as plankton net tows.

### ADMINISTRATIVE ISSUES

- |  |                 |
|--|-----------------|
| 1. Letters to agencies for "permits"                                     | Stacey          |
| 2. Hiring of the Acoustic Explorer (late January)                        | Stacey          |
| 3. Purchase of dye (16 gallons of 20%)                                   | Stacey          |
| 4. Chemical analyses Stub (BOD/TOC/NUTS)                                 | Stacey/Bart     |
| 5. MESC equipment repairs (cable, haul-out, fathometer)                  | Brad/Chuck/Bart |
| 6. Get paper 100 lbs (also ask where likely pump location will be)       | Stacey/Bart     |
| 7. Helicopter Imaging/photos   | Stacey          |
| 8. Check on other operations in the area (including flight restrictions) | Jim             |
| 9. Fax to Coast Guard  | Jim             |
| 10. Arrange for NaCl from inventory                                      | Chuck           |
| 11. Purchase fluorometer filter kits                                     | Chuck           |
| 12. Obtain drifters  | Bart            |
| 13. Tracor Setup   | Stacey          |
| 14. Sampling Bottles   | Chuck           |
| 15. Talk with Dave re: nutrient analyses                                 | Bart            |

### EXPERIMENT SCALING

#### Initial Scaling

Basis for comparison: Carrier operating pulper discharge @ 1500 lbs/hr (680 kg/hr)

Assumed Carrier beam\*draft = 10 m \* 40 m = 400 m<sup>2</sup>

Acoustic Explorer beam\*draft = 8 m \* 3 m = 24 m<sup>2</sup>

Scaling Factor: 400/24 = 17

Discharge rate off of Acoustic Explorer: (680 kg/hr)/17 = 40 kg/hr

Length scale of the wake: @10 kts (5m/s) = 300 m/min

Revision: 17 January, 1995

Discharging over 3.3 minutes gives a length scale of 1000 m

Discharge over the 3.3 minutes = 2.2 kg paper

Using 10% TSS as guide: 22 kg paper, 222 kg seawater mix = 230 L (61 gal)

Requires pump rate of 70 L/min

Near-Field Wake cross-section size:  $8 \times \text{beam} \times \text{draft} = 8 \times 3 \times 8 = 192 \text{ m}^2$

Volume of plume field =  $192 \text{ m}^2 \times 1000 \text{ m} = 1.92 \times 10^5 \text{ m}^3 = 1.92 \times 10^8 \text{ L}$

Concentration scaling for dye: 16 gallons of 20% Rhodamine WT = 12.1 L dye

Near-field Concentration:  $12.1 / 1.92 \times 10^8 = 6.3 \times 10^{-8} = 63 \text{ ppb}$

### Final Scaling

Pump for 3.3 minutes to create plume of 1 km in length

Use 60.6 L (16 gals) 20% dye

Pump mixture = 61 L of 20% dye, 22 kg paper, 155 L seawater

Total Volume of Pump Mixture = 230 L (61 gal)

Pumping Rate Required: 70 L/min (18.4 gpm)

Starting Concentration of TSS = 10 % (22kg/220 kg)

Theoretical Maximum Dye Concentration in wake = 63 ppb ( $60.6 \text{ L} \times 20\% / 1.92 \times 10^8 \text{ L}$ )

Additional salt needed for salinity adjustment: 2.0 kg ( $33 \text{ g/L} \times 61 \text{ L} = 2013 \text{ g}$ )

## EXPERIMENT CHRONOLOGY

### Date/Time

Date: 24 January 1995

Tide: Low @ 10:10, High @ 16:08, Height @ 1.0 - 3.2'

Times: Leave Dock @ 0700, Deploy Array @ 0800, Start Pre-DischargeMapping @ 0815, Begin Discharge and Mapping @ 0945

### Alternate Date/Time

Date: 25 January 1995

Tide: Low @ 11:29, High @ 17:47, Height @ 0.3 - 3.4'

Times: Leave Dock @ 0800, Deploy Array @ 0900, Start Pre-DischargeMapping @ 0915, Begin Discharge and Mapping @ 1045

### Location of Discharge Line

Start Position: 32° 36.94' N, 117° 11.18' W

End Position: 32° 37.21' N, 117° 10.62' W

Line Direction: 240°

Expected Current Direction: 330°

Expected Current Speed: 0.5 kts

Line Length: 1 km (3.3 mins @ 10 kts)

Water Depth: 20-22 m

Distance to site from NRaD: ~ 11 km

Time to Site from NRaD: ~ 1 hr

### **Location of Current Meter Array Mooring**

Position: 32° 37.09'N, 117° 10.90' W  
Water Depth: 20 m  
Vertical: S4 Current Meters @ 3 m, 10 m, 17 m  
RTM @ 1 m, 3 m, 10 m, 17 m

### **General Chronology**

- 1) Deploy current meter array
- 2) Map out area with ECOS, collecting discrete samples for TSS, NUTS, BOD, and Chl-*a* before arrival of ACOUSTIC EXPLORER; use skiff or ECOS for plankton net tows
- 3) Discharge pulped material and dye from ACOUSTIC EXPLORER, then sample and map plume dispersion with ECOS, skiff, and helicopter, until parameters of interest are back to background

### **EXPERIMENT SPECIFICS**

#### ***Pre-Discharge Mapping***

After deployment of the current meter array, the ECOS will run surface mapping transects parallel to the Discharge Line and then a single perpendicular transect line doing a series of tow-yos (see attached Figure). The perpendicular line will be run such that it crosses near the current meter array and follows the expected direction of the plume (surface current). Seven complete verticals will be performed along this transect. The mapping will cover 13.75 line-kilometers and take approximately 1.5 hrs to complete. All ECOS instrumentation will be used during these transects (see below). Discrete samples will be collected from the towed system for BOD, TSS, TOC, NUTS, and Chl-*a* at locations shown in the attached Figure. Vertical plankton net tows will be run on the skiff and/or the ECOS.

#### ***Discharge Mapping***

The ACOUSTIC EXPLORER will run along the Discharge Line dumping the cellulose and dye waste stream. The discharge will go for approximately 3.3 minutes over a distance of 1 km. As the current meter array is passed, a drifter will be thrown over the side to begin marking the surface advective field.

The ECOS will wait for the ACOUSTIC EXPLORER to pass and dump the drifter, then will begin to map the dispersion of the plume mainly by drifting back and forth through the patch (perpendicular to Discharge Line) performing vertical profiles with its towed system. The Drifter and helicopter sightings will be used to position the ECOS through the center of the plume as it slowly transits back and forth through the patch. The speed through the plume must take into account the space scale expected for the near field plume which is  $3 * \text{beam} = 24 \text{ m}$ . Discrete samples will again be collected from the towed system for BOD, TSS, TOC, NUTS, and Chl-*a* at the locations shown in the attached Figure. Vertical plankton net tows may also be run on the ECOS.

The Helicopter will be used for photographing the dispersion of the plume as well as for sighting the ECOS in the visible patch.

Revision: 17 January, 1995

The skiff will be used to transfer personnel from the ACOUSTIC EXPLORER to the ECOS and make measurements such as vertical plankton net tows.

## **INSTRUMENTATION AND ANALYSES**

1. Hull mounted pumping system with dye fluorometer, Optical Backscatter Sensor if available
2. Standard tow body configuration with CTD, transmissometer, flow-through dye fluorometer
3. Third dye fluorometer @ 3-5 m stationary depth along with transmissometer
4. ADCP
5. Tracor package attached to tow system
6. Three each S4 Current Meters
7. Four each RTMs
8. Plankton net tows from either ECOS or SKIFF
9. 30 each 1 L polycarbonate water bottles for collection of seawater samples for TSS and Nutrients
10. 15 each 0.5 L polyethylene water bottles for collection of seawater samples for Chlorophyll-*a*
11. Sample Bottles for ATI analysis:
  - 30 BOD
  - 30 TOC
  - 5 Nuts (duplicates with in-house measurements)
  - 5 TSS (duplicates with in-house measurements)

## **PERSONNEL**

R/VACOUSTIC EXPLORER: Dumping- Chuck, Other?  
R/V ECOS: Bradley, Andy, Bart, Stacey, Tracor personnel, Schoonmaker  
SKIFF: John, Others (Scripps...)  
HELICOPTER: Photographer, other?  
OTHER: Gerhard (S4 setup)

## **ACOUSTIC EXPLORER**

Identify DGPS capability or install ours  
Obtain vessel specifications including beam, draft, speed, navigation, space, power  
Identify dumping point, available power for pump, deck space for 100 gallon container


## **SKIFF**

Determine if DGPS needed and install  
Setup for hand plankton tows


## **Helicopter Photos**

Identify vendor-Photoshop  
Determine flight restrictions-Done by helo vendor  
Identify photographer and photography system (video/stills include filters etc) - photoshop

**Transect Lines**



**Vertical Profiles**



**Discharge Line**



**Pre-discharge Survey Plan**

## MESC SURVEY DATA ACQUISITION REPORT GAR1

**GOAL:** Field test for Environmental Assessment of Shipboard Discharges Program. The ECOS, skiff, and a helicopter were used to track the dispersion of rhodamine dye and pulper material dumped along a 1 km trackline (by the ACOUSTIC EXPLORER) off the west coast of Coronado.

**DATE:** 27 January, 1995 JD 27

**TIME:** 06:57 - 18:15, Left Dock @ 07:08, Use data from 08:20 - 16:39

**RECORDS:** RTAPS: 13590 (2 second acquisition rate), LABVIEW 30021 (1 second acquisition rate)

**PERSONNEL:** ECOS: Chadwick, Curtis, Davidson, Katz, Patterson, Schoonmaker, Samilo (Tracor)  
22' SKIFF: Groves, Fransham  
ACOUSTIC EXPLORER: Rohr, Katz (during discharge only)  
Helicopter: Jerry Mosley

### SENSOR SYSTEMS CONFIGURATION:

#### TOW SYSTEM

SEABIRD Model 19 CTD w/ *in-situ* pump assembly

SEABIRD pH/DO<sub>2</sub> Sensors (SN 220261)

SEA TECH Transmissometer (SN 298)

TRACOR TAPS (attached to cage)

On-board centrifugal pump with standard teflon hose tow cable # 1 (4 wire only)

Flow: Hose > pump > overflow > bubble trap > Model 10 AU oil fluorometer (SN 5110) > Model 10 chl- $\alpha$  fluorometer (SN 401) > Model 10 AU rhodamine fluorometer (SN 5109)

DELAY: 60 seconds to overflow; 80 seconds to rhodamine fluorometer

#### HULL SYSTEM

WETLABS Multi-wavelength Transmissometer (Schoonmaker's)

Model 10 Rhodamine Fluorometer (SN 0126) w/ broken range output voltage

Submersible bilge pump with teflon hose > overflow valve > fluorometer > transmissometer

DELAY: 10 seconds to fluorometer (use same to overflow valve)

#### BOW SYSTEM (3 m fixed tow depth)

SEA TECH Transmissometer (Lapota's)

Model 10 Rhodamine Fluorometer (SN 5555)

Submersible bilge pump with teflon hose direct to fluorometer with no bypass flow

Transmissometer and pump attached to wire winch off of bow davit (starboard) using a 2' V-fin depressor

DELAY: 38 seconds to fluorometer

### CURRENT METER ARRAY

Three S4s, and four TempMentors deployed at center of discharge line, 32° 37.059' N, 117° 10.968' W  
in 21.7 m of water @ 0755 ~1700

1 RTM SN 900804  
3 m S4 SN 05451194, RTM SN 900958  
10 m S4 SN 04590867, RTM SN 900805  
17 m S4 SN 05451203, RTM SN 900806

### OTHER SENSOR SYSTEMS

ECOS: ADCP- standard hull-mounted configuration

WIND- standard configuration

FATHOMETER- standard configuration

DGPS- standard configuration

SKIFF: DGPS standard self-logging configuration, vertical plankton net tows (30 um)

HELICOPTER: Photos/videos with time stamp

ACOUSTIC EXPLORER: GPS and photos/videos

### **DATA STREAM CONFIGURATION:**

#### TOW SYSTEM

CTD, pH/DO<sub>2</sub>, transmissometer, fathometer to RTAPS

Model 10 AU oil fluorometer (SN 5110) to RTAPS and LABVIEW

Model 10 chl-*a* fluorometer (SN 401) to RTAPS and LABVIEW

Model 10 AU rhodamine fluorometer (SN 5109) to LABVIEW

TRACOR TAPS (self-logging acoustic package)

#### HULL SYSTEM

WETLABS Multiwavelength Transmissometer to standalone computer/files (Schoonmaker)

Model 10 Rhodamine Fluorometer (SN 0126) to LABVIEW

#### BOW SYSTEM (3 m fixed tow depth)

SEA TECH Transmissometer to LABVIEW

Model 10 Rhodamine Fluorometer (SN 555) to LABVIEW

CURRENT METER ARRAY- S4s internal logging, RTMs internal logging

## OTHER

ADCP- to standalone computer/files

WIND- to LABVIEW

FATHOMETER- to RTAPS

DGPS- to RTAPS, LABVIEW, and ADCP

SKIFF- DGPS standard self-logging configuration, discrete samples from plankton tows

HELICOPTER- Photos/videos with time stamp

ACOUSTIC EXPLORER- GPS file, photos/videos

## **DISCRETE SAMPLES:**

30 TSS + 5 (ATI)

22 Chlorophyll *a*

30 Nutrients + 4 (ATI)

30 BOD (ATI)

30 TOC (ATI)

31 Plankton Net Vertical Tows

**TIDE:** Ebb-Slack-Flood; High @ 06:06, Low @ 13:17, High @ 19:41, Range ~ 7.7' (calendar)

## **GENERAL WEATHER CONDITIONS:**

Weather during the survey day was partly cloudy and calm prior to departure. Seas remained calm until and after the discharge of the dye. Afternoon breezes came up and created a 2-3' chop along with a long and 1-2' swell. Previous days of the week were rainy and stormy. The result of the rains was a highly turbid layer, about 3 m deep, in the survey area.

## **GENERAL SURVEY NOTES:**

PRE-DISCHARGE. The ECOS departed the dock with the skiff following @ 0708. The current meter array was deployed @ 0757 and a pre-discharge mapping survey ensued. The ECOS collected both hull and tow system data at the surface. A set of tow-yos were also performed along a track expected to be traveled following the dye release (last transect of the mapping survey). The skiff performed pre-discharge vertical net tows during this time period.

DISCHARGE. At 10:28:55, the ACOUSTIC EXPLORER began discharging pulped paper and dye (see mixture information below). The discharge occurred along a 1 km trackline which ran more or less SW-NE, and passed the current meter array (about midpoint of the trackline). A current drogue was also discharged when passing the current meter array. The discharge ceased at 10:32:52. During the discharge, photos and videos were taken aboard the ACOUSTIC EXPLORER and from a helicopter.

POST-DISCHARGE. After the discharge, the ECOS and skiff moved slowly into the area of the ship wake, transecting the wake in a perpendicular direction. The ECOS mapped the dispersion of dye/pulp etc., attempting to continually cross perpendicular to the wake while following the advective field (using the drogue). The skiff followed the general pattern of the ECOS collecting vertical tows inside and outside the wake area. The post-discharge mapping went on for approximately 6 hours after the discharge. The current meter array was picked up at the end of the mapping. During the first couple of hours after the discharge, photos and videos were taken aboard the ACOUSTIC EXPLORER and from a helicopter.

**GENERAL.** The mixture dumped from the ACOUSTIC EXPLORER was as follows: 61L 20% Rhodamine WT, 27.7 kg (wet weight) of pulped paper (@ 6.3 kg wet/1 kg dry measured), 2 kg NaCl, mixed to a total volume of 231 L in seawater. This mixture provided starting concentrations of 53g/kg (~5%) dye and 19 g/L pulped paper (~2%). The salt was added to bring the salinity up to background, ~33 psu. The paper was obtained from NSW, Carderock. The mixture was pumped out of a large vat using a Jabsco pump @ 59 L/min (measured). See the attached Solid Waste Field Sampling Plan (previous deliverable) for scaling considerations.

Discrete samples for BOD, TSS, Chl-*a*, Nutrients, and TOC were collected throughout the survey aboard the ECOS from either the hull or towed systems. These samples were generally filled in the following order: BOD, TSS, Chl-*a*, Nutrients (2 bottles), TOC from the TSS bottle. When duplicates were taken, they were filled immediately after the first bottle of the same sample type. The first 8 sets of samples were obtained during the pre-discharge mapping, while the remaining 22 sets were obtained after the discharge. Not all sample types were collected during each sampling (in particular, Chl-*a*). Exact start times for each sample can be retrieved from the "Green Book".

Manual range scale changes were made for the hull fluorometer system because of a bad range voltage output. The range changes for this fluorometer were as follows:

TIME RANGE (coarse sensitivity)

0826 31.6 (\*100)  
 1125 31.6 (\*1)  
 1134 31.6 (\*100)  
 1158 1 (\*100)  
 1205 31.6 (\*100)  
 1209 1 (\*100)  
 1552 3.16(\*100)

Sample Suite	Start Time	Comment	Sample Suite	Start Time	Comment
1	08:51:30	Filled from one large container	16	11:23:53	Tow system
2	09:00:00	TSS, NUTS duplicates	17	11:32:26	
3	09:32:20		18	11:52:21	
4	09:52:25		19	11:53:34	TSS, NUTS duplicates (no chl- <i>a</i> )
5	09:56:00	Tow system	20	11:56:20	Tow system
6	10:01:25		21	12:00:00	No chl- <i>a</i>
7	10:05:40	Tow system	22	12:25:04	
8	10:12:41		23	12:26:40	No chl- <i>a</i>
9	10:35:21	After Discharge	24	12:29:00	Tow system, no chl- <i>a</i>
10	10:38:29		25	12:31:53	TSS, NUTS duplicates (no chl- <i>a</i> )
11	10:43:44		26	13:28:14	
12	10:53:46		27	13:30:18	TSS duplicates, no chl- <i>a</i>
13	10:58:12	Tow system	28	13:33:30	Tow system, no chl- <i>a</i>
14	11:18:52	TSS, NUTS duplicates	29	14:20:26	
15	11:22:09		30	14:25:56	No chl- <i>a</i>

**DATA ACQUISITION FILES CREATED:**

ADCP:                   GAR1.CFG Note: Blanking = 1 m, Bottom bin set to 30 m, Salinity to 33.3 psu,  
FN00001 added to file (Also note: possible changes from previous surveys for  
ADCP commands: V 12 cf. 16; J 10 cf. 5; O 109 cf. 77; R 291 cf. 200)  
GAR1001R.000  
GAR1001P.000  
GAR1002R.000 - GAR1002R.008  
GAR1002P.000 - GAR1002P.008

RTAPS:                   GAR1.SDA  
                          GAR1.STA  
                          GAR1.RND  
                          GAR1.INI  
                          GAR1.INM  
                          GAR1.INR

LABVIEW:                GAR1LV.CSV

TRACOR TAPS:           Not yet available

WETLABS:               Not yet available

CURRENT METERS:        GAR01S4.XLS

TEMPMENTORS:          GAR1TEMP.ASC

SKIFF DGPS:            16260270.ASC, .DAT, .EPH, .ION, .MES, .SSF

HELICOPTER:            Photos w/ negatives  
                          Video Tapes (1)

ACOUSTIC EXPLORER:    FDG19027.100 (GPS file)  
                          Photos w/ negatives  
                          Video Tapes (2)

**CALIBRATIONS:**

OIL:   OILCAL.XLS  
CHL:   CHLCAL.XLS  
TSS:   TSSCAL.XLS  
RHO:   GARCALRH.XLS

Full CTD including DO2 and pH Sensor SN 220261 recalibrated for this survey on 24 January 1995.

DO2:           SOC: 2.2267, BOC: -0.0110  
pH:            m: 4.5527, b: 2.6153  
TRANS:        lm: 19.91 lb: -0.279 based on: 0%: 0.014, 100%: 4.60 (91.3% Full Scale in Air)  
OIL:           Calibrated 1 February (After PRF8)

Copied from **GAR1.ini**:

cal

sc,12

oxsocboc,0,2.2267,-0.0110

phmb,0,4.5527,2.6153

trmb,0,19.91,-.279

pressure,0,81.07479,-2.167891e-2,1.813005e-8

cond,0,7.2996e-6,5.0228e-1,-4.1821,5.0761e-4,4.5

temperature,0,3.6756e-3,5.7471e-4,8.3729e-6,-1.774e-6,2350.06

ln,3,63.1037,-5.186

xx,19,-1

fr,6,7

xx,7,100

## **APPENDIX K**

### **SEWAGE EMISSIONS DATA**

Source: Sewage Emissions Data.  
San Diego, California  
Naval Command, Control & Ocean Surveillance  
Center, RDTE Division, Code 522, 1995

WATER QUALITY SAMPLE SUMMARIES

SPT :- 016940AV0F2C NGR :- SJ 74763 96525 DAVYHULME STW FINAL EFFLUENT REQUEST NO 1  
 -----  
 FOR PERIOD 01/01/1990 00:01 TO 04/04/1995 14:01

SAMPLED BY RIVERS

PURP	70	81	82	81	77	172	102	111	118	117	180	92	85	135	143	182	38	47
Temp	Diss	Diss	pH	COND	CL	ALKAL	NH3	N02	N03	Ortno	CCD	BOD	5	Solids	Sols	Silic	Flow	Flow
PURP	Water	O2	O2	as Cl	CaCO3	as N	as N	PO4-	as O	ATU	Susped	N/V	ate	Mean	Inst.			
	-OF	Deg	C	%	mg/L	mg/L	mg/L	mg/L	mg/L	ML/d	ML/d							
MEAN	9.833	7.27	1329	149.4	199.5	10.87	.1135	.5778	1.926	84.42	13.98	19.16	4.434					396.2
S.D.	1.443	.0678	164.1	73.22	4.949	5.349	.2242	1.052	1.714	26.9	9.344	13.13	5.817					175
MAX	11.5	7.35	1560	259	203	31.8	3.34	14	17.2	175	86	158	39					1078
MIN	9	7.2	1172	82	196	<.22	.012	<.1	<.05	33	<2.8	<2	1					<1E-4
NO.OF OCCURS.	3	4	4	5	2	244	240	240	226	192	241	246	79					141
NO. GT.											1							
NO. LT.											13	2	13					

DATA RECORDED AS BEING LESS THAN LIMIT OF DETECTION - TAKEN AS 2/3 RECORDED VALUE  
 IF NO: OF LT'S EXCEEDS 33% TREAT DATA WITH CAUTION--IF MORE THAN 50% CONSULT STATISTICS DEPT.  
 END OF LISTING OF FILE :NHQTEBI.DAT(1,\*,1).JAM1035(1) FOR USER :NHQTEBI AT 1995/04/04 14:10:57

WATER QUALITY SAMPLE SUMMARIES

SPT :- 010925101240102 NGR :- SJ 49.10 2055 WIDNES STW FINAL EFFLUENT SAMPLED AT OUTLET OF PRIMARY TANKS  
 WIDNES STW OFF HALE GATE LANE HALE BANK WIDNES  
 FOR PERIOD 01/01/1990 00:01 TO 06/04/1995 17:08 REQUEST NO 1

PURP	76	81	82	01	77	172	102	111	118	117	100	92	85	135	143	182	38	47	
Temp	Diss	Diss	pH	Cond	CL	Alkal	NH3	NO2	NO3	Ortno	COO	BOD	5	Solid	Sols	Silic	Flow	Flow	
PURP	Water	O2	O2	as	CL	CaCO3	as	N	as	N	P04	as	C	ATU	Suspd	N/V	ate	Mean	Inst.
OF	Deq	C %	mg/L	ML/d	ML/d	ML/d	ML/d												
MEAN	3.444	8.49	3995	649	38.02	1.31	7.302	9.590	397.6	176.6	112.8	47.16	68.93						
STD	.527	1.12	190.0	111.7	17.09	3.037	12.45	7.32	120.5	71.71	36.71	20.58	34.06						
MAX	9	9.22	4130	717	91.0	44.4	69.5	67	830	400	210	90	90						
MIN	8	7.2	3060	523	<.25	<.02	<.05	<.25	121	<20	32	13	3.6						
NO.OF OCCURS.	9	3	2	3	240	235	233	220	197	245	247	79	6						
NO: GT.																			
NO: LT.					2	01	70	3											

DATA RECORDED AS BEING LESS THAN LIMIT OF DETECTION - TAKEN AS 2/3 RECORDED VALUE

IF NO: OF LT'S EXCEEDS 30% TREAT DATA WITH CAUTION--IF MORE THAN 50% CONSULT STATISTICS DEPT.

END OF LISTING OF FILE :NHQTEHI.DAT(1,\*,1).JAMCAB(1) FOR USER :NHQTEHI AT 1995/04/07 12:17:54

\*\*\*\*\*

SPT :- 01093503LIV NGR :- SJ 32434 93168 LIVERPOOL STW PRIMARY TANK EFFLUENT TO RIVER

-----  
 TAKEN BEFORE SYPHON TO RIVER

FOR PERIOD 01/01/1990 00:01 TO 06/04/1995 17:15

REQUEST NO 1

WATER QUALITY SAMPLE SUMMARIES

PURP	76	81	82	61	77	172	102	111	116	117	180	92	85	135	143	182	38	47	
Temp	Diss	pH	Cond	Cl	Alkal	NH3	NO2	NO3	Ortho	COD	BOD	S	Solid	Sols	Silic	Flow	Flow	Flow	
PURP	Water	O2	at 25C	as	Cl	CaCO3	as	N	as	N	PO4	as	O	ATU	Susp	N/V	ate	Mean	Inst.
OF	Deg	C	%	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	M/d	M/d

MEAN	0.9	4140	477.6	105	12.01	.2194	.5768	2.943	274.2	107.7	87.06	19.62	170	167
S.D.	0.0	479.4	0.0	5.303	.3921	1.875	2.167	117.2	51.92	35.09	12.7	0.0	86.8	
MAX	0.9	4140	1030	165	30.6	2.32	24.2	24.4	752	290	355	60	170	583
MIN	0.7	4140	169	165	<.25	<.005	<.05	85	18	17	5	170	9.331	
NO. OF OCCURS.	1	1	3	1	178	177	175	176	103	174	179	24	1	156

NO. GT. 0  
 NO. LT. 5

DATA RECORDED AS BEING LESS THAN LIMIT OF DETECTION - TAKEN AS 2/3 RECORDED VALUE  
 IF NO: OF LT'S EXCEEDS 30% TREAT DATA WITH CAUTION--IF MORE THAN 50% CONSULT STATISTICS DEPT.  
 END OF LISTING OF FILE :NHQTEBI.DATA(1,\*,1).JAMCAB3(1) FOR USER :NHQTEBI AT 1995/04/J7\_\_12:16:51

WATER QUALITY SAMPLE SUMMARIES

SPT :- 010925101100104 NGR :- SJ 50084 85026 WARRINGTON SOUTH STW BELLHOUSE LANE ACTON GRANGE MOORE 259

FINAL EFFLUENT FINAL TANK OUTLET BESTEL DEAN SAMPLER

FOR PERIOD 01/01/1990 00:01 TO 04/04/1995 14:01

REQUEST NC 1

PURP	76	81	82	61	77	172	162	111	118	117	180	92	85	135	143	182	38	47
Temp	Diss	Diss	PH	Cond	CL	Alkal	NH3	NO2	NO3	Ortho	CO <sub>2</sub>	as O	ATU	Suspd	N/V	ate	Mean	Inst.
SAMPLE	PURP	Water	O <sub>2</sub>	O <sub>2</sub>	D 25C	as Cl	CaCO <sub>3</sub>	as N	as N	P04	as O	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	ML/d
OF	Deg	C %	mg/L	mg/L	US/cm	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	mg/L	ML/d	
MEAN	12.53	9.58	7.585	2457	18607	100	16.48	.4002	2.418	6.007	55.52	8.597	12.29	4.724	3.5	31	10.78	
S.D.	3.655	0.0	.2109	2962	37061	0.0	8.745	.429	3.172	2.925	33.66	7.624	10.79	6.803	0.0	0.0	4.057	
MAX	19.4	9.58	7.9	6900	74200	100	34	3.06	16.1	13	278	37	77	48	3.5	31	18.9	
MIN	8	9.58	7.4	907	72	100	.26	.01	<.1	.05	27	<2	<2	1	3.5	31	4.59	
NO.OF OCCURS.	13	1	4	4	4	1	118	118	118	117	99	133	133	52	1	1	13	
NO. GT.																		
NO. LT.																		

DATA RECORDED AS BEING LESS THAN LIMIT OF DETECTION - TAKEN AS 2/3 RECORDED VALUE  
 IF NO: OF LT'S EXCEEDS 30% TREAT DATA WITH CAUTION--IF MORE THAN 50% CONSULT STATISTICS DEPT.

SPT :- 01693S003HWD NGR :- SJ 48917 02364 HALEWOOD STW PRIMARY TANK EFFLUENT  
 -----  
 WATER QUALITY SAMPLE SUMMARIES  
 -----  
 SAMPLED BY EPIC SAMPLER FROM CHAMBER ADJ. NO6 SED TANK  
 FOR PERIOD 01/01/1990 00:01 TO 06/04/1995 17:12

REQUEST NO 1

PURP	76	01	02	01	77	172	102	111	116	117	180	92	85	135	143	182	38	47																								
	Temp		Diss		Conc		CL		ALCAL		NH3		NO2		NO3		Ortho		COJ		3CD		5		SOLID		SOLS		SILIC		FLOW											
SAMPLE	PURP		Water		O2		as		CL		CaCO3		as		N		as		N		P04-		as		O		ATU		Susod		N/V		ate		Mean		Inst.					
OF	Deg		C		%		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L		mg/L							
MEAN	8.375	8.162	2570	314.3	6.59	1.1	2.447	16.15	676.9	270.7	86.2	32.1	.04	43.09	0.0	347.3	0.04	3470	0.04	<1E-4	1	103	17	243	80	243	243	80	243	243	80	243	243	80	243	243	80					
S.D.	.5175	3559	288.4	145.4	38.76	1.32	4.235	13.53	322.7	192.7	36.78	17.06																														
MAX	9	8.55	2810	478	288	10.3	43.6	81.1	2730	2550	380	96																														
MIN	8	7.8	2250	>200	5.6	<.02	1.47	215	17.5	26	12																															
NO.OF OCCURS.	8	4	3	3	234	232	229	221	187	240	243	80																														
NO. GT.	1																																									
NO. LT.					55	85																																				

DATA RECORDED AS BEING LESS THAN LIMIT OF DETECTION - TAKEN AS 2/3 RECORDED VALUE  
 IF NO: OF LT'S EXCEEDS 30% TREAT DATA WITH CAUTION--IF MORE THAN 50% CONSULT STATISTICS DEPT.  
 END OF LISTING OF FILE :NHGTE6I.DAT(1,\*,1).JAMCA31(1) FOR USER :NHOTE6I AT 1995/04/07\_12:15:59

\*\*\*\*\*

SPT :- J17475C06J121 VGR :- SD 16341 70309 BARROW IN FURNESS NORTH SCALE STW EFFLUENT (SETTLEMENT ONLY)

\*\*\*\*\*  
OUTFALL TO WALNEY CHANNEL

FOR PERIOD 01/01/1990 TO 04/04/1995 14:01

REQUEST NO 1

WATER QUALITY SAMPLE SUMMARIES

PURP 76 81 82 81 77 172 102 111 118 117 180 92 85 135 143 182 38 47

Tempo Diss OH Cond Cl Alkal NH3 NO2 NO3 Ortho COD BOD 5 SOLID Sols Silic Flow Flcw

SAMPLE PURP Water O2 O2 mg/l OF Deg C % uS/cm mg/l ml/d ml/d

MEAN	7.573	7.45	50.94	26.5	.0748	.4246	6.125	347.8	149.4	105.5	34.13	.0741
S.D.	.4638	144.5	15.36	13.32	.1649	.7738	3.892	150.6	78.27	63.1	34.06	.2222
MAX	9.1	1198	157	48.5	1.11	4.26	22	916	405	325	176	<1
MIN	6.8	447	24	6.5	<.01	<.05	1.2	109	37	27	5	0.0

NO.OF OCCURS. 44 29 51 53 53 53 53 52 53 53 51 9

NO. GT. 2

NO. LT. 5 40 1 1

DATA RECORDED AS BEING LESS THAN LIMIT OF DETECTION - TAKEN AS 2/3 RECORDED VALUE  
IF NO. OF LT'S EXCEEDS 30% TREAT DATA WITH CAUTION--IF MORE THAN 50% CONSULT STATISTICS DEPT.

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 52

SAMPLE POINT - S03WITHAM DP WITHAM STM. FINAL EFFLUENT.  
SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	28 0	7.6439	0.1026	7.8511-	7.47	8.01	7.645
SS 105 C	0135 MG/L	48 3<	11.7208	>5.3179	21.9458 (LOG NORMAL)	4.0	27.5	11.25
BOD TOTAL	0088 MG/L O	47 12<	7.0887	>3.786	14.8924 (LOG NORMAL)	<3.0	18.34	7.0
BOD+ATU T	0085 MG/L O	1 1<	5.0	0.0	<5.0 (ONLY VALUE)	<5.0	<5.0	<5.0
AMMONIA N	0111 MG/L N	48 8<	0.8398	>1.6401	3.054 (LOG NORMAL)	<0.2	9.3017	0.3895
TON AS N	0116 MG/L N	28 0	14.4483	6.1579	26.0273 (LOG NORMAL)	3.82	28.7	13.1
P. SOL-REAC	0191 MG/L AS P	28 0	5.5279	2.1455	9.5431 (LOG NORMAL)	0.421	9.0018	5.325
CHLORIDE	0172 MG/L CL	48 0	14.4042	15.1595	170.2428 (LOG NORMAL)	92.8	161.0	147.0
CD TOTAL	9265 UG/L CD	15 9<	0.0513	>0.0192	0.1161 (LOG NORMAL)	<0.1	0.16	<0.1
1240H3CL3	7410 UG/L	1 1<	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
1230H6CL3	7411 UG/L	1 1<	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
INST. FLOW	9072 L/S	37 0	108.3513	26.2119	155.9016 (LOG NORMAL)	57.0	167.0	116.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 24

SAMPLE POINT - S03WITHAM DP WITHAM STM. FINAL EFFLUENT.  
SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	24 0	7.66	0.0886	7.8345-	7.49	7.86	7.63
SS 105 C	0135 MG/L	24 4<	8.3125	>4.8375	17.886 (LOG NORMAL)	<5.0	24.0	7.75
BOD TOTAL	0088 MG/L O	24 9<	5.6041	>4.9197	15.6171 (LOG NORMAL)	<2.6	26.2	6.05
AMMONIA N	0111 MG/L N	24 2<	0.4191	>0.3614	1.0924 (LOG NORMAL)	<0.2	1.75	0.2875
NITRITE N	0118 MG/L N	5 0	0.0386	0.0345	0.136 (MAX VALUE)	0.009	0.136	0.016
NITRATE N	0117 MG/L N	5 0	13.21	6.9267	24.6 (MAX VALUE)	7.74	24.6	10.9
TON AS N	0116 MG/L N	24 0	12.7554	5.3537	22.815 (LOG. NORMAL)	5.79	24.7	11.3
P. SOL-REAC	0191 MG/L AS P	24 0	5.047	1.1406	7.1066 (LOG NORMAL)	2.66	7.17	5.08
INST FLOW	9072 L/S	24 0	115.9156	23.3746	157.8072 (LOG NORMAL)	77.0	163.0	115.5

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 25

SAMPLE POINT - SO3WITHAM DP WITHIN STW. FINAL EFFLUENT.  
SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD.DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	22 0	7.6759	0.1237	7.9197-	7.5	8.0	7.7
SS 105 C	0135 MG/L	21 2<	7.5952-	>2.6571	12.8573(LLOG NORMAL)	4.4	14.5	<8.0
BOD TOTAL	0088 MG/L O	20 4X	18.6473-	>51.7993	71.0465(LLOG NORMAL)	<2.0	238.0	6.25
AMMONIA N	0111 MG/L N	21 1<	0.3904-	>0.3286	1.0014(LLOG NORMAL)	0.1	1.5	0.3
NITRITE N	0118 MG/L N	5 1<	0.154-	>0.1268	0.37 (MAX VALUE)	<0.05	0.37	0.13
NITRATE N	0117 MG/L N	5 0	9.94	4.0586	14.3 (MAX VALUE)	5.6	14.3	10.6
TON AS N	0116 MG/L N	22 0	13.65	5.5303	24.0232(LLOG NORMAL)	5.8	25.9	13.0
P SOL.REAC	0191 MG/L.AS P	22 0	6.28	1.8276	9.6381(LLOG NORMAL)	2.2	10.2	6.25
INST FLOW	9072.L/S	22 0	136.409	47.5461	224.8231(LLOG NORMAL)	47.0	310.0	123.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 99

SAMPLE POINT - SO6NORWICH0PO  
SAMPLE TYPE - DP

NORWICH (WHITLINGHAM) STW FINAL EFF COMB  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TG 27900 07800

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95XILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	72 0	7.4359	0.1408	7.7135- 7.1584	7.13	8.06	7.43
SS	105 C 0135 MG/L	98 29<	5.2013	>3.9562	13.4442(LOG NORMAL)	1.0	35.0	5.5
BOD+ATU	T 0085 MG/L O	98 86<	1.0022	>1.9415	8.5118(LOG NORMAL)	3.24	19.7	<9.105
AMMONIA	N 0111 MG/L N	98 46<	0.1844	>0.1477	0.5102(LOG NORMAL)	<0.2	1.3	0.2125
TON AS	N 0116 MG/L N	71 0	16.4438	4.7547	25.1812(LOG NORMAL)	7.54	33.7	16.3
P SOL-REAC	0191 MG/L AS P	72 0	4.032	0.83	5.5215(LOG NORMAL)	1.64	6.2	4.085
INST-FLOW	9072 L/S	84 0	747.5357	165.3459	1045.652 (LOG NORMAL)	510.0	1780.0	735.5

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 52

SAMPLE POINT - SO6NORWICH0PO  
SAMPLE TYPE - DP

NORWICH (WHITLINGHAM) STW FINAL EFF COMB  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TG 27900 07800

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95XILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	49 0	7.4734	0.1345	7.7384- 7.2084	7.19	7.8	7.45
SS	105 C 0135 MG/L	49 18<	5.0306	>2.5754	10.7951(LOG-NORMAL)	<5.0	14.0	5.5
BOD+ATU	T 0085 MG/L O	49 46<	0.471	>2.5692	8.7303(LOG-NORMAL)	<2.9	<10.1	<7.7
AMMONIA	N 0111 MG/L N	49 20<	0.4943	>1.6032	2.0534(LOG-NORMAL)	<0.2	11.4	0.219
NITRITE	N 0118 MG/L N	11 0	0.1795	0.2506	0.5784(LOG-NORMAL)	0.011	0.802	0.048
NITRATE	N 0117 MG/L N	11 0	17.8036	6.1709	29.2747(LOG-NORMAL)	8.14	30.8	16.3
TON AS	N 0116 MG/L N	49 0	17.9349	5.9088	28.883 (LOG-NORMAL)	8.19	31.8	17.4
P SOL-REAC	0191 MG/L AS P	49 0	3.6798	0.8589	5.2341(LOG-NORMAL)	1.37	5.39	3.81
INST-FLOW	9072 L/S	43 0	796.4814	258.8886	1275.736 (LOG NORMAL)	12.7	1620.0	772.0

DATE PRODUCED  
15/06/95

N.R.-A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

PAGE NO. 3

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 47

SAMPLE POINT - SO6NORMICHDP0  
SAMPLE TYPE - DP

NORWICH (WHITTINGHAM) STW FINAL EFF COMB  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TG 27900 07800

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	46 0	7.4265	0.1492	7.7205-	7.2	8.0	7.4
SS	0135 MG/L	46 12<	5.3826-	>4.9886	15.3622(L06 NORMAL)	1.5	26.0	5.55
BOD*ATU T	0085 MG/L O	46 17<	2.2282-	>1.5285	5.6334(L06 NORMAL)	1.4	9.2	<2.95
AMMONIA N	0111 MG/L N	46 39<	0.1065-	>0.5133	0.6491(L06 NORMAL)	<0.05	3.6	<0.15
NITRITE N	0118 MG/L N	10 7<	0.014.-	>0.0133	0.0539(L06 NORMAL)	<0.02	0.06	<0.05
NITRATE N	0117 MG/L N	11 0	15.3181	6.8822.	28.288 (L06 NORMAL)	0.2	24.9	16.3
TON AS N	0116 MG/L N	45 0	18.5733	5.3948	28.4847(L06 NORMAL)	8.5	29.9	18.4
P SOL-REAC	0191 MG/L AS P	46 0	4.576	0.8058	6.0076(L06 NORMAL)	2.9	6.4	4.65
INST FLOW	9072 L/S	33 0	789.9636	146.5441.	1051.1433(L06 NORMAL)	526.0	1310.0	765.7

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 32

GRID REF - TA 26300 10900

PYEPIPE PUMPING STATION CRUDE SEWAGE  
CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS)

SAMPLE POINT - S04PYEWIPEDB  
SAMPLE TYPE - DB

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
INST FLOW	0037 M3/S	28 0	0.93	1.6292	3.236 (LOG NORMAL)	0.152	9.17	0.6495
PH	0061 PH UNITS	30 0	6.95	0.3908	7.72 - 6.1799	6.07	7.72	6.955
SS	0135 MG/L	31 0	445.1935	235.7052	891.2534(LOG NORMAL)	136.0	1210.0	414.0
BOD+ATU	0085 MG/L	31 1<	599.6067-	>215.2123	1302.8499(LOG NORMAL)	<150.0	1140.0	601.5
AMMONIA N	0111 MG/L N	31 0	25.9523	7.7238	40.1663(LOG NORMAL)	8.0	49.7	24.146
TON AS N	0116 MG/L N	29 20<	0.4176-	>0.5939	1.6944(LOG NORMAL)	<0.6	3.01	<0.6
P SOL-REAC	0191 MG/L AS P	30 0	5.5954	1.8958	9.1146(LOG NORMAL)	2.06	9.48	5.1066
INST FLOW	9072 L/S	7 0	469.2665	282.3322.	866.0 (MAX VALUE)	0.866	866.0	458.0

DATE PRODUCED 15/06/95

N-R-A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

PAGE NO. 3

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 13

GRID REF - TA 26300 10900

PYEPIPE PUMPING STATION CRUDE SEWAGE  
CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS)

SAMPLE POINT - S04PYEWIPEDB  
SAMPLE TYPE - DB

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
INST FLOW	0037 M3/S	13 0	0.5756	0.2556	1.0572(LOG NORMAL)	0.178	0.993	0.536
PH	0061 PH UNITS	13 0	7.0607	0.452	7.9512- 6.1702	6.32	7.72	7.13
SS	0135 MG/L	13 0	323.2307	127.6839	562.4106(LOG NORMAL)	65.0	558.0	320.0
BOD+ATU	0085 MG/L	13 0	439.4384	172.3623	762.1871(LOG NORMAL)	171.0	793.0	414.0
AMMONIA N	0111 MG/L N	13 0	19.4923	9.6004	37.6378(LOG NORMAL)	5.04	33.9	19.2
NITRITE N	0118 MG/L N	6 0	0.0931	0.0335	0.118 (MAX VALUE)	0.0597	0.118	0.1055
NITRATE N	0117 MG/L N	6 4<	0.2021-	>0.0509	0.652 (MAX VALUE)	<0.5	0.652	<0.6
TON AS N	0116 MG/L N	13 9<	0.297 -	>0.3235	1.1153(LOG NORMAL)	<0.6	1.78	<0.6
P SOL-REAC	0191 MG/L AS P	13 0	3.56427	1.5643	6.5852(LOG NORMAL)	0.436	6.05	4.15
INST.FLOW	9072 L/S	5 0	529.4	248.5715	942.0 (MAX VALUE)	280.0	942.0	463.0

DATE PRODUCED  
15/06/95

N.R.A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

PAGE NO. 3

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 13

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

GRID REF - TA 26300 10900

SAMPLE POINT - S04PYEWIPEDB  
SAMPLE TYPE - DB

PYEWIPE PUMPING STATION (CRUDE SEWAGE  
CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS))

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
INST FLOW	M <sup>3</sup> /S	12	0-6796	0.2859	1-217 (LOG NORMAL)	0.33	1-223	0.637
PH	PH UNITS	13	7-0615	0.4958	8-0384-	5.9	7.9	7.1
SS	MG/L	13	287-2307	122.9201	518-4034 (LOG NORMAL)	114.0	542.0	260.0
BOD+ATU T	MG/L	12	424-9181*	>257.5443	911-9136 (LOG NORMAL)	>67.4	1030.0	377.0
AMMONIA-N	MG/L	13	24-8538	10.5486	44-6849 (LOG NORMAL)	3.7	36.6	26.3
NITRITE-N	MG/L	6	0-2236-	>0.454	1-17 (MAX VALUE)	<0.05	1-17	0.06
NITRATE-N	MG/L	6	0-303.-	>0.496	1-53 (MAX VALUE)	<0.1	1-53	<0.6
TOTAL-N	MG/L	13	0-2 --	>0.633	1-3394 (LOG NORMAL)	<0.2	2.6	<0.6
P SOL-REAC	MG/L AS P	13	5-3976	2.45	10-0167 (LOG NORMAL)	0.8	8.3	6.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 58

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

GRID REF - TL 22260 98160

SAMPLE POINT - S07FLAGFENDP FLAG FEN STW COMBINED FINAL EFFLUENT  
SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD.DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	48	7.6229	0.1248	7.8688- 7.377	7.16	7.84	7.64
SS	0135 MG/L	57	20.3473	>12.9786	44.9875(LOG NORMAL)	<5.0	78.5	16.0
BOD+ATU	0095 MG/L O	57	11.765	>11.4462	34.0127(LOG NORMAL)	<5.0	71.8	10.5
AMMONIA	0111 MG/L N	57	22.5414	7.4573	36.3623(LOG NORMAL)	10.1	61.6	23.134
TON AS N	0116 MG/L N	45	1.4416	>6.2746	6.2196(LOG NORMAL)	<0.6	42.8	<0.6
P SOL.REAC	0191 MG/L AS P	48	3.9891	1.907	7.5908(LOG NORMAL)	1.31	11.7	3.997
INST FLOW	9072 L/S	37	833.8648	386.0525	1562.1963(LOG NORMAL)	368.0	2000.0	662.0

DATE PRODUCED  
15/06/95

N.R.A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

PAGE NO. 1

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 49

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

GRID REF - TL 22260 98160

SAMPLE POINT - S07FLAGFENDP FLAG FEN STW COMBINED FINAL EFFLUENT  
SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD.DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	48	7.616	0.0726	7.7591- 7.4729	7.38	7.77	7.63
SS	0135 MG/L	48	23.927	7.8435	38.4551(LOG NORMAL)	9.5	44.0	24.25
BOD+ATU	0095 MG/L O	48	17.8583	>6.9631	30.9675(LOG NORMAL)	<7.5	38.7	18.4
AMMONIA	0111 MG/L N	48	22.077	4.8254	30.7703(LOG NORMAL)	10.8	28.9	22.45
NITRITE	0118 MG/L N	10	0.1025	0.2897	0.3913(LOG NORMAL)	0.0073	0.927	0.011
NITRATE	0117 MG/L N	10	0.598	>0.572	1.7836(LOG NORMAL)	<0.5	2.33	<0.618
TON AS N	0116 MG/L N	48	0.4815	>0.4087	1.4026(LOG NORMAL)	<0.6	3.26	<0.622
P SOL.REAC	0191 MG/L AS P	48	4.2897	1.3086	6.7021(LOG NORMAL)	1.69	7.69	4.215
INST FLOW	9072 L/S	41	703.8536	152.6192	978.6547(LOG NORMAL)	335.0	1100.0	684.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 50

SAMPLE POINT - S07FLAGFENDP  
SAMPLE TYPE - DP

FLAG FEN STM COMBINED FINAL EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TL 22260 98160

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD.DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH	47	7.5336	0.1705	7.8696-	7.0	8.11	7.5
SS 105 C	0135 MG/L	47	26.8063-	>11.0347	47.7867(LOG NORMAL)	<1.0	52.0	27.0
BOD*ATU T	0085 MG/L	45	16.2295-	>7.3092	30.7881(LOG NORMAL)	2.9	>40.2	17.6
AMMONIA N	0111 MG/L	47	20.8869-	>5.6767	31.2613(LOG NORMAL)	<0.2	30.1	20.8
NITRITE N	0118 MG/L	11	0.0081-	>0.0177	0.0524(LOG NORMAL)	0.01	0.05	0.02
NITRATE N	0117 MG/L	11	0.4727-	>1.4783	2.3373(LOG NORMAL)	<0.1	5.2	<0.5
TON AS N	0116 MG/L	46	0.2695-	>1.2764	1.836 (LOG NORMAL)	<0.1	8.9	<0.6
P SOL-REAC	0191 MG/L AS P	47	4.2208-	>1.5853	7.1852(LOG NORMAL)	<0.3	9.6	4.2
INST FLOW	9072 L/S	37	818.7567	228.1502	1236.714 (LOG NORMAL)	391.0	1490.0	788.0

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 54

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

GRID REF - TF 45900 14900

WISBECH STW COMBINED EFFLUENT FROM 1/8/85  
FINAL SEDIMENTATION TANK EFFLUENT

SAMPLE POINT - S07WISBECHDP  
SAMPLE TYPE - DP

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD. DEV.	95XILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	44	7.8081	0.1774	8.1577 - 7.4585	7.22	8.12	7.85
SS 105 C	0135 MG/L	50	30.746	26.0321	78.6354 (LOG NORMAL)	9.0	133.0	22.75
BOD+ATU T	0085 MG/L O	50	10.3689	>25.5197	59.4252 (LOG NORMAL)	<12.0	>151.0	<20.0
AMMONIA N	0111 MG/L N	50	1.1808	26.7356	3.9675 (LOG NORMAL)	<0.2	9.13	0.5253
TON AS N	0116 MG/L N	43	17.2476	11.3062	38.0261 (LOG NORMAL)	0.759	44.1	16.6
P SOL+REAC	0191 MG/L AS P	44	7.7934	2.0023	11.4409 (LOG NORMAL)	4.09	12.8	7.3681
CHLORIDE	0172 MG/L CL	50	204.6056	34.0337	264.858 (LOG NORMAL)	108.0	315.0	206.56
CD TOTAL	9265 UG/L CD	4	0.0525	>0.055	0.21 (MAX VALUE)	<0.1	0.21	<0.1
HG TOTAL	9269 UG/L HG	12	0.005	>0.0028	0.0329 (LOG NORMAL)	<0.05	0.06	<0.05
1246H3CL3	7410 UG/L	1	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
123C6H6CL3	7411 UG/L	1	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
INST FLOW	9072 L/S	41	206.0734	151.3172	489.2169 (LOG NORMAL)	18.0	672.0	184.0

N.R.A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

DATE PRODUCED  
15/06/95

PAGE NO. 3

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 25

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

GRID REF - TF 45900 14900

WISBECH STW COMBINED EFFLUENT FROM 1/8/85  
FINAL SEDIMENTATION TANK EFFLUENT

SAMPLE POINT - S07WISBECHDP  
SAMPLE TYPE - DP

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD. DEV.	95XILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	24	7.8129	0.1284	8.066 - 7.5597	7.53	8.02	7.805
SS 105 C	0135 MG/L	24	53.3333	155.9816	204.2291 (LOG NORMAL)	<5.0	784.0	22.5
BOD+ATU T	0085 MG/L O	24	11.2291	>40.0554	62.578 (LOG NORMAL)	<5.2	208.0	<10.85
AMMONIA N	0111 MG/L N	24	0.7385	>0.7466	2.0768 (LOG NORMAL)	<0.2	2.99	0.4465
NITRITE N	0118 MG/L N	3	0.3063	0.027	0.533 (MAX VALUE)	0.279	0.333	0.307
NITRATE N	0117 MG/L N	3	33.6656	16.778	51.3 (MAX VALUE)	17.9	51.3	31.8
TON AS N	0116 MG/L N	24	28.51	11.8845	50.8335 (LOG NORMAL)	2.34	51.6	31.3
P SOL+REAC	0191 MG/L AS P	24	8.27	2.2133	12.3133 (LOG NORMAL)	2.86	12.7	8.52
CHLORIDE	0172 MG/L CL	24	250.3353	75.0038	390.3903 (LOG NORMAL)	150.0	535.0	238.0
CD TOTAL	9265 UG/L CD	5	0.0	>0.0	<0.1 (MAX VALUE)	<0.1	<0.1	<0.1
HG TOTAL	9269 UG/L HG	6	0.0	>0.0	<0.05 (MAX VALUE)	<0.05	<0.05	<0.05
TECHNAZINE	7366 MG/L	1	1	0.0	<25.0 (ONLY VALUE)	<25.0	0.0	0.0
MASS SPEC	7422	1	0.0	0.0	0.0 (ONLY VALUE)	0.0	0.0	0.0
C6H3CL4N	7472 NG/L	1	1	0.0	<25.0 (ONLY VALUE)	<25.0	<25.0	<25.0
C7H4CL4S	7473 NG/L	1	1	0.0	<25.0 (ONLY VALUE)	<25.0	<25.0	<25.0
CL-PROPAM	7528 NG/L	1	0	0.0	441.0 (ONLY VALUE)	441.0	441.0	441.0
INST FLOW	9072 L/S	22	156.7727	62.4255	273.759 (LOG NORMAL)	36.0	308.0	141.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 24

SAMPLE POINT - S07WISBECHDP  
SAMPLE TYPE - DP

WISBECH STW COMBINED EFFLUENT FROM 1/8/85  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TF 45900 14900

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD. DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	23 0	7.806	0.1878	8.1761-	7.5	8.31	7.8
SS 105 C	0135 MG/L	23 2<	22.5608-	>14.1637	49.7894(LOG NORMAL)	7.0	60.5	20.0
BOD+ATU T	0085 MG/L O	21 6<	8.5428	>4.1174	15.4076(LOG NORMAL)	<2.0	17.2	8.0
AMMONIA N	0111 MG/L N	22 13X	0.1619-	>1.1545	2.4427(LOG NORMAL)	<0.1	>20.0	<0.2
NITRITE N	0118 MG/L N	3 1<	0.1333-	>0.0871	0.21 (MAX VALUE)	<0.05	0.21	0.19
NITRATE N	0117 MG/L N	3 1<	21.2	>18.5971	35.1 (MAX VALUE)	<0.1	35.1	28.5
TON AS N	0116 MG/L N	23 2<	24.1782-	>13.5296	49.8053(LOG NORMAL)	<0.2	40.0	28.7
P SOL.-REAC	0191 MG/L AS P	23 0	9.0426	2.7654	14.1411(LOG NORMAL)	4.28	13.6	9.3
CHLORIDE	0172 MG/L CL	22 0	263.7636	49.0102	351.1217(LOG NORMAL)	134.0	322.0	274.0
CD TOTAL	9265 UG/L CD	13 7<	0.0823-	>0.1187	0.3631(LOG NORMAL)	<0.1	<0.5	<0.2
HG TOTAL	9269 UG/L HG	3 2<	0.0183-	>0.0028	0.055 (MAX VALUE)	<0.05	0.055	<0.05
INST FLOW	9072 L/S	20 0	142.9	50.4859	236.8597 (LOG NORMAL)	20.0	234.0	130.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY  
 REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT  
 REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061, PH UNITS	37 0	6.6281	0.3549	7.3273-5.9288	5.99	7.59	6.57
SS 105 C	0135 MG/L	45 0	156.7593	227.1643	532.1164(LOG NORMAL)	50.5	1640.0	128.0
BOD+ATU T	0085 MG/L O	45 1>	440.482 *	>199.3103	816.1914(LOG NORMAL)	30.3	1310.0	446.47
AMMONIA N	0111 MG/L N	45 0	30.8359	9.7404	48.8333(LOG NORMAL)	9.56	57.6	29.7
TON AS N	0116 MG/L N	35 23<	0.8021-	>1.8774	3.5456(LOG NORMAL)	<0.6	8.93	<0.6
P SOL.REAC	0191 MG/L AS P	37 0	7.7044	2.545	12.4206(LOG NORMAL)	1.68	14.0	7.8561
CHLORIDE	0172 MG/L CL	45 0	508.4382	247.0309	975.1811(LOG NORMAL)	168.0	1034.1	445.06
CD TOTAL	9265 UG/L CD	19 0	0.2294	0.1008	0.4193(LOG NORMAL)	0.11	0.45	0.19
SILVER TOT	9198 UG/L	19 3<	2.2524-	>2.1839	6.2244(LOG NORMAL)	<0.5	8.95	1.2
ST 40 LC50	7070 X CONC.	1 0	11.0	0.0	11.0 (ONLY VALUE)	11.0	11.0	11.0
1240GH3CL3	7410 UG/L	1 1<	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
1230GH6CL3	7411 UG/L	1 1<	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
INST FLOW	9071 M3/HR	2 0	318.0	62.2254	362.0 (MAX VALUE)	274.0	362.0	318.0
INST FLOW	9072 L/S	14 0	250.9371	126.0055	489.1999(LOG NORMAL)	121.58	442.75	203.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY  
 REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT  
 REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	38 0	6.5913	0.2946	7.1717-6.0109	6.0	7.36	6.56
SS 105 C	0135 MG/L	38 0	147.7105	49.902	240.3215(LOG NORMAL)	84.0	336.0	133.0
BOD+ATU T	0085 MG/L O	38 2>	482.4 *	>179.6892	817.8493(LOG NORMAL)	32.4	982.0	476.0
AMMONIA N	0111 MG/L N	38 0	31.2239	16.5512	62.5462(LOG NORMAL)	6.51	105.0	30.35
NITRATE N	0118 MG/L N	11 0	0.1274	0.0502	0.2216(LOG NORMAL)	0.02	0.195	0.125
NITRITE N	0117 MG/L N	11 4<	1.465 -	>2.6275	5.3439(LOG NORMAL)	<0.2	9.09	<0.6
TON AS N	0116 MG/L N	38 17<	0.8489-	>1.5105	3.2687(LOG NORMAL)	<0.2	9.29	0.623
P SOL.REAC	0191 MG/L AS P	38 1<	6.9007-	>3.8077	14.1123(LOG NORMAL)	<0.3	20.4	6.68
CHLORIDE	0172 MG/L CL	38 1<	481.7368-	>306.1648	1059.8734(LOG NORMAL)	<3.3	1430.0	398.5
CD TOTAL	9265 UG/L CD	26 4<	0.1731-	>0.2402	0.6545(LOG NORMAL)	0.11	<1.0	0.185
HG TOTAL	9269 UG/L HG	8 5<	0.0498-	>0.0813	0.282 (MAX VALUE)	<0.05	0.282	<0.05
AS TOTAL	9261 UG/L AS	8 1<	2.3212-	>1.3333	4.7 (MAX VALUE)	<0.5	4.7	2.155
SILVER TOT	9198 UG/L	26 15<	0.6393-	>5.1539	6.2752(LOG NORMAL)	<0.5	>99999.0	<1.0
E COLI P	2549 NO/100ML	1 1>	99999.0	0.0	>99999.0 (ONLY VALUE)	>99999.0	>99999.0	<99999.0
ST 40 LC50	7070 X CONC.	1 1<	8.0	0.0	<8.0 (ONLY VALUE)	<8.0	<8.0	<8.0
S FAEC P	7092 NO/ML	4 2>	31300.0 *	>16381.5952	>50300.0 (MAX VALUE)	>10000.0	>50000.0	31300.0
INST FLOW	9072 L/S	11 0	295.1818	168.8086	614.676 (LOG NORMAL)	86.0	562.0	290.0

DATE PRODUCED  
15/06/95

N.R.A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

PAGE NO. 1

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 39

SAMPLE POINT - SOZKINGLYNDP  
SAMPLE TYPE - DP

KING'S LYNN STW F/E  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	36	6.543	0.432	7.3942-	5.6	7.3	6.55
SS	105 C	37	127.3297	42.5286	206.1988(LOG NORMAL)	39.2	246.0	126.0
BOD+ATU	T 0085 MG/L	35	471.1269*	>254.4349	952.7076(LOG NORMAL)	30.3	1210.0	<417.0
AMMONIA	N 0111 MG/L	33	27.4983*	>8.2118	42.6137(LOG NORMAL)	0.5	40.6	28.7
NITRITE	N 0118 MG/L	11	0.1018	0.0244	0.146 (LOG NORMAL)	0.07	0.14	0.1
NITRATE	N 0117 MG/L	11	0.0	>0.2085	0.5281(LOG NORMAL)	<0.05	<0.5	<0.5
TON AS	N 0116 MG/L	34	0.4181-	>1.3678	2.2241(LOG NORMAL)	0.016	8.0	<0.6
P SOL.REAC	0191 MG/L AS P	34	7.306	3.235	13.399 (LOG NORMAL)	0.297	20.1	7.1

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 54

SAMPLE POINT - S07SPALDNGDP  
SAMPLE TYPE - DP

SPALDING STW F/E  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TF 26100 25200

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	41 0	7.8295	0.132	8.0896-	7.48	8.12	7.84
SS 105 C	0135 MG/L	51 0	32.7941	19.0439	68.8319(LOG NORMAL)	10.0	117.0	29.0
D.O	0082 MG/L	1 0	8.6	0.0	8.6 (ONLY VALUE)	8.6	8.6	8.6
BOD+ATU T	3085 MG/L	51 16<	22.0484-	>16.8127	55.9849(LOG NORMAL)	11.2	103.0	22.63
AMMONIA N	0111 MG/L	51 0	12.3625	5.0354	21.8101(LOG NORMAL)	3.59	23.701	12.6
TON AS N	0116 MG/L	40 0	10.7656	5.1248	20.4639(LOG NORMAL)	2.02	25.426	9.5312
P SOL-REAC	0191 MG/L AS P	41 0	2.9837	1.0905	5.0175(LOG NORMAL)	0.972	4.7537	3.2
CHLORIDE	0172 MG/L CL	51 0	192.2703	37.9948	260.2629(LOG NORMAL)	113.0	302.9	190.41
HG TOTAL	9269 UG/L HG	4 1<	0.0417-	>0.0085	0.067 (MAX VALUE)	<0.05	0.067	0.05
DICHLORVOS	0507 UG/L	12 12<	0.0	>0.0277	0.0753(LOG NORMAL)	<0.025	<0.1	<0.045
MALATHION	0535 UG/L	12 12<	0.0	>0.0529	0.0753(LOG NORMAL)	<0.025	<0.1	<0.045
PARATHION	0543 UG/L	12 12<	0.0	>0.0277	0.0736(LOG NORMAL)	<0.03	<0.1	<0.045
124C6H3CL3	7410 UG/L	1 1<	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
123C6H6CL3	7411 UG/L	1 1<	0.1	0.0	<0.1 (ONLY VALUE)	<0.1	<0.1	<0.1
AZINPH-ETH	7440 UG/L	12 12<	0.0	>0.0591	0.0719(LOG NORMAL)	<0.03	<0.1	<0.05
FENTHION	7441 UG/L	12 12<	0.0	>0.0529	0.0753(LOG NORMAL)	<0.025	<0.1	<0.045
PARATH-MET	7442 UG/L	12 12<	0.0	>0.0529	0.0753(LOG NORMAL)	<0.025	<0.1	<0.045
INST FLOW	9072 L/S	45 0	80.8644	35.5228	147.7475(LOG NORMAL)	14.5	150.0	73.8

DATE PRODUCED 15/06/95 N.R.A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

PAGE NO. 2

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 24

SAMPLE POINT - S07SPALDNGDP  
SAMPLE TYPE - DP

SPALDING STW F/E  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TF 26100 25200

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	24 0	7.8912	0.0903	8.0692-	7.67	8.14	7.88
SS 105 C	0135 MG/L	24 0	27.5	8.9721	44.1133(LOG NORMAL)	7.5	49.0	27.5
BOD+ATU T	3085 MG/L	24 7<	15.7541-	>7.9017	32.5411(LOG NORMAL)	<9.7	41.4	18.15
AMMONIA N	0111 MG/L	24 0	7.634	4.9673	17.004 (LOG NORMAL)	0.508	17.0	8.03
NITRATE N	0118 MG/L	4 0	1.1012	0.4517	1.58 (MAX VALUE)	0.667	1.58	1.079
NITRITE N	0117 MG/L	4 0	11.425	4.4591	17.8 (MAX VALUE)	7.44	17.8	10.23
TON AS N	0116 MG/L	24 0	14.5145	5.1749	24.1511(LOG NORMAL)	4.96	22.1	15.45
P SOL-REAC	0191 MG/L AS P	24 0	3.605	0.8689	5.1809(LOG NORMAL)	1.96	5.15	3.81
CHLORIDE	0172 MG/L CL	24 0	176.375	18.5983	208.5241(LOG NORMAL)	118.0	214.0	178.0
HG TOTAL	9269 UG/L HG	4 2<	0.0355-	>0.0163	0.083 (MAX VALUE)	<0.05	0.083	<0.0505
DICHLORVOS	0507 UG/L	5 5<	0.0	>0.0164	<0.05 (MAX VALUE)	<0.02	<0.05	<0.02
MALATHION	0535 UG/L	5 5<	0.0	>0.0054	<0.05 (MAX VALUE)	<0.04	<0.05	<0.04
PARATHION	0543 UG/L	5 5<	0.0	>0.0054	<0.05 (MAX VALUE)	<0.04	<0.05	<0.04
AZINPH-ETH	7440 UG/L	5 5<	0.0	>0.0054	<0.05 (MAX VALUE)	<0.04	<0.05	<0.04
FENTHION	7441 UG/L	5 5<	0.0	>0.0054	<0.05 (MAX VALUE)	<0.04	<0.05	<0.04
PARATH-MET	7442 UG/L	5 5<	0.0	>0.0054	<0.05 (MAX VALUE)	<0.04	<0.05	<0.04
INST FLOW	9072 L/S	24 0	94.075	45.5931	180.2132(LOG NORMAL)	20.7	182.0	92.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES -- 23

SAMPLE POINT - S07SPALDNGDP  
SAMPLE TYPE - DP

SPALDING STW F/E  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TF 26100 25200

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD-DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	22 0	7.8577	0.1432	8.14 - 7.5754	7.4	8.0	7.9
SS 105 C	0135 MG/L	22 1<	24.3954	>13.7577	50.5502(LOG NORMAL)	<5.0	73.0	21.6
BOD+ATU T	0085 MG/L 0	22 3X	17.445	>11.9845	40.6245(LOG NORMAL)	<2.0	60.4	16.35
AMMONIA N	0111 MG/L N	22 0	9.0727	6.9782	22.0707(LOG NORMAL)	0.3	27.4	7.35
NITRITE N	0118 MG/L N	3 1>	1.19 *	>0.02	1.2 (MAX VALUE)	>1.16	1.2	1.18
NITRATE N	0117 MG/L N	2 0	17.3	8.7681	23.5 (MAX VALUE)	11.1	23.5	17.3
TON AS N	0116 MG/L N	22 0	13.009	4.2475	20.8745(LOG NORMAL)	6.9	23.7	12.4
P SOL-REAC	0191 MG/L AS P	22 0	3.7636	1.0513	5.6899(LOG NORMAL)	2.4	6.0	3.6
CHLORIDE	0172 MG/L CL	22 0	192.5454	27.9348	241.6089(LOG NORMAL)	125.0	270.0	196.0
HG TOTAL	9269 UG/L HG	4 3<	0.0075	>0.0191	<0.05 (MAX VALUE)	<0.01	<0.05	<0.04
DICHLORVOS	0507 UG/L	4 4<	0.0	>0.004	<0.028 (MAX VALUE)	<0.02	<0.028	<0.02
MALATHION	0535 UG/L	3 3<	0.0	>0.0	<0.02 (MAX VALUE)	<0.02	<0.02	<0.02
PARATHION	0543 UG/L	3 3<	0.0	>0.0	<0.02 (MAX VALUE)	<0.02	<0.02	<0.02
AZINPH-ETH	7440 UG/L	3 3<	0.0	>0.0	<0.02 (MAX VALUE)	<0.02	<0.02	<0.02
FENTHION	7441 UG/L	2 2<	0.0	>0.0	<0.02 (MAX VALUE)	<0.02	<0.02	<0.02
PARATH-MET	7442 UG/L	4 4<	0.0	>0.0025	<0.02 (MAX VALUE)	<0.015	<0.02	<0.02
INST FLOW	9072 L/S	21, 0	121.9456	63.076	241.2798(LOG NORMAL)	35.6	277.0	117.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 42

SAMPLE POINT - S04BOSTON DP  
SAMPLE TYPE - DP

BOSTON HUMUS TANK EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TF 35600 40900

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95XILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH	35	8.0228	0.1124	8.2443-7.8014	7.84	8.29	8.01
SS	105 C	41	23.3414	>12.9273	47.8724(LOG NORMAL)	<5.0	66.0	20.0
BOD5	0085 MG/L	41	6.6209	>2.5316	13.1606(LOG NORMAL)	7.45	17.51	<10.0
AMMONIA	N 0111 MG/L	41	1.2784	1.1609	3.3889(LOG. NORMAL)	0.221	5.8674	0.868
TON AS	N 0116 MG/L	34	27.6447	9.0705	44.4465(LOG NORMAL)	10.4	46.9	26.2705
P SOL.REAC	0191 MG/L	35	5.0279	0.8909	6.6113(LOG NORMAL)	3.18	7.8268	4.82
INST FLOW	9072 L/S	5	172.44	83.3205	259.0 (MAX VALUE)	77.4	259.0	183.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 12

SAMPLE POINT - S04BOSTON DP  
SAMPLE TYPE - DP

BOSTON HUMUS TANK EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF - TF 35600 40900

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95XILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH	12	8.0	0.1133	8.2232-7.7768	7.84	8.22	7.98
SS	105 C	12	30.4156	15.2029	59.1609(LOG NORMAL)	11.5	58.5	29.5
BOD5	0085 MG/L	12	11.6583	>5.1926	22.0515(LOG NORMAL)	5.5	21.6	12.7
AMMONIA	N 0111 MG/L	12	1.447	1.537	4.143 (LOG NORMAL)	0.264	4.74	0.64
NITRITE	N 0118 MG/L	10	0.1716	0.104	0.3683(LOG NORMAL)	0.053	0.333	0.154
NITRATE	N 0117 MG/L	10	24.88	7.7918	39.2686(LOG NORMAL)	15.2	36.9	24.0
TON AS	N 0116 MG/L	12	25.0916	7.0666	38.0468(LOG NORMAL)	15.4	37.0	25.3
P SOL.REAC	0191 MG/L	12	4.955	1.2661	7.1827(LOG NORMAL)	2.54	6.59	5.21
INST FLOW	9072 L/S	24	134.8996	98.8314	319.8694(LOG NORMAL)	1.175	302.0	121.25

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 12

GRID REF - TF 35600 40900

SAMPLE POINT - SO4BOSTON DP BOSTON HUMUS TANK EFFLUENT  
 SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(O.R RANGE)	STD. DEV.	95%ILE(O.R RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	12 0	7.8566	0.2055	8.2616-	7.4	8.2	7.875
SS	0135 MG/L	12 0	21.3156	11.7475	43.5533(LLOG NORMAL)	8.4	46.0	18.85
BOD+ATU	T 3085 MG/L O	12 0	10.9166	5.9686	22.2144(LLOG NORMAL)	2.6	23.2	9.35
AMMONIA	N 0111 MG/L N	12 0	1.4	0.8984	3.0958(LLOG NORMAL)	0.1	2.7	1.05
NITRITE	N 0118 MG/L N	11 0	0.4545	0.155	0.7424(LLOG NORMAL)	0.2	0.74	0.4
NITRATE	N 0117 MG/L N	11 0	19.4454	9.0574	36.5368(LLOG NORMAL)	7.4	32.7	17.9
TON AS	N 0115 MG/L N	12 0	20.4916	9.2315	37.8905(LLOG NORMAL)	7.8	32.7	19.0
P SOL.REAC	0191 MG/L AS P	12 0	5.4583	1.2471	7.7118(LLOG NORMAL)	2.2	7.4	5.45

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 13

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

GRID REF - TA 32100 07100

CLEETHORPES SOUTHERN OUTFALL CRUDE SEW CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS)

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
INST FLOW	0037 M <sup>3</sup> /S	4	0.33	0.0739	0.44 (MAX VALUE)	0.28	0.44	0.3
PH	0061 PH UNITS	13	7.693	0.2962	8.2767- 7.1093	7.38	8.19	7.65
SS	105 C	13	323.7692	58.592	428-0323(LOG NORMAL)	220.0	440.0	330.0
BOD+ATU	T 0085 MG/L	13	322.0807	53.71	417-1774(LOG NORMAL)	221.0	385-26	344.0
AMMONIA	N 0111 MG/L	13	37.9116	10.412	56.9666(LOG NORMAL)	26.2	62.875	36.64
TON AS N	0116 MG/L	13	0.1611	>0.056	0.4902(LOG NORMAL)	<0.6	0.779	<0.6
P SOL.REAC	0191 MG/L AS P	13	7.5048	2.0629	11.2805(LOG NORMAL)	5.23	12.4	7.44
M D FLOW	9073 M <sup>3</sup> /D	13	15198.5384	2916.9108	20408.5366(LOG NORMAL)	11800.0	20300.0	14400.0

DATE PRODUCED 15/06/95

N.R.A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 13

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

GRID REF - TA 32100 07100

CLEETHORPES SOUTHERN OUTFALL CRUDE SEW CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS)

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
INST FLOW	0037 M <sup>3</sup> /S	1	2.09	0.0	2.09 (ONLY VALUE)	2.09	2.09	2.09
PH	0061 PH UNITS	13	7.7023	0.1933	8.0832- 7.3214	7.24	8.01	7.72
SS	105 C	13	318.5384	160.2537	621-5727(LOG NORMAL)	93.0	752.0	306.0
BOD+ATU	T 0085 MG/L	13	286.0769	104.1517	480-2699(LOG NORMAL)	126.0	454.0	282.0
AMMONIA	N 0111 MG/L	13	27.8407	9.7872	46.0502(LOG NORMAL)	5.73	40.1	30.1
NITRITE	N 0118 MG/L	6	0.0977	0.0545	0.155 (MAX VALUE)	0.0334	0.155	0.099
NITRATE	N 0117 MG/L	6	0.3156	>0.297	1.3 (MAX VALUE)	<0.5	1.3	<0.6
TON AS N	0116 MG/L	13	1.3554	>2.0132	4.7467(LOG NORMAL)	<0.6	7.56	0.627
P SOL.REAC	0191 MG/L AS P	13	5.353	2.2584	9.5976(LOG NORMAL)	1.49	8.84	5.14
M D FLOW	9073 M <sup>3</sup> /D	13	15513.0769	4488.9375	23757.9786(LOG NORMAL)	9670.0	26900.0	14200.0

PAGE NO. 2

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 12

SAMPLE POINT - S04CLEETHPDB CLEETHORPES SOUTHERN OUTFALL CRUDE SEW  
 SAMPLE TYPE - DB CRUDE SEWAGE(AJ SEWAGE TREATMENT WORKS)

GRID REF - TA 32100 07100

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD. DEV.	95XILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	7 0	7.7857	0.3387	7.4 - 8.5	.74	8.5	7.7
SS 105 C	0135 MG/L	7 0	293.2857	166.2886	650.0 (MAX VALUE)	118.0	650.0	259.0
BOD*ATU T	0085 MG/L.0	7 1>	311.6656*	>157.9059	534.0 (MAX VALUE)	65.0	534.0	287.0
AMMONIA N	0111 MG/L N	7 0	36.2	15.347	67.4 (MAX VALUE)	24.2	67.4	37.6
NITRITE N	0118 MG/L N	4 3<	0.015	>0.005	0.06 (MAX VALUE)	<0.05	0.06	<0.05
NITRATE N	0117 MG/L N	4 3<	0.05	>0.15	<0.5 (MAX VALUE)	0.2	<0.5	<0.5
TON AS N	0116 MG/L N	7 6<	0.0857	>0.1496	0.6 (MAX VALUE)	<0.2	0.6	<0.6
P SOL.REAC	0191 MG/L AS P	7 0	7.3671	2.3867	11.5 (MAX VALUE)	4.5	11.5	7.9
M. D FLOW	9073 M3/D	8 0	1.0912.5	6721.9764	19900.0 (MAX VALUE)	190.0	19900.0	11405.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 45

GRID REF - TM 17200 41900

CLIFF QUAY FINAL EFFLUENT  
PRIMARY SEDIMENTATION TANK EFFLUENT

SAMPLE POINT - S06CLIFFQUDJ0  
SAMPLE TYPE - DJ

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD. DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	36 0	7.3397	0.2372	7.8071-	6.85	7.86	7.33
SS	105 C 0135 MG/L	41 0	202.9268	71.5812	336.1342(LOG NORMAL)	38.0	448.0	206.0
BOD+ATU	T 0085 MG/L	41 1<	324.5339-	>113.7765	537.3529(LOG NORMAL)	96.5	623.79	318.0
AMMONIA	N 0111 MG/L	41 0	35.5302	10.3085	54.4676(LOG NORMAL)	7.74	50.9	39.4
TOTAL	AS N 0116 MG/L	35 27<	0.2118-	>0.2015	0.8231(LOG NORMAL)	<0.6	1.5673	<0.6
P SOL	REAC 0191 MG/L	AS P 36 0	8.7978	2.717-	13.8104(LOG NORMAL)	2.33	14.0	8.7573

ATE PRODUCED 15/06/95

N.R.-A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 24

GRID REF - TM 17200 41900

CLIFF QUAY FINAL EFFLUENT  
PRIMARY SEDIMENTATION TANK EFFLUENT

SAMPLE POINT - S06CLIFFQUDJ0  
SAMPLE TYPE - DJ

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD. DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	24 0	7.3616	0.2423	7.839 -	6.95	7.85	7.38
SS	105 C 0135 MG/L	24 0	154.6791	70.8357	288.2673(LOG NORMAL)	23.0	284.0	144.0
BOD+ATU	T 0085 MG/L	24 2X	252.4652-	>109.2794	465.4071(LOG NORMAL)	62.7	>426.3	258.5
AMMONIA	N 0111 MG/L	24 0	35.327	11.15	55.9278(LOG NORMAL)	9.35	57.3	38.25
NITRITE	N 0118 MG/L	11 0	0.0992	0.0859	0.2568(LOG NORMAL)	0.012	0.258	0.078
NITRATE	N 0117 MG/L	11 8<	0.1663-	>0.0752	0.5057(LOG NORMAL)	<0.5	0.753	<0.6
TOTAL	AS N 0116 MG/L	24 15<	0.2805-	>0.1041	0.6558(LOG NORMAL)	<0.6	1.06	<0.6
P SOL	REAC 0191 MG/L	AS P 24 0	7.4662	3.4922	14.037 (LOG NORMAL)	1.71	17.3	6.805

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94 NO. OF SAMPLES - 23

SAMPLE POINT - S06CLIFFROUDJO  
SAMPLE TYPE - DJ

CLIFF QUAY FINAL EFFLUENT  
PRIMARY SEDIMENTATION TANK EFFLUENT

GRID REF - TM 17200 41900

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD.DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	22 0	7.4018	0.262	7.9179-	6.94	8.1	7.35
SS 105	0135 MG/L.	22 0	155.9636	64.7142	277.4934(LOG NORMAL)	29.0	278.0	151.5
BOD+ATU T	0085 MG/L 0	22 3X	245.695	>84.7956	408.9201(LOG NORMAL)	28.9	395.0	251.5
AMMONIA N	0111 MG/L N	22 0	41.8	13.4477	66.6776(LOG NORMAL)	16.8	71.4	39.75
NITRITE N	0118 MG/L N	11 1<	0.0772-	>0.0337	0.1429(LOG NORMAL)	<0.05	0.17	0.08
NITRATE N	0117 MG/L N	11 9<	0.1063-	>0.2863	0.69 (LOG NORMAL)	<0.01	0.97	0.2
TON AS N	0116 MG/L N	22 22<	0.0	>0.1956	0.5687(LOG NORMAL)	<0.2	<0.6	<0.55
P SOL.REAC	0191 MG/L AS P	22 0	8.8045	2.3032	12.4232(LOG NORMAL)	4.6	12.2	9.05

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 52

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

GRID REF -

CHELMSFORD MIXED FINAL EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

SAMPLE POINT - S03CHELMSDDP  
SAMPLE TYPE - DP

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	37 0	7.5213	0.2115	7.938 - 7.1046	7.31	8.6	7.5
SS	105 C 0135 MG/L	49 0	18.9993	7.9315	33.8987(LOG NORMAL)	7.5	53.5	18.0
BOD+ATU	T 0085 MG/L	49 5<	10.0391	>5.1997	20.2277(LOG NORMAL)	<5.0	28.5	9.3
AMMONIA	N 0111 MG/L	49 0	3.06	2.2076	7.1972(LOG NORMAL)	0.4073	9.54	2.79
TON AS	N 0116 MG/L	36 0	16.9356	3.1643	22.5779(LOG NORMAL)	9.78	22.671	17.483
P SOL-REAC	0191 MG/L AS P	37 0	4.7543	1.0739	6.6934(LOG NORMAL)	2.23	6.58	4.9797

DATE PRODUCED 15/06/95

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 25

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

GRID REF -

CHELMSFORD MIXED FINAL EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

SAMPLE POINT - S03CHELMSDDP  
SAMPLE TYPE - DP

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	24 0	7.5429	0.1378	7.8145 - 7.2712	7.13	7.86	7.555
SS	105 C 0135 MG/L	24 0	22.9156	16.4902	53.8263(LOG NORMAL)	9.0	66.0	15.5
BOD+ATU	T 0085 MG/L	24 3<	12.6458	>7.4698	27.1331(LOG NORMAL)	<3.8	31.1	10.35
AMMONIA	N 0111 MG/L	24 0	2.5379	1.7673	5.8589(LOG NORMAL)	0.694	6.52	1.955
NITRITE	N 0118 MG/L	8 0	0.2716	0.1536	0.589 (MAX VALUE)	0.117	0.589	0.236
NITRATE	N 0117 MG/L	8 0	17.55	3.531	23.2 (MAX VALUE)	13.1	23.2	17.15
TON AS	N 0116 MG/L	24 0	18.9291	3.6288	25.4103(LOG NORMAL)	13.1	25.5	19.7
P SOL-REAC	0191 MG/L AS P	24 0	4.96	0.9301	6.6188(LOG NORMAL)	3.07	6.81	4.83

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 26

SAMPLE POINT - S03CHELMSDDP  
SAMPLE TYPE - DP

CHELMSFORD MIXED FINAL EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(COR RANGE)	STD.-DEV.	95%ILE(COR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	25	7.4896	0.2955	8.0718~	7.2	8.8	7.4
SS	105 C 0135 MG/L	25	13.552 -	>10.2385	33.1113(NORMAL)	4.5	39.0	11.0
BOD+ATU	T 0085 MG/L O	25	6.668 -	>4.6577	15.6887(NORMAL)	2.6	20.1	5.4
AMMONIA	N 0111 MG/L N	25	5.344 -	>4.4227	13.5098(NORMAL)	<0.1	14.7	4.5
NITRITE	N 0118 MG/L N	11	0.239	0.0832	0.3938(NORMAL)	0.13	0.38	0.23
NITRATE	N 0117 MG/L N	11	12.8636 -	>5.7062	23.6336(NORMAL)	<0.5	21.6	13.5
TON AS	N 0116 MG/L N	24	16.1	3.45	22.307 (LOG NORMAL)	9.5	21.8	15.15
P SOL-REAC	0191 MG/L AS P	25	4.884 -	>2.0528	8.7453 (LOG NORMAL)	<0.2	10.3	4.7
INST FLOW	9072 L/S	13	458.8461	63.6032	569.9087 (LOG NORMAL)	350.0	616.0	458.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 12

SAMPLE POINT - S06LOWEST0080  
SAMPLE TYPE - DB

LOWESTOFT NESS POINT SEA OUTFALL  
CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS)

GRID REF - TM 55600 93700

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	12 0	6.4716	0.841	8.1285-	4.67	7.99	6.615
SS	105 C 0135 MG/L	12 0	405.875	235.768	852.0297(LOG NORMAL)	116.5	975.0	393.0
BOD+ATU	T 0085 MG/L O	12 0	677.8058	271.1097	1185.9897(LOG NORMAL)	356.6	1150.0	592.5
AMMONIA	N 0111 MG/L N	12 0	25.0567	8.2065	40.2562(LOG NORMAL)	5.69	35.0	27.176
TON AS	N 0116 MG/L N	11 6<	0.693 -	>0.6215	2.0208(LOG NORMAL)	<0.6	2.1286	<0.6
P SOL-REAC	0191 MG/L AS P	12 0	5.9152	2.4679	10.5512(LOG NORMAL)	1.27	10.4	5.5529

JATE PRODUCED 15/06/95

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 12

SAMPLE POINT - S06LOWEST0080  
SAMPLE TYPE - DB

LOWESTOFT NESS POINT SEA OUTFALL  
CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS)

GRID REF - TM 55600 93700

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	12 0	6.6283	0.9071	8.4154-	4.35	7.63	6.76
SS	105 C 0135 MG/L	12 0	752.8333	637.5461	1925.6425(LOG NORMAL)	266.0	2560.0	530.0
BOD+ATU	T 0085 MG/L O	12 0	890.8333	751.9721	2274.8139(LOG NORMAL)	404.0	3190.0	706.5
AMMONIA	N 0111 MG/L N	12 0	27.5083	8.5437.	43.2766(LOG NORMAL)	11.1	44.4	28.65
NITRITE	N 0118 MG/L N	11 0	0.3718	0.7075	1.3236(LOG NORMAL)	0.0076	2.23	0.095
NITRATE	N 0117 MG/L N	11 5<	0.8841-	>0.9147	2.6698(LOG NORMAL)	<0.5	2.66	<0.6
TON AS	N 0116 MG/L N	12 5<	1.357 -	>1.4917	4.1351(LOG NORMAL)	<0.6	4.85	0.6695
P SOL-REAC	0191 MG/L AS P	12 0	5.7008	1.1837	7.826 (LOG NORMAL)	2.79	7.51	5.725

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

NO. OF SAMPLES - 12

SAMPLE POINT - SD6LOWESTDBO LOMESTOFT NESS POINT SEA OUTFALL  
 SAMPLE TYPE - DB CRUDE SEWAGE(AT SEWAGE TREATMENT WORKS)

GRID REF - TM 55600 93700

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD. DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	12 0	6.8916	0.4962	7.8693-	5.7	7.6	7.0
SS	0135 MG/L	12 0	425.5	144.6409	694.0394(LOG NORMAL)	88.0	623.0	419.5
BO+ATU T	0085 MG/L	10 1>	501.5535*	>218.4829	912.7724(LOG NORMAL)	>70.7	875.0	661.0
AMMONIA N	0111 MG/L	12 0	25.6333	14.5336	55.1795(LOG NORMAL)	2.18	51.4	25.65
NITRITE N	0118 MG/L	11 2<	0.0745-	>0.0366	0.1463(LOG NORMAL)	<0.007	0.14	0.08
NITRATE N	0117 MG/L	11 11<	0.0	>0.2018	0.5241(LOG NORMAL)	<0.1	<0.5	<0.5
TION AS N	0116 MG/L	12 8<	0.7358-	>0.9918	2.5859(LOG NORMAL)	<0.1	3.1	<0.6
P SOL.REAC	0191 MG/L AS P	10 0	6.442	2.792	11.6958(LOG NORMAL)	1.82	10.0	6.55

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

SAMPLE POINT - S03BURNHAMDP BURNHAM FINAL EFFLUENT  
 SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	22 0	7.5972	0.1466	7.8862- 7.3083	7.32	7.92	7.6
SS	105 C 0135 MG/L	24 2<	13.5	>12.071	35.7746(LOG NORMAL)	<5.0	59.0	10.0
BOD+ATU	T 0085 MG/L O	24 10<	5.4687	>11.7814	25.7888(LOG NORMAL)	2.2	<60.0	<5.61
AMMONIA	N 0111 MG/L N	24 1<	14.0211	>14.8888	40.1445(LOG NORMAL)	<0.2	38.9	5.29
TON AS	N 0116 MG/L N	21 4<	12.7661	>12.0317	34.5581(LOG NORMAL)	<0.6	34.344	9.79
P SOL.REAC	0191 MG/L AS P	22 1<	6.1865	>2.5848	11.049 (LOG NORMAL)	<0.3	9.4	7.0287

DATE PRODUCED 15/06/95 N.R.A. ANGLIAN REGION - CHEMICAL DATA PROCESSING SYSTEM PAGE NO. 1

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

SAMPLE POINT - S03BURNHAMDP BURNHAM FINAL EFFLUENT  
 SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	23 0	7.4713	0.1386	7.7445- 7.198	7.31	7.78	7.46
SS	105 C 0135 MG/L	23 10<	5.413	>4.7473	15.3874(LOG NORMAL)	<5.0	25.5	5.5
BOD+ATU	T 0085 MG/L O	23 14<	2.7304	>3.2591	10.9503(LOG NORMAL)	<3.1	14.3	5.3
AMMONIA	N 0111 MG/L N	23 0	2.9016	4.5624	9.7581(LOG NORMAL)	0.249	20.7	0.996
NITRITE	N 0118 MG/L-N	3 0	0.585	0.1646	0.712 (MAX VALUE)	0.399	0.712	0.644
NITRATE	N 0117 MG/L-N	3 0	16.3656	3.3005	19.1 (MAX VALUE)	12.7	19.1	17.3
TON AS	N 0116 MG/L N	23 0	15.6152	4.7503	24.3701(LOG NORMAL)	5.5	23.7	17.2
P SOL.REAC	0191 MG/L AS P	23 0	6.3447	1.3187	8.7124(LOG NORMAL)	3.08	9.13	6.22
CHLORIDE	0172 MG/L CL	23 0	173.5652	27.7797	224.5979(LOG NORMAL)	131.0	225.0	166.0
HG TOTAL	9269 UG/L HG	3 3<	0.0	>0.0	<0.05 (MAX VALUE)	<0.05	<0.05	<0.05
E COLI P	2549 NO/100ML	2 0	58250.0	46174.0727	90900.0 (MAX VALUE)	25600.0	90900.0	58250.0
CHLOROFORM	9067 UG/L	10 0	1.453	0.6201	2.6191(LOG NORMAL)	0.79	2.4	1.35
CHBR2CL	9068. UG/L	10 0	0.511	0.511	1.424 (LOG NORMAL)	0.16	1.9	0.33
CHBR2CL	9069 UG/L	10 0	0.331	0.0911	0.4977(LOG NORMAL)	0.19	0.46	0.325
BROMOFORM	9070 UG/L	10 10<	0.0	>0.0431	0.1391(LOG NORMAL)	<0.1	<0.2	<0.1
INST FLOW	9072 L/S	5 0	187.0	25.9711	225.0 (MAX VALUE)	166.0	225.0	173.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 24

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

SAMPLE POINT - S03BURNHAMDP BURNHAM FINAL EFFLUENT  
SAMPLE TYPE - DP FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD. DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	21 0	7.4819	0.1693	7.8154- 7.1483	7.2	7.8	7.5
SS 105 C	0135 MG/L	21 4<	7.8285-	>13.0783	27.8902(LOG NORMAL)	1.2	65.0	6.0
BOD+ATU T	0085 MG/L.0	21 9X	2.235 -	>8.0768.	13.3771(LOG NORMAL)	<1.0	>39.7	<3.0
AMMONIA N	0111 MG/L N	22 6<	3.8727-	>7.3953	13.8536(LOG NORMAL)	<0.1	33.7	1.0
NITRITE N	0118 MG/L N	2 1>	0.15 *	>0.502	>0.86 (MAX VALUE)	0.15	>0.86	>0.505
NITRATE N	0117. MG/L N	2 0	14.95	2.192	16.5 (MAX VALUE)	13.4	16.5	14.95
TON AS N	0116 MG/L N	22 1<	12.7409-	>6.9138	25.8318(LOG NORMAL)	<0.2	34.6	13.05
P SOL-REAC	0191 MG/L AS P	22 0	6.3045	2.2312	10.6495(LOG NORMAL)	3.6	10.0	5.6
CHLORIDE	0172 MG/L CL	22 0	170.0	35.3351.	233.4435(LOG NORMAL)	117.0	277.0	160.5
HG TOTAL	9269 UG/L HG	3 2<	0.0043-	>0.0213	<0.05 (MAX VALUE)	0.013	<0.05	<0.05
COLI P	9193 NO/100ML	2 1>	13930.0 *	>4313.3513	>20300.0 (MAX VALUE)	13900.0	>20000.0	>16950.0
S FAEC P	2346 NO/100ML	1 0	1730.0	0.0	1700.0 (ONLY VALUE)	1700.0	1700.0	1700.0
E COLI P	2549 NO/100ML	2 1>	8530.0 *	>8131.7279	>20300.0 (MAX VALUE)	8500.0	>20000.0	>14250.0
CHLOROFORM	9067 UG/L	8 0	1.2375	0.4068	2.2 (MAX VALUE)	0.9	2.2	1.1
CHBRCL2	9068 UG/L	8 5<	0.2562-	>0.1751	1.0 (MAX VALUE)	<0.5	1.0	<0.5
CHBR2CL	9069 UG/L	8 6<	0.135 -	>0.0875	0.7 (MAX VALUE)	0.38	0.7	<0.5
BROMOFORM	9070 UG/L	8 8<	0.0	>0.3412	<1.0 (MAX VALUE)	<0.1	<1.0	<1.0
INST FLOW	9072 L/S	17 0	189.3235	24.7019	232.4627(LOG NORMAL)	146.0	228.0	190.0

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/92 TO 31/12/92

NO. OF SAMPLES - 54

SAMPLE POINT - S03COLCHTRDP  
COLCHESTER COMBINED EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	44	7.6584	0.1269	7.9085-	7.35	8.07	7.65
SS	0135 MG/L	52	16.1826	>22.2836	52.6915(LOG NORMAL)	4.5	143.0	10.75
BOD+ATU T	0085 MG/L O	51	14.7309	>11.4552	34.5756(LOG NORMAL)	2.93	73.5	10.9
AMMONIA N	0111 MG/L N	52	13.8426	14.8539	39.8235(LOG NORMAL)	0.386	78.068	5.935
TON AS N	0116 MG/L N	44	8.3176	>5.4389	18.5966(LOG NORMAL)	<0.6	16.6	8.395
P SOL.REAC	0191 MG/L AS P	44	6.1224	2.1939	10.2091(LOG NORMAL)	0.341	12.9	6.2982

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

REPORT COVERS PERIOD(S) FROM 01/01/93 TO 31/12/93

NO. OF SAMPLES - 34

SAMPLE POINT - S03COLCHTRDP  
COLCHESTER COMBINED EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

GRID REF -

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	32	7.8009	0.1052	8.0083-	7.53	8.02	7.81
SS	0135 MG/L	32	21.125	36.9936	73.4994(LOG NORMAL)	6.0	219.0	13.75
BOD+ATU T	0085 MG/L O	32	13.9968	>28.6531	53.306 (LOG NORMAL)	<4.2	169.0	9.5
AMMONIA N	0111 MG/L N	32	22.2662	7.339	35.9041(LOG NORMAL)	4.6	33.4	25.65
NITRITE N	0118 MG/L N	10	0.7599	0.6315	1.9561(LOG NORMAL)	0.111	2.27	0.589
NITRATE N	0117 MG/L N	10	1.931	1.1081	4.0282(LOG NORMAL)	0.873	4.55	1.675
TON AS N	0116 MG/L N	32	2.5272	>2.3224	6.7646(LOG NORMAL)	<0.6	10.8	1.81
P SOL.REAC	0191 MG/L AS P	32	6.1265	5.3426	15.9091(LOG NORMAL)	1.74	34.2	5.205

OUTPUT FROM GENERAL DATA ABSTRACTION FACILITY

REPORT TYPE 712 - PART 3 - STATISTICAL SUMMARY REPORT

NO. OF SAMPLES - 27

REPORT COVERS PERIOD(S) FROM 01/01/94 TO 31/12/94

GRID REF -

COLCHESTER COMBINED EFFLUENT  
FINAL SEDIMENTATION TANK EFFLUENT

SAMPLE POINT - SOICOLCHTRDP  
SAMPLE TYPE - DP

DETERMINAND	UNITS	NO. OF VALUES	MEAN VALUE(OR RANGE)	STD.DEV.	95%ILE(OR RANGE)	MINIMUM	MAXIMUM	MEDIAN
PH	0061 PH UNITS	26 0	7.5915	0.2527	8.0893-	6.6	7.9	7.635
SS	105 C 0135 MG/L	26 2<	20.15 -	>33.9433	70.018 (LOG NORMAL)	4.0	166.0	12.0
BOI+ATU	T 0085 MG/L O	24 2>	10.6727*	>10.6522	29.6556(L06 NORMAL)	2.8	57.2	8.8
AMMONIA	N 0111 MG/L N	26 0	20.8038	8.994	37.7263(L06 NORMAL)	4.0	35.7	19.85
NITRITE	N 0118 MG/L N	10 0	0.613	0.6305	1.7284(L06 NORMAL)	0.03	1.79	0.4
NITRATE	N 0117 MG/L N	10 3<	1.953 -	>2.7772	6.4846(L06 NORMAL)	0.3	8.8	0.75
TON AS	N 0116 MG/L N	25 6<	3.416 -	>3.761	10.0575(L06 NORMAL)	<0.6	10.4	1.0
P SOL-REAC	0191 MG/E AS P	26 0	5.6484	2.0635	9.4966(L06 NORMAL)	1.9	11.3	5.55

# CITY OF TAMPA

Sandra W. Freedman, Mayor

Department of Sanitary Sewers

Advanced Wastewater Treatment Plant



FAX NO. 813-248-5269

Advanced Primary  
OR  
Secondary Treatment

which one

- Dump directly into Gulf?  
OR Tampa Bay  
- Where does your  
outfall go?  
- Industrial Load - Is it  
Part of the Total?

FAX TRANSMITTAL SHEET

TO: NCCOSC RDTE DIV 5221  
ATTN: Bart Chadwick  
FROM: John Drapp


We are sending 3 pages, including coversheet.

Per request of Melissa!



2700 Maritime Boulevard • Tampa, Florida 33605 • 813/247-3451

- Secondary Treatment  
- Discharges into Tampa Bay which runs into the Gulf



Post-It® Fax Note	7671	Date	2-9-95	# of pages	1
To	Melissa	From	John Drapp		
Co./Dept.	WCCOPE RDTE	Co.	City of Tampa		
Phone #		Phone	(813) 247-3451		
Fax #	(619) 553-5404	Fax #			



# PERMIT REQUIREMENTS

**PERMITTEE:**

 City of Tampa  
 Hookers Point AWTP

 GMS ID NO.: 4029M03950  
 PERMIT NO.: DO29-184532B

**SPECIFIC CONDITIONS:**

1. Drawings, plans, documents or specifications submitted by the permittee, not attached hereto, but retained on file at the Southwest District Office, are made a part hereof.

2. In accordance with Chapter 17-602, F.A.C., the required certified operator on site time is: A Class C or better operator for 24 hours/day, and 7 days a week. The lead/chief operator must be Class A.

3. The effluent shall be sampled in accordance with Chapter 17-601, F.A.C. and shall meet the following limitations:

Parameter	Unit	Minimum	Maximum	Type	Sample Frequency
CBOD5 and Suspended Solids	mg/l	-	5 annual avg	**fpc	Daily 7 days/wk
Influent CBOD5 and Suspended Solids	mg/l	Report	-	**fpc	Daily 7 days/wk
Fecal Coliform	#100	*Non- detectable	25	grab	Daily 7 days/wk
Total Nitrogen	mg/l	-	3 annual avg	**fpc	Daily 7days/wk
Ortho Phosphorous	mg/l	Report	-	**fpc	Daily 7days/wk
Total Phosphorous	mg/l	Report	-	**fpc	Daily 7days/wk
Total Residual Chlorine	mg/l	-	0.01	grab	Hourly/24 any sample hrs/day
Flow	mgd	-	70.0 ADF	***rmf&t	Continuous
pH	STD UN	6.00	8.50	****meter	Continuous
DO	mg/l	5.00	-	****meter	Continuous

\* Non-detectable in at least seventy-five (75%) of samples collected during the monthly operating period (e.g. 23 per 30 samples)

\*\* fpc-Flow Proportional Composite (24 Hours)

\*\*\* rmf&t-Recording Flowmeter and Totalizer

\*\*\*\* hourly measurements for 24 hours may be substituted for continuous measurement

The results shall be reported monthly on DER Form 17-601.900(1) to

27 de Mayo a Julio de 1995  
 13:00

COMISION DE SERVICIOS DE AGUA DEL ESTADO  
 SISTEMA DE ALEJAMIENTO Y TRATAMIENTO DE AGUAS RESIDUALES  
 DE LA CIUDAD DE TIJUANA, B.C.

CALIDAD DE AGUA RECIBIDA Y PRODUCIDA POR EL SISTEMA.

UNIDAD	PARAMETRO	AFLUENTE DE LA PLANTA DE TRATAMIENTO	EFLUENTE DE LA PLANTA DE TRATAMIENTO
mg/l	D.B.O.5 (total)	427.7	54.6
mg/l	OXIGENO DISUELTO	0.3	2.8
mg/l	DEMANDA QUIMICA DE OXIGENO (total)	811.0	117.0
mg/l	GRASAS Y ACEITES	244.3	50.9
mg/l	SOLIDOS SUSPENDIDOS TOTALES TSS	302.7	41.5
ml/l	SOLIDOS SEDIMENTABLES	5.4	0.0
mg/l	FOSFORO TOTAL P <sub>T</sub>	7.3	3.8
mg/l	NITROGENO AMONICAL NH <sub>3</sub>	47.1	20.6
mg/l	DETERGENTES S.A.A.M.	29.8	16
NMP/100 ml	COLIFORMES TOTALES	24000E6	<250
NMP/100 ml	COLIFORMES FECALES	24000E6	<250
°C	TEMPERATURA	21.3	21.0
PH	POTENCIAL DE HIDROGENO	7.2	7.3

**PART 1: INFLUENT FLOW/LOADINGS**

A. List the average monthly volumetric flows and BOD<sub>5</sub> loadings received at your facility during your 12 month MWFP reporting period. (Influent sampling should be at the same frequency as the required effluent sampling.)

MWFP Reporting Period		Col. 1 Average Monthly Influent Flow	Col. 2 Average Monthly Influent BOD <sub>5</sub> Concentrations	Col. 3 Average Monthly Influent BOD <sub>5</sub> Loading
Year	Month	(MGD)	(mg/l)	(pounds per day)
92	July	10.72	150	12,046
92	August	13.68	111	10,765
92	September	15.67	89	10,887
92	October	10.18	155	12,881
92	November	16.74	89	10,534
92	December	14.41	114	12,765
93	January	14.57	103	11,859
93	February	10.95	143	13,217
93	March	13.26	121	13,664
93	April	11.68	143	13,019
93	May	11.17	141	12,525
93	June	9.58	168	13,055

Give source of data listed above: Data obtained from laboratory testing of daily influent monitoring performed by the Jefferson Parish Department of Environment and Development Control.

1

11.5 12,775

11.05

1.25

# New Orleans Treatment Plant

## PART 2: EFFLUENT QUALITY/PLANT PERFORMANCE

A. For the permitted parameters, list the average monthly effluent concentration and average monthly loading produced by your facility during your 12 month RPP reporting period. Disregard any data which are not applicable to your permit. Circle whether you are measuring ammonia nitrogen ( $\text{NH}_3\text{-N}$ ), nitrate nitrogen ( $\text{NO}_3\text{-N}$ ), or

(1) Concentration

Year	Month	BOD <sub>5</sub> (mg/l)	TSS (mg/l)	$\text{NH}_3\text{-N}$ or $\text{NO}_3\text{-N}$ (mg/l)	Total Phosphorus (mg/l)	Fecal Coliform (count/ 100 ml)	pH (Lowest/ Highest)	Other Cl Resid
92	July	13.1	7.8			11	7.04 / 7.72	1.07
92	August	18.2	6.5			31	7.07 / 7.82	0.65
92	September	14.3	5.9			29	7.07 / 7.74	1.04
92	October	5.7	3.6			9	7.08 / 7.78	1.03
92	November	5.5	3.8			25	7.09 / 7.84	1.31
92	December	7.5	5.1			9	6.70 / 7.84	1.88
93	January	8.0	6.1			11	7.11 / 7.75	1.16
93	February	9.6	4.8			9	6.70 / 7.84	1.9
93	March	11.5	8.1			12	7.10 / 7.90	1.6
93	April	14.2	13.2			11	9.98 / 7.69	1.1
93	May	12.7	10.2			12	9.93 / 7.45	1.2
93	June	16.7	7.7			9	7.02 / 7.37	1.1

(2) Average Monthly Mass Loading

MWPP Reporting Period						
Year	Month	BOD <sub>5</sub> (lbs/day)	TSS (lbs/day)	NH <sub>3</sub> -N or NO <sub>3</sub> -N (lbs/day)	Total Phosphorus (lbs/day)	Other
92	July	1,119	673			
92	August	2,010	709			
92	September	1,912	856			
92	October	488	301			
92	November	773	556			
92	December	904	691			
93	January	983	780			
93	February	884	462			
93	March	1,276	883			
93	April	1,415	1,396			
93	May	1,189	989			
93	June	1,318	611			

B. List the monthly permit limits for the facility in the blanks below. Circle whether your permit lists ammonia nitrogen (NH<sub>3</sub>-N) or nitrate nitrogen (NO<sub>3</sub>-N).

(1) Concentration (Attach additional sheets for other if necessary.)

Fecal Coliform (Count/100 ml)	BOD <sub>5</sub> (mg/l)	TSS (mg/l)	NH <sub>3</sub> -N or NO <sub>3</sub> -N (mg/l)	Total Phosphorus (mg/l)	Other CL Residual (mg/l)	Other	Other
200/400	30/45	30/45		1.43			
Permit Limiter:							
90% of the Permit Limiter:	27	27		1.29			

(2) Average Monthly Mass Loading (Attach additional sheets for other if necessary.)

BOD <sub>5</sub> (lbs/day)	TSS (lbs/day)	NH <sub>3</sub> -N or NO <sub>3</sub> -N (lbs/day)	Total Phosphorus (lbs/day)	Other	Other	Other
2,402	2,402					
Permit Limiter:						
90% of the Permit Limiter:	2,162					

Houston Treatment Plant

NO<sub>2</sub> DO<sub>2</sub>

HARRIS COUNTY W.C.I.D. #1  
WASTEWATER TREATMENT PLANT  
1994

	<u>CBOD</u>	<u>TSS</u>	<u>NAM</u>	<u>TKN</u>	<u>D.O.</u>	<u>PH</u>	<u>SS.EFF</u>	<u>CL2.RES</u>	<u>SB2RES</u>	<u>SOLIDS</u>	<u>AERA.D.O.</u>	<u>FLOW MD. AVG</u>	<u>GALS PER MONTH</u>
<u>JAN.</u>	3.80	9.0	0.4	1.50	8.70	7.5	0.00	2.20	0.01	540	4.30	1,285,000	39,846,000
<u>FEB.</u>	2.80	6.0	0.2	1.20	8.80	7.6	0.00	1.80	0.02	470	4.50	1,452,000	40,677,000
<u>MAR.</u>	3.30	8.0	0.1	1.00	8.30	7.4	0.01	2.20	0.01	550	3.90	1,457,000	45,157,000
<u>APR.</u>	2.80	4.9	0.3	1.70	8.00	7.1	0.00	2.60	0.01	770	1.80	988,000	29,651,000
<u>MAY</u>	2.80	6.0	0.0	0.90	7.70	7.1	0.00	1.80	0.01	540	3.30	1,709,000	53,008,000
<u>JUN.</u>	3.10	5.3	0.2	1.00	7.80	7.0	0.00	3.00	0.01	600	2.30	1,152,000	34,869,000
<u>JUL.</u>	2.30	6.0	0.6	0.80	6.80	7.1	0.00	2.60	0.01	650	2.30	910,000	28,211,000
<u>AUG.</u>	2.50	3.8	0.0	0.70	6.60	7.2	0.00	4.00	0.01	410	4.00	851,000	26,692,000
<u>SEPT.</u>	2.80	5.0	0.0	0.80	6.70	7.0	0.00	6.00	0.02	540	3.40	757,800	22,733,000
<u>OCT.</u>	3.00	7.0	0.0	1.10	7.30	6.90	0.00	6.20	0.02	440	4.50	2,001,200	62,038,000
<u>NOV.</u>	3.30	6.0	0.0	0.80	6.60	7.10	0.00	3.20	0.02	610	2.90	843,000	25,313,000
<u>DEC.</u>	3.20	5.0	0.0	0.80	7.30	7.20	0.00	2.40	0.02	590	2.60	1,761,000	54,600,000
<u>TOTAL</u>	<u>35.70</u>	<u>72.0</u>	<u>1.80</u>	<u>12.30</u>	<u>90.60</u>	<u>86.20</u>	<u>0.01</u>	<u>38.00</u>	<u>0.17</u>	<u>6710</u>	<u>39.80</u>	<u>15,191,200</u>	<u>462,795,000</u>
<u>AVG.</u>	3.00	6.00	0.20	1.00	7.60	7.20	0.00	3.20	0.01	560	3.30	1,265,900	38,566,250

NO<sub>2</sub> DO<sub>2</sub>

23

553-5404

# FAX

City of San Diego  
Metropolitan Wastewater Department

## Wastewater Chemistry Laboratory



5530 Kiowa Drive  
La Mesa, CA 91942  
FAX: (619) 668-3250



Please deliver to: MARILISSA CABALLERO

Organization: NAVY

FAX Number: 553-5404

Total pages sent: 5  
(including this one)

Sent by: WALTER KONOPKA

Phone: 668-3205

### Notes/Comments:

*the Point Loma Outfall  
These are the pages from the California  
Discharge Permit  
that give waste discharge requirements.*

*Effluent*

*1994 Avg annual 64.5 mg/L      came into Plant 78.4 mg/L*

- Monthly JAN - 54.8*
- Feb - 55.3*
- MARCH - 71.0*
- April - 76.8*
- MAY - 71.3*
- JUNE - 65.7*
- JULY - 80.5*
- AUG - 56.9*
- SEPT - 54.0*
- OCT - 61.7*
- NOV - 77.6*
- DEC - 58.5*

*TCC = 04.5 mg/L  
N = -  
P = 3.0 ...*

*16  
=*

Order No. 90-32

18

B. DISCHARGE SPECIFICATIONS<sup>1</sup>

1.a. The discharge of waste through the Point Loma Ocean Outfall containing pollutants in excess of the following effluent limitations are prohibited:

Constituent	Units	6-Month <sup>2</sup>	30-Day <sup>3</sup>	7-Day <sup>4</sup>	Daily <sup>5</sup>	Instantaneous <sup>6</sup>
		Median	Average	Average	Maximum	Maximum
Biochemical	mg/L	---	30	45	50	50
Oxygen Demand	lb/day	---	55,000	88,000	91,000	91,000
BOD <sub>5</sub> @ 20° C						
Suspended	mg/L	---	30	45	50	50
Solids	lb/day	---	55,000	88,000	91,000	91,000
pH	pH Units	---	Within the limits of 6.0 - 9.0 at all times			
Grease & Oil	mg/L	---	25	40	75	75
	lb/day	---	46,000	73,000	140,000	140,000
Settleable	ml/L	---	1.0	1.5	3.0	3.0
Solids						
Turbidity	NTU	---	75	100	225	225
Acute Toxicity <sup>8</sup>	TUa	---	1.5	2.0	2.5	2.5
Arsenic	mg/L	0.005	---	---	0.02	0.05
	lb/day	10	---	---	40	100
Cadmium	mg/L	0.011	---	---	0.044	0.11
	lb/day	20	---	---	80	200
Chromium	mg/L	0.05	---	---	0.2	0.5
(Hexavalent) <sup>9</sup>	lb/day	90	---	---	360	900
Copper	mg/L	0.07	---	---	0.28	0.7
	lb/day	130	---	---	520	1300
Lead	mg/L	0.1	---	---	0.4	1.0
	lb/day	160	---	---	640	1600
Mercury	mg/L	0.0005	---	---	0.002	0.005
	lb/day	0.9	---	---	3.6	9
Nickel	mg/L	0.06	---	---	0.24	0.6
	lb/day	110	---	---	440	1100
Selenium	mg/L	0.0014	---	---	0.0056	0.014
	lb/day	2.3	---	---	9.2	23

Order No. 90-32

19

Constituent	Units <sup>7</sup>	6-Month <sup>2</sup> Median	30-Day <sup>3</sup> Average	7-Day <sup>4</sup> Average	Daily <sup>5</sup> Maximum	Instantaneous <sup>6</sup> Maximum
Silver	mg/L lb/day	0.03 50	---	---	0.12 200	0.3 500
Zinc	mg/L lb/day	0.08 140	---	---	0.32 560	0.8 1400
Cyanide <sup>10</sup>	mg/L lb/day	0.01 20	---	---	0.04 80	0.1 200
Total Residual Chlorine (TRC) <sup>11</sup>	mg/L lb/day	0.23 320	---	---	0.9 1300	6.9 9,700
Ammonia (ex- pressed nitrogen)	mg/L lb/day	60 80,000	---	---	260 360,000	600 800,000
Chronic Toxicity <sup>12</sup>	TUc	---	---	---	113	---
Phenolic Com- pounds (nonchlorinated)	mg/L lb/day	0.014 26	---	---	0.056 100	0.14 260
Chlorinated Phenolics	mg/L lb/day	0.0036 6.8	---	---	0.014 27	0.036 68
Endosulfan <sup>13</sup>	ug/L lb/day	0.07 0.13	---	---	0.14 0.26	0.21 0.39
Endrin	ug/L lb/day	0.006 0.011	---	---	0.012 0.022	0.018 0.033
HCH <sup>14</sup>	ug/L lb/day	0.13 0.24	---	---	0.26 0.48	0.39 0.72
Radioactivity	Not to exceed limits specified in Title 17, Chapter 5, Subchapter 4, Group 3, Article 3, Section 30269 of the California Code of Regulations.					

Note:

- mg/L = milligrams per liter
- NTU = Nephelometric turbidity units
- TUa = Acute toxicity units
- TUc = Chronic toxicity units
- ug/L = micrograms per liter
- lb/day = pounds per day

Order No. 90-32

20

## B.1.b. LIMITATIONS FOR PROTECTION OF HUMAN HEALTH--NONCARCINOGENS

Constituent	30-day Average	
	(ug/L)	(lb/day)
Acrolein	7	13
Antimony	320	580
Bis(2-chloroethoxy)methane	53	100
Bis(2-chloroisopropyl) ether	57	100
Chlorobenzene	60	110
Chromium (III)	70	130
di-n-butyl phthalate	25	45
Dichlorobenzenes <sup>14</sup>	44	80
1,1-dichloroethylene	28	50
Diethyl phthalate	19	35
Dimethyl phthalate	16	30
4,6-dinitro-2-methylphenol	240	430
2,4-dinitrophenol	420	760
Ethylbenzene	72	130
Fluoranthene	22	40
Hexachlorocyclopentadiene	4	7
Isophorone	22	40
Nitrobenzene	19	35
Thallium	400	730
Toluene	100	180
1,1,2,2-tetrachloroethane	69	130
Tributyltin	0.10	0.2
1,1,1-trichloroethane	38	70
1,1,2-trichloroethane	50	90

## B.1.c. LIMITATIONS FOR PROTECTION OF HUMAN HEALTH -- CARCINOGENS

Acrylonitrile	3	5
Aldrin	0.0025	0.005
Benzene	22	40
Benzidine	0.008	0.015
Beryllium	3.7	6.8
Bis(2-chloroethyl) ether	0.005	0.009
Bis(2-ethylhexyl) phthalate	10	20
Carbon Tetrachloride	14	25
Chlordane <sup>15</sup>	0.003	0.005
Chloroform	50	90
DDT <sup>16</sup>	0.02	0.04
1,4-dichlorobenzene	22	40
3,3-dichlorobenzidine	0.9	1.6
1,2-dichloroethane	14	25
Dichloromethane	450	820
1,3-dichloropropene	30	45

Order No. 90-32

21

B.1.C CONT'D

Constituent	30-day Average	
	(ug/L)	(lb/day)
Dieldrin	0.005	0.009
2,4-dinitrotoluene	29	50
1,2-diphenylhydrazine	18	30
Halomethanes <sup>17</sup>	24	45
Heptachlor <sup>18</sup>	0.02	0.04
Hexachlorobenzene	0.02	0.04
Hexachlorobutadiene	5	8
Hexachloroethane	8.0	15
N-nitrosodimethylamine	250	450
N-nitrododiphenylamine	9.5	20
PAHs <sup>19</sup>	1.0	1.8
PCBs <sup>20</sup>	0.002	0.004
TCDD equivalents <sup>21</sup>	0.0004	0.0007
Tetrachloroethylene	50	90
Toxaphene	0.02	0.04
Trichloroethylene	9.5	20
2,4,6-trichlorophenol	14	25
Vinyl Chloride	0.90	1.6

2. The arithmetic mean of biochemical oxygen demand and total suspended solids values, by weight, for effluent samples collected in the period of 30 consecutive days shall not exceed 15 percent of the arithmetic mean of values, by weight, for influent samples collected at approximately the same times during the same period.
3. Waste discharged through the Point Loma Ocean Outfall must be essentially free of:
  - (a) Material that is floatable or will become floatable upon discharge.
  - (b) Settleable material or substances that form sediments which degrade<sup>23</sup> benthic communities or other aquatic life.
  - (c) Substances which will accumulate to toxic levels in marine waters, sediments or biota.
  - (d) Substances that significantly<sup>24</sup> decrease the natural light to benthic communities and other marine life.
  - (e) Materials that result in aesthetically undesirable discoloration of the ocean surface.

## **APPENDIX L**

### **BIBLIOGRAPHY**

Source: Bibliography  
San Diego, California  
Naval Command, Control & Ocean Surveillance  
Center, RDTE Division, Code 522, 1995

## BIBLIOGRAPHY

At the onset of this project, a literature review was conducted to obtain information useful to understanding the fate and effects of solid wastes discharges to the ocean in general, and within Special Areas in particular. Some of this information has already been used in the previous sections. The literature was examined for previous studies performed within Special Areas, as well as for studies that related to more general issues such as the characteristics of discharged materials, regulations, and naval operations. As a result, the search covered a broad range of topics including:

- Regulatory Framework and Regulations
- Oceanographic and Meteorological Conditions
- Environmental Conditions
- Ecology and Fisheries
- General Waste Dumping
- Pulp Mill Discharges
- Sewage Treatment Plant Discharges
- River and Other Industrial Discharges
- Operations Conditions including ship traffic patterns, ship types, and waste generation
- Composition of Glass, Metal, and Paper Material
- Corrosion Processes
- Dispersion and Dispersion Modeling

The University of California library system was the main system used to search and acquire literature. This was done using the library's computerized search capability and utilizing the large holdings available at the University of California, San Diego (UCSD). The NRaD library system was also used during this search, particularly when interlibrary loans were required.

The literature was first searched using the names of five seas (Mediterranean, Baltic, North, Caribbean, and Antarctic) in the search parameter (subject search). The database of books, research journal articles, and government reports generated under this search included thousands of titles. These were subjectively reviewed for titles that seemed appropriate for the study at hand, resulting in an annotated list composed of hundreds of titles. An attempt was made to obtain all the citations from this annotated bibliography, although this was not always possible or time effective. The literature obtained was reviewed for useful content, photocopied as necessary, and returned to the library.

The bibliography presented in this section is a result of the above searches and contains all titles that were obtained for review. The bibliographic information is therefore more extensive than just those references noted in the report.

- 1991 MARPOL Amendments Enter into Force. *Marine Pollution Bulletin*, V26, N6, pp. 294-295, 1993.
- 1993 Annual Report. Waste Management Institute, Stony Brook, New York, 1993.
- A Review of The Past and Present Measurements of Selected Trace Metals in Sea Water in the OSLO Commission and ICNAF (NAFO) Areas*. Copenhagen: International Council for the Exploration of the Sea, 1980.
- Abelmann A., R. Gersonde. *Biosiliceous Particle Flux In The Southern Ocean*. *Marine Chemistry*, V35 N1-4:503-536, 1991.
- Accumulation and Transformation of Chemical Contaminants by Biotic and Abiotic Processes in the Marine Environment, La Spezia, Italy*. FAO/UNEP/IAEA Consultation Meeting on the Accumulation and Transformation of Chemical Contaminants by Biotic and Abiotic Processes in the Marine Environment (1990: La Spezia, Italy) Proceedings of the FAO/UNEP/IAEA Consultation Meeting on the Athens UNEP, 1991.
- Aijaz, S., S. A. Jenkins. *On the Electrokinetics of the Dynamic Shear Stress in Fluid Mud Suspensions*. *Journal of Geophysical Research*, V99, NC6, pp. 12, 697-12,706, 1994.
- Ailor, W.H. Jr. *Aluminum Alloys After Five Years in Seawater. Materials Performance and the Deep Sea*. ASTM STP 445, 115-130, American Society for Testing and Materials, 1969.
- Airborne Pollution of The Mediterranean Sea: Report And Proceedings of The Second IMO/UNEP Workshop*. Athens, UNEP, 1992.
- Aitsam, A., J. Elken, L. Talsepp, J. Laanemets. *Topographically Induced Variability in the Baltic Sea*. Institute of Thermophysics and Electrophysics Academy of Sciences of Estonian SSR.
- Akerblom, A. *3rd Soviet-Swedish Symposium on the Pollution of the Baltic*. The National Swedish Environment Protection Board, Stockholm, 1975.
- Akerblom, A. *3rd Soviet-Swedish Symposium on the Pollution of the Baltic*. Sovetskoshvedskii Simpozium po Zagriazneniiu Baltiiskogo Moria, Oslo: National Swedish Environment Protection Board, Irvington-on-Hudson, N.Y., Columbia University Press, 1978.
- Alig, C.S. *Dispersion of Sanitary Wastewater Discharges From Navy Ships*. Naval Ship Research and Development Center Annapolis, Maryland, August, 1974.
- Allredge, A.L., C.C. Gotshalk. *Bacterial Carbon Dynamics on Marine Snow*. *Continental Shelf Research*, V10, pp. 41-58, 1990.
- Allen W. *Increased Dangers To Caribbean Marine Ecosystems - Cruise Ship Anchors and Intensified Tourism Threaten Reefs*. *Bioscience*, V42 N5, pp. 330-335, 1992.
- Aller, R.C. *Diagenetic Processes Near the Sediment-Water Interface of Long Island Sound I. Decomposition and Nutrient Element Geochemistry (S, N, P)*. Ed. B. Saltzman, Estaurine

- Physics and Chemistry: Studies in Long Island Sound, Advances in Geophysics, V22, Academic Press, 1980.
- Amann, R.I., B.J. Binder, R.J. Olson, S.W. Chisolm, R. Devereux, D.A. Stahl. *Combination Of 16S Rrna-Targeted Oligonucleotide Probes With Flow Cytometry For Analyzing Mixed Microbial Populations*. Applied Environmental Microbiology. V56, pp.1919-1925, 1990.
- American Meteorological Society. *Bibliography On Meteorology, Climatology, and Physical/Chemical Oceanography*. Washington [US National Oceanographic Data Center], 1970.
- American Society for Testing and Materials (ASTM). *Annual Book of ASTM Standards, Standard A 624*. ASTM, Philadelphia, Pennsylvania, 1984.
- American Society for Testing and Materials. *Annual Book of ASTM Standards, ASTM Standard Practice G1-81*. ASTM, Philadelphia, Pennsylvania, 1981.
- Andrulewicz, E., Rohde, K.H. *Harmful Substances: Petroleum Hydrocarbons*. Helsinki: Baltic Marine Environment Protection Commission, Baltic Sea Environment Proceedings, N17 B, 351, pp. 170-198, 1987.
- Anglophone and Francophone Caribbean Performance History - A Select Bibliography*. Theatre Research International, V15 N3, pp. 260-261, 1990.
- Annex V of MARPOL 73/78, Regulations for the Prevention of Pollution by Garbage from Ships*. International Maritime Organization.
- Antarctic Environmental Protection Act Of 1993*. Hearing Before The Subcommittee On Science, Space, and Technology, U.S. House Of Representatives, 103 Congress, First Session, N21, 1993.
- Apel, J.R. *Principles of Ocean Physics*. International Geophysics Series 38, Academic Press, 1-634, 1987.
- apRheinallt, T. *Impact of Sewage Sludge*. FWR Environment, U.K.
- Armi, L.A. *Effects of Variations in Eddy Diffusivity on Property Distributions in the Oceans*. Journal of Marine Research, V37, N3, pp. 515-530, 1979.
- Arntz W., M. Gorny. *Shrimp (Decapoda, Natantia) Occurrence and Distribution In The Eastern Weddell Sea, Antarctica*. Polar Biology, V11 N3, pp. 169-177, 1991.
- Atmospheric Transport Of Contaminants In The Mediterranean Region /IMO*. IMO/FAO/UNESCO/WHO/IAEA/UN/UNEP Joint Group of Experts on The Scientific Aspects of Marine Pollution. UNESCO/IMO/WHO/IAEA/UN/UNEP Joint Group of Experts on the Scientific Aspects of Marine Pollution GESAMP, Geneva: World Meteorological Organization, 1985.
- Atwood D., J. Hendee, A. Mendez. *An Assessment Of Global Warming Stress On Caribbean Coral Reef Ecosystems*. Bulletin of Marine Science, V51 N1, pp. 118-130, 1992.

- Augier, H. *Protected Marine Areas: The Example Of France: Appraisal and Prospects, Basis Of A Strategy For The Creation Of A European Network Of Marine Protected Areas In The Mediterranean Region*. Strasbourg Council of Europe, Croton, Publishing. Company, Distributor, N.Y., Manhattan, 1985.
- Ayrton, W.R., S. Oldridge. *Littoral Zones, Amenities, and Tourism*. Scarborough Borough Council, England, for the Institution of Environmental Health Officers, London, England.
- Bak R., A. Boldrin, G. Nieuwland, S. Rabitti. *Biogenic Particles and Nano Picoplankton In Water Masses Over The Scotia-Weddell Sea Confluence, Antarctica*. Polar Biology, V12 N2:219-224, 1992.
- Bak, R.P.M. *Variations In Coral Reefs in Response to Gradients Limiting Primary Production*. Netherlands Institute for Sea Research, The Netherlands.
- Ball, J., T. Reynolds. *Dispersion of Liquid Waste From a Moving Barge*. Journal WPCF, V48, N11, pp. 2541-2547, November, 1976.
- Balopoulos, E.T., A.E. James. *Surface Currents in the N.W. Aegean Sea (Greece) As Shown From the Movement of Driftcards*. Institute of Oceanographic and Fisheries Research, Department of Oceanography, University College, Singleton Park Swansea S A2 8PP, S. Wales (U.K.), VII Journees Etud. Pollutions, 1984.
- Baltic Sea Monitoring Symposium (1986 : Tallinn, Estonia)*. Baltic Sea Monitoring Symposium, Tallinn, USSR, 10-15 March 1986. Helsinki: Baltic Marine Environment Protection Commission, Helsinki Commission, 1986.
- Baltic Sea*. New Encyclopedia Britannica 15th Edition, Encyclopedia Britannica Inc., 1974.
- Baltic, Marine Environment Protection Commission. *Reception Of Wastes From Ships In The Baltic Sea Area A MARPOL 73/78 Special Area*. Baltic Sea Environment Proceedings, (28), pp. 84, 1989.
- Barrow 1990-1995 Sewage Discharge Monitoring Report*. National Rivers Authority, Outfall to Walney Channel, United Kingdom, 1990-1995.
- Bathmann U., G. Fischer, P. Muller, D. Gerdes. *Short-Term Variations In Particulate Matter Sedimentation Off Kapp Norvegia, Weddell Sea, Antarctica - Relation To Water Mass Advection, Ice Cover, Plankton Biomass and Feeding Activity*. Polar Biology, V11 N3, pp. 185-195, 1991.
- Beavers, J.A., G.H. Koch, W.E. Berry. *Corrosion of Metals in Marine Environments*. MCIC Report 86-50, Battelle Columbus Division, Columbus, Ohio, July, 1986.
- Becker, C.D. *Regional Survey Of Marine Biota For Bioassay Standardization Of Oil And Oil Dispersant Chemicals : Research Report*. American Petroleum Institute Committee on Environmental Affairs. Richland, Washington: Battelle Pacific Northwest Laboratories, 1973.

- Becker, C.D., J.A. Lichatowich, M.J. Schneider, J.A. Stand. *Regional Survey of Marine Biota for Bioassay Standardization of Oil and Oil Dispersing Chemicals*. American Petroleum Institute Committee on Environmental Affairs, API Publication N4167, April, 1973.
- Beckmann, M., G. Liebezeit. *Organic Carbon In The North Sea In May/June 1986: Distribution and Controlling Factors*. Scientific, Common Problems Of The Environment United, Nations Environment Programme, Hamburg (FRG) International Carbon Unit. Biogeochemistry and Distribution of Suspended Matter In The North Sea And Implications-To-Fisheries-Biology, N65 pp. 99-116, 1986.
- Benner, R., A.E. Maccubbin, R.E. Hodson. *Temporal Relationship Between the Deposition and Microbial Degradation Of Lignocellulosic Detritus in a Georgia Salt Marsh and the Okefenokee Swamp*. Microbiology Ecology, V12, pp. 291-298, 1986.
- Benner, R., M.A. Moran, R.E. Hodson. *Effects of pH and Plant Source on Lignocellulose Biodegradation Rates In Two Wetland Ecosystems, The Okefenokee Swamp and A Georgia Salt Marsh*. Limnology Oceanography V30, pp. 489-499, 1985.
- Benner, R., R.E. Hodson. *Microbial Degradation Of The Leachable and Lignocellulosic Components Of Leaves And Wood From Rhizophora Mangle In A Tropical Mangrove Swamp*. Marine Ecology Progress Series, V23, pp. 221-230, 1985.
- Benner, R., S.Y. Newell, A.E. Maccubbin, R.E. Hodson. *Relative Contributions Of Bacteria and Fungi To Rates of Degradation of Lignocellulosic Detritus in Salt Marsh Sediments*. Applied and Environmental Microbiology, V48, pp. 36-40, 1984.
- Benninghoff, W.S., W.N. Bonner. *Man's Impact on The Antarctic Environment: A Procedure For Evaluation Impacts From Scientific and Logistic Activities*. Scientific Committee on Antarctic Research, Scott Polar Research Institute, 1985.
- Berkman P., M. Nigro. *Trace Metal Concentrations In Scallops Around Antarctica - Extending The Mussel Watch Programme To The Southern Ocean*. Marine Pollution Bulletin, N6:322-323 V24, 1992.
- Bermuda Biological Station For Research. *Primary Productivity In The Tropical North Atlantic Off Barbados And The Caribbean Sea Off Jamaica*, Final Report/Bermuda Biological Station, St. George's West, Bermuda, 1965.
- Berner, R.A. *Early Diagenesis, A Theoretical Approach*. Princeton University Press, Princeton, New Jersey, 1980.
- Bernhard, M. *Heavy Metals and Chlorinated Hydrocarbons in the Mediterranean*. VIII.
- Bertels, A. *R/V Cruise No. 75 , 16 Oct.-19 Dec. 1967: Summary Cruise Report Track Charts, Bathymetry, And Location Of Observations*. North Atlantic Ocean-Caribbean Sea, Lesser Antillean Arc And Mid-Atlantic Ridge Survey.
- Bethoux J., P. Courau, E. Nicolas, D. Ruizpino. *Trace Metal Pollution In The Mediterranean Sea*. Oceanologica Acta, V13 N4, pp. 481-488, 1990.

- Bethoux, J.P. *Phosphorus And Nitrogen In The Mediterranean Sea: Budgets and Potential Fertility*. Marine Chemistry, 10(2), pp. 141-158, 1981.
- Beusekom, J., E.E. Von. *Distribution Of Dissolved Aluminum In The North Sea: Influence Of Suspended Matter*. Eds. S. Kempe, G. Liebezeit, V. Dethlefsen, U. Harms, Scientific Committee on Problems of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry and Distribution of Suspended Matter in The North Sea And Implications To Fisheries Biology, N65 pp. 117-136, 1988.
- Bianchi F., A. Boldrin, F. Cioce, G. Dieckmann, H. Kuosa, A.M. Larsson, E.M. Nothig, P.I. Sehlstedt, G. Socal, E.E. Syvertsen. *Phytoplankton Distribution In Relation To Sea Ice, Hydrography and Nutrients In The Northwestern Weddell Sea In Early Spring 1988 During EPOS*. Polar Biology, V12 N2, pp. 225-235, 1992.
- Bidigare R., M. Ondrusek, J. Brooks. *Influence Of The Orinoco River Outflow On Distributions Of Algal Pigments In The Caribbean Sea*. Journal Of Geophysical Research-Oceans, V98 pp. 2259-2269, 1993.
- Bingel, F., D. Avsar, M. Unsal *A Note on Plastic Materials In Trawl Catches In The Northeastern Mediterranean*. Meeresforschung (Reports on Marine Research) 31(3-4), pp. 227-233. Institute of Marine Science, Erdemli-Icel, Turkey, 1987.
- Birkhoff, G., Zarantonello. *Jets, Wakes and Cavities*. Academic Press, New York, pp. 353, 1957.
- Bjornsen P., J. Kupařinen. *Growth and Herbivory By Heterotrophic Dinoflagellates In The Southern Ocean, Studied By Microcosm Experiments*. Marine Biology, V109 N3, pp. 397-405, 1991.
- Blaxter, J.H.S., J.R. Hunter. *The Biology of the Clupeoid Fishes*. Off Print from Advances in Marine Biology, V20, pp.1-223, 1982.
- Bleaching Of Coral Reefs In The Caribbean*. United States Congress Senate Committee on Appropriations Subcommittee on Commerce, Justice, State, The Judiciary, and Related Agencies. Hearing Before a Subcommittee of The Committee on Appropriations, United States Senate, One Hundredth Congress, First Session: Special Hearing, Washington, 1988.
- Blough N., O. Zafiriou, J. Bonilla. *Optical Absorption Spectra of Waters From The Orinoco River Outflow-Terrestrial Input of Colored Organic Matter to The Caribbean*. Journal of Geophysical Research-Oceans, V98 Nc2, pp. 2271-2278, February, 1993.
- Boltovskoy D., V. Alder, F. Spinelli. *Summer Weddell Sea Microplankton - Assemblage Structure, Distribution and Abundance, With Special Emphasis On The Tintinnina*. Polar Biology, V9 N7, pp. 447-456, 1989.
- Boltovskoy D., V. Alder. *Microzooplankton and Tintinnid Species-Specific Assemblage Structures - Patterns Of Distribution and Year-To-Year Variations In The Weddell Sea (Antarctica)*. Journal Of Plankton Research, V14 N10, pp. 1405-1423, 1992.

- Boston Humus 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Boston Humus Treatment Works, United Kingdom, 1992-1994.
- Bourne, W.R.P. *Oil and Garbage in the Gulf*. Marine Pollution Bulletin, V20, N2, 1989.
- Boysenennen E., W. Hagen, G. Hubold, U. Piatkowski. *Zooplankton Biomass In The Ice-Covered Weddell Sea, Antarctica*. Marine Biology, V111 N2, pp. 227-235, 1991.
- Branica, M., G. Kniewald. *International Symposium on The Chemistry of The Mediterranean (10th: 1988: Primosten, Croatia) Physico-Chemical Characteristics of The Aquatic System: 10th International Symposium 'Chemistry Of The Mediterranean,' 4-12 May, 1988, Primosten, Yugoslavia*. Amsterdam, New York: Elsevier, 1989.
- Branica, M., G. Kniewald. *International Symposium On The Chemistry Of The Mediterranean (9th: 1986: Primosten, Croatia) Reliability of Environmental Analytical Data: Ix International Symposium 'Chemistry Of The Mediterranean,' 1-7 May, 1986, Primosten, Yugoslavia*. Amsterdam, Netherlands, 1987.
- Breslin, V.T., S.E. Tisdell. *Thermoplastic Stabilization of MSW Combustor Ash*. Journal of Environmental Engineering, V120, N1, January/February, 1994.
- Breslin, V.T. *Degradation of Starch-Plastic Composites in a Municipal Solid Waste Landfill*. Journal of Environmental Polymer Degradation, V1, N2, 1993.
- Breslin, V.T., F.J. Roethel, V.P. Schaeperkoetter. *Physical and Chemical Interactions of Stabilized Incineration Residue with the Marine Environment*. Marine Pollution Bulletin, V19, N11B, pp. 628-632, 1988.
- Brey T., J. Gutt. *The Genus Sterechinus (Echinodermata, Echinoidea) On The Weddell Sea Shelf and Slope (Antarctica) - Distribution, Abundance and Biomass*. Polar Biology, V11 N4, pp. 227-232, 1991.
- Brick, R.M., J.J. Daly, E.L. Koehler, A.G. Skibbe. *The Selection of Tin Coatings for Steel Containers*. Ed. T. Lyman Metals Handbook, V1, pp. 1133-1141, Ohio, 1961.
- Brisou, J. *An Environmental Sanitation Plan for the Mediterranean Seaboard*. Pollution and Human Health, World Health Organization, Geneva, 1976.
- Brockmann U., L. Rwpm, H. Postma. *Cycling of Nutrient Elements in The North Sea*. Netherlands Journal of Sea Research, V26 N2-4, pp. 239-264, 1990.
- Broman, D. *Transport and Fate of Hydrophobic Organic Compounds in The Baltic Aquatic Environment : Polycyclic Aromatic Hydrocarbons, Polychlorinated Dibenzodioxins and Dibenzofurans*. University of Stockholm, Stockholm, 1990.
- Brooks, N.H. R.G. Arnold, R.C.Y. Koh, G.A. Jackson, W.K. Faisst. *A Research Plan for Deep-Ocean Disposal of Sewage Sludge*. Robert E. Krieger Publishing Company, Malabar, Florida, 1987.

- Brügmann L, P. Bernard, R.Vangrieken. *Geochemistry of Suspended Matter From The Baltic Sea .2. Results of Bulk Trace Metal Analysis By AAS*. Marine Chemistry, V38 N3-4 pp. 303-323, 1992.
- Brunnee J. *International Conference On Ecology And Law In The Baltic Sea Area - Sources And Developments, Held In The University-Of-Latvia, Riga, Latvia, USSR, During 26-31 August 1990*. Environmental Conservation, V17 N4 pp. 375-375, 1990.
- Budzichawski, T. *Tan Book*. United States Navy, Carderock Division Naval Surface Warfare Center Annapolis Detachment Solid Waste Management Branch, 1995.
- Burmester, H. *Vessel Source Pollution: The Integration of International and Domestic Responses in the Search for an Effective Legal Framework*. Oceans Policy Study 1:6, 1978.
- Burnham 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Burnham Treatment Works, United Kingdom, 1992-1994.
- Burns, D.A. *Hydrographic Measurements In The Grenada Basin, Southeastern Caribbean Sea*. 1981.
- Burns, D.A., M. Car. *Current Meter Data Report For The Eastern Part Of The Caribbean Sea*. 1975.
- Burton J., M. Althaus, G. Millward, A. Morris. *Processes Influencing The Fate Of Trace Metals In The North Sea*. Philosophical Transactions of The Royal Society of London Series A-Physical Sciences and Engineering, V343 N1669, pp. 557-568, 1993.
- Busch, H.J. *Exports of Wastes By Sea*. Gefaehrliche Ladung, V34, N1, pp. 9-10, 1989.
- Bythell J., C. Sheppard. *Mass Mortality of Caribbean Shallow Corals*. Marine Pollution Bulletin, V26 N6, pp. 296-297, 1993.
- Cadee G. *Organic Carbon In The Upper Layer and Its Sedimentation During The Ice-Retreat Period In The Scotia-Weddell Sea*, 1988. Polar Biology, V12 N2, pp. 253-259, 1992.
- Calamari, D. *Selected Bibliography On Studies And Research Relevant To Pollution In The Mediterranean = Bibliographie Selectionnee Sur Les Etudes Et Recherches Relatives A La Pollution En Mediterranee*. Rome: Food and Agriculture Organization of the United Nations, 1977.
- Capuzzo, J.M., D.R. Kester. *Oceanic Processes in Marine Pollution Volume I, Biological Processes and Wastes in the Ocean*. Robert E. Krieger Publishing Company, Malabar, Florida, U.S., 1987.
- Carde, K.L., R.G. Steward. *Optical Means Of Measuring Particle-Sea Interactions In The Caribbean Sea And Gulf Of Mexico*. 1983.
- Caribbean Geological Evolution: Report of A Workshop To Define Caribbean Geological Problems, Needed Investigations, and Initiatives For Ocean Drilling*, 1987.

- Caribbean Sea*. New Encyclopedia Britannica 15th Edition, Encyclopedia Britannica Inc., 1974.
- Caribbean Study Project Working Papers And Selection From Dialogue At Preparatory Conference*, Jamaica, October 1972, [Malta]: Published For The International Ocean Institute By Malta University Press, 1974.
- Caribbean Zooplankton*. Office Of Naval Research, Department of The Navy U.S. Government Printing Office Washington, 1976.
- Carlson, Herbert. *Quality status of the North Sea*. Report, International Conference on the Protection of the North Sea, Bremen, 1984.
- Cederwall H., R. Elmgren. *Biological Effects Of Eutrophication In The Baltic Sea, Particularly The Coastal Zone*. *Ambio*, V19 N3 pp. 109-112, 1990.
- Cevasco, F.M. *Coastal Characteristics and Their Effect on Tanker Discharge Operations*. U.S. Army Mobility Equipment Research and Development Command, Fort Belvoir, Virginia, January, 1977.
- Chadwick, D.B., M.H. Salazar. *Integrated Technologies for Monitoring the Marine Environment*. Ocean Proceedings 1991 Conference, pp. 343-350, 1991.
- Charnock, H., K.R. Dyer, J.M. Huthnance, P.S. Liss, J.H. Simpson, P.B. Tett. *Understanding the North Sea System*. Chapman & Hall for the Royal Society, 1994.
- Chelmsford 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Chelmsford Treatment Works, United Kingdom, 1992-1994.
- Chester, R. *Marine Geochemistry*. Unwin Hyman Ltd, London, England, 1990.
- Chrást, R.J., H. Rai. *Ectoenzyme Activity And Bacterial Secondary Production In Nutrient-Impooverished And Nutrient-Enriched Freshwater Mesocosms*. *Microbial Ecology*, V25, pp. 131-150, 1993.
- CICAR Ichthyoplankton Workshop (1974 : Mexico City, Mexico) Ichthyoplankton: Report Of The CICAR Ichthyoplankton Workshop, Mexico City, 17-26 July 1974*. Paris, France: Unesco, 1975.
- City of Tampa 1994 Annual Sewage Discharge Monitoring Report*. City of Tampa Sewage Treatment Works, Florida, 1994.
- Clark, C.D. *Satellite Remote Sensing of Marine Pollution*. *Int. J. Remote Sensing*, V14, N16, pp. 2985-3004, 1993.
- Clark, C.D. *Satellite Remote Sensing for Marine Pollution Investigations*. *Marine Pollution Bulletin*, Vol. 26, No. 7, pp. 357-368, 1993.
- Clark, R. B. *Marine Pollution, Third Edition*. Clarendon Press, Oxford, New York, 1992.
- Cleethorpes Southern 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Cleethorpes Southern Treatment Works, United Kingdom, 1992-1994.

- Cliff Quay 1992-1994 Sewage Discharge Monitoring Report.* National Rivers Authority, Cliff Quay Treatment Works, United Kingdom, 1992-1994.
- Coast Guard Oil, Noxious Liquid Substances, Garbage, Municipal or Commercial Waste and Ballast Water Regulations.* The Bureau of National Affairs, Inc., February, 1995.
- Colchester 1992-1994 Sewage Discharge Monitoring Report.* National Rivers Authority, Colchester Treatment Works, United Kingdom, 1992-1994.
- Colwell, R.R. *Microbiological Effects of Ocean Pollution.* University of Maryland, USA.
- Committee on Merchant Marine and Fisheries, House of Representatives One Hundred Third Congress First Session. *The Effectiveness of the Coast Guard's Efforts to Enforce Laws Which Prohibit the Dumping of Garbage into the Oceans and Examine the Two Publicized Cases of Illegal Dumping.* U.S. Government Printing Office, 1993.
- Common Measures Adopted By The Contracting Parties To The Convention For The Protection Of The Mediterranean Sea Against Pollution.* Athens: UNEP, 1990.
- Connor, M.S. *At-Sea Incineration: Up in Smoke?* Research Fellow Interdisciplinary Programs in Health, Harvard School of Public Health.
- Contributions to Antarctic Research I. V50 Antarctic Series,* American Geophysical Union, Washington D.C., 1990.
- Convention For The Protection And Development Of The Marine Environment Of The Wider Caribbean Region (1983)* Convention For The Protection And Development Of The Marine Environment Of The Wider Caribbean Region: U.S.G.P.O., 1984.
- Cooperative Investigation Of The Caribbean And Adjacent Regions. Guide To CICAR Standard Observations : With Special Reference To CICAR Survey Months And Standard Sections / Compiled By The Operations Coordinator Of CICAR, Tom Van 'T Hof.* Curacao, Netherlands Antilles: CICAR, 1974.
- Cooperative Research Report. *A Review of the Past and Present Measurements of Selected Trace Metals in Sea Water in the OSLO Commission and ICNAF (NAFO) Areas.,* 1980.
- Cooperative Research Report. *The ICES Coordinated Monitoring Programme for Contaminants in Fish and Shellfish, 1978 and 1979 and Six-Year Review of ICES Coordinated Monitoring Programs,* 1984.
- Coral Reefs, Seagrass Beds, and Mangroves: Their Interaction in the Coastal Zones of the Caribbean.* Report of a Workshop, West Indies Laboratory, St. Croix, U.S. Virgin Islands, Unesco, 1983.
- Cormaci M., G. Furnari, B. Scammacca. *The Benthic Algal Flora Of Terra Nova Bay (Ross Sea, Antarctica).* Botanica Marina, V35 N6, pp. 541-552, 1992.
- Cota G., S. Kottmeier, D. Robinson, W. Smith. *Bacterioplankton In The Marginal Ice Zone Of The Weddell Sea - Biomass, Production and Metabolic Activities During Austral*

- Autumn. Deep-Sea Research Part A-Oceanographic Research Papers*, V37 N7, pp. 45-1167, 1990.
- Cripps G. *Baseline Levels Of Hydrocarbons In Seawater Of The Southern Ocean - Natural Variability and Regional Patterns*. *Marine Pollution Bulletin*, V24 N2, pp. 109-114, 1992.
- Cripps G. *Hydrocarbons In The Seawater and Pelagic Organisms Of The Southern Ocean*. *Polar Biology*, V10 N5, pp. 393-402, 1990.
- Cruz G., V. Lopez, C. Sosa. *Pollution By Solid Wastes Carried By Marine Currents To The Caribbean Coast Of Honduras*. *Revista De Biologia Tropical*, V38 N2, pp. 339-342. Language: Spanish, 1990.
- Cruz, G.A., V. Lopez, C. Sosa. *Contaminacion por Desechos Solidos Llevados por Corrientes Marinas a la Costa Caribena de Honduras*. Departamento de Biologia, Tegucigalpa, Honduras, 1990.
- Csanady, G.T, J.H. Churchill. *Environmental Engineering Analysis of Potential Dumpsites*. Robert E. Krieger Publishing Company, Malabar, Florida, 1987.
- Csanady, G.T. *Accelerated Diffusion in the Skewed Shear Flow of Lake Currents*. *J. Geophysics Research*, 71, pp. 411-420, 1966.
- Csanady, G.T. *An Analysis of Dumpsite Diffusion Experiments: Ocean Dumping of Industrial Wastes*. Woods Hole Oceanographic Institution Woods, Hole, Plenum Press, New York and London, pp. 109-129, 1981.
- Csanady, G.T. *An Analysis of Dumpsite Diffusion Experiments*. Eds. B.H. Ketchum, D.R. Kester, P.K. Park, Plenum Press, New York & London, 1979.
- Csanady, G.T. *Turbulent Diffusion in the Environment*. *J. Atmospheric Science*, 26, pp. 414-426, 1973.
- Cubit, J.D., et al. *Hydrographic And Meteorological Studies Of A Caribbean Fringing Reef At Punta Galeta, Panama: Hourly And Daily Variations For 1977-1985*. Washington, D.C.: Smithsonian Institution Press, 1988.
- Cuomo, V., V. Pane, P. Contesso. *Marine Litter: Where does the Problem Lie?*. Research Laboratory, Ciba-Geigy A/S, Torre Annunziata, Naples, Italy, 1988.
- Daan, N., P. Bromley, J. Hislop, N. Nielsen. *Ecology Of North Sea Fish*. *Netherlands Journal Of Sea Research*, V26 N2-4, pp. 343-386, 1990.
- Dahl, J.B., O. Tollan. *Measurement of Dilution of Tank Waste Water in the Wake of M/T "Esso Bergen"*. Institute for Atomic Energy, Kjeller, Norway, 1972.
- Dahl, J.B., U. Haagensen, C. Qvenild, O. Tollan. *Dilution in the Ship's Wake of Tank Washings Released From M/T "Esso Slagen"*. Institute of Atomic Energy, Kjeller, Norway, 1973.

- Dahl, J.B., U. Haagensen, C. Qvenild, O. Tollan. *Measurements of the Dilution of Tank Washings in the Wake of M/T "Esso Slagen"*. Institute of Atomic Energy, Kjeller, Norway, 1975.
- Data Profiles For Chemicals For The Evaluation Of Their Hazards To The Environment Of The Mediterranean Sea*. Franklin Institute VI, Geneva: International Register of Potentially Toxic Chemicals, United Nations Environment Programme, 1978.
- Davaki, A. *Directory Of Mediterranean Marine Environmental Centres*. MAP Technical Reports Series N65, Athens, UNEP, 1992.
- Davidson, L., G. Kristina. *Special Area Status for the Wider Caribbean Region Under Annex I of the International Convention for the Prevention of Pollution From Ships*. U.S. Coast Guard, 1989.
- Davidson, S. *Atmospheric Depositions*. International Journal of Estuarine and Coastal Law.
- Davidson, S. *Atmospheric Depositions. The North Sea: Perspectives On Regional Environmental Cooperation*. Eds. D. Freestone, T. Ijlstra, V5, N1-3 pp. 288-299, 1990.
- Davis, R. *Sea Surface Films*. In a Report to the MITRE Corporation, JSR-86-105, McLean, Virginia, November, 1986.
- Davyhulme 1990-1995 Sewage Discharge Monitoring Report*. National Rivers Authority, Davyhulme, Sampled by Rivers, United Kingdom, 1990-1995.
- Day, R.H., D.G. Shaw. *Patterns in the Abundance of Pelagic Plastic and Tar in the North Pacific Ocean, 1976-1985*. Marine Pollution Bulletin, V18 N6B, pp. 311-316, 1987.
- De Armas, J.D. *Pelagic Tar in the Western Mediterranean*. Instituto Espanol de Oceanografia, Laboratorio de Caraias, Islas Canarias, VII Jounees Etud. Pollutions, 1984.
- De Bievre A.F.M. *Shipping and the North Sea Environment*. Werkgroep Noordzee.
- De Flora, S., M. Bagnasco, P. Zanicchi. *Genotoxic, Carcinogenic, and Teratogenic Hazards in The Marine Environment, With Special Reference To The Mediterranean Sea*. Mutation Research, V258 N3, pp. 285-320, 1991.
- De Wolfe, P., Zijlstra, J.J. *The Ecosystem*. Eds W. Salomons, B.L. Bayne, E.K. Duursma & U. Forstner, North Sea Pollution, pp. 118-151, Berlin, Springer-Verlag, 1988.
- Defense Mapping Agency. *Ocean Basin Environment. Sailing Directions (Planning Guide) for the North Atlantic Ocean*. Defense Mapping Agency, 3rd Edition, 1988.
- Defense Mapping Agency. *Ocean Basin Environment. Sailing Directions (Planning Guide) for the North Sea and Baltic Sea*. Defense Mapping Agency, 3rd Edition, 1990.
- Defense Mapping Agency. *Ocean Basin Environment. Sailing Directions (Planning Guide) for Antarctica*. Defense Mapping Agency, 2nd Edition, 1992.

- Defense Mapping Agency. *Ocean Basin Environment. Sailing Directions (Planning Guide) for the Mediterranean*. Defense Mapping Agency, 5th Edition, 1991.
- Dehairs F., N. Stroobants, L. Goeyens. *Suspended Barite As A Tracer Of Biological Activity In The Southern Ocean*. Marine Chemistry, V35 N1-4, pp. 399-410, 1991.
- Dekker, L. *Management Of Toxic Materials In An International Setting : A Case Study Of Cadmium In The North Sea*. Rotterdam, Netherlands, Brookfield, VT, USA: A.A. Balkema, 1987.
- Delft Hydraulics Laboratory. *Dilution of Liquids Discharged From A Ship*. Report M 1312. Delft Hydraulics Laboratory, Delft, The Netherlands, 1975.
- Delille D. *Marine Bacterioplankton At The Weddell Sea Ice Edge, Distribution Of Psychrophilic and Psychrotrophic Populations*. Polar Biology, V12 N2, pp. 205-210, 1992.
- Delvigne, G.A.L. *Experiments on The Dilution Capacity of Wakes From Tankers Dumping In The North Sea*. Malabar, Krieger Publishing Company, pp. 235, Florida, 1987.
- Dethlefsen V. *Fish In The Polluted North Sea*. Dana-A Journal of Fisheries and Marine Research, V8:109-129, 1989.
- Devine, M. *Analysis of Mixing and Dispersion of Industrial Wastes Dumped into the Deep Sea*. Oceanic Processes in Marine Pollution, Robert E. Krieger Publishing Co., V2, pp. 63-70, Malabar, Florida, 1979.
- Dexter, S.C. *Effect of Variation in Sea Water Upon the Corrosion of Aluminum*. Corrosion, V36, pp. 423-432, 1980.
- Dietrich S. ed. *Scientific Seminar On Antarctic Ocean Variability and Its Influence On Marine Living Resources, Particularly Krill (1987: Paris, France) Antarctic Ocean and Resources Variability*. [Scientific Seminar On Antarctic Ocean Variability and Its Influence On Marine Living Resources, Particularly Krill, Paris, 2-6 June 1987] Berlin, New York: Springer-Verlag, 1988.
- Dijkstra F. *North Sea Pollution - Technical Strategies For Improvement*. Proceedings Of The Iawprc International Conference, Held In Amsterdam, The Netherlands, 10-14 September 1990 - Preface. Water Science And Technology, V24 N10:R9-R9, 1991.
- Disposal*. United Press International.
- Dixon, T. R. *Operational Discharges From Ships And Platforms [Garbage, Packaged Dangerous Or Harmful Goods And Pyrotechnics]*. pp.45 47, 1990 North Sea Report. London: The Marine Forum for Environmental Issues, pp.164, 1990.
- Dixon, T.R., T.J. Dixon. *Marine Litter Surveillance*. Marine Pollution Bulletin, V14, N4, pp. 145-148, 1983.
- Donohue, G.L., J.W. Hoyt. *Harbor Pollution From Large Ships*. Fleet Engineering Department, Naval Undersea Center, San Diego, California, Oct., 1973.

- Dorten W., F. Elbazpoulichet, L. Mart, J. Martin. *Reassessment Of The River Input of Trace Metals Into The Mediterranean Sea*. *Ambio*, V20 N1, pp. 2-6, 1991.
- Drake, J., S. Gill, K. Bayer. *Technical Evaluation (TECHEVAL) Test Report for the Large Pulper Installed Onboard the U.S.S. George Washington (CVN 73)*. Carderock Division, Naval Surface Warfare Center, CARDIVNSWC-TM-63-94/4, Washington. D.C., 1994.
- Du Bois, P.B., P. Guegueniat, R. Gandon, R. Leon, Y. Baron. *Percentage Contribution of Inputs From the Atlantic, Irish Sea, English Channel and Baltic into the North Sea During 1988: A Tracer-Based Evaluation Using Artificial Radionuclides*. *Netherlands Journal of Sea Research*, 1993.
- Dubinsky, Z. *Coral Reefs*. Elsevier Science Publishers, Amsterdam, 1990.
- Ducklow, H.W., D.L. Kirchman, G.T. Rowe. *Production and Vertical Flux of Attached Bacteria In The New York Bight As Studied With Floating Sediment Traps*. *Applied Environmental Microbiology*, V43, pp. 769-776, 1982.
- Duckworth, D., D. Lapota, D.E. Rosenberger, J.H. Grovhoug. *Bioluminescence, Chlorophyll Fluorescence, and Phototaxis Inhibition in the Dinoflagellate, Gongaulax Polyedra*. *Bioluminescence and Chemiluminescence*, John Wiley & Sons, Chichester, pp. 540-543, 1994.
- Durst, C.S. *The Relationship Between Current and Wind*. *Quart. J. R. Met. Society*, V50, pp. 113, London, 1924.
- Duursma, E.K. *Assessment of Environmental Impact of Nutrients*. Netherlands Institute for Sea Research, 1987.
- Earthscan. *Environmental Wars & Environmental Refugees: The Political Background To The Cartagena Convention*. London: Earthscan, International Institute For Environment And Development, Series Title: Earthscan Press Briefing Document, No. 34B, 1983.
- Earthscan. *The Improbable Treaty: The Cartagena Convention and The Caribbean Environment*. London: International Institute for Environment and Development, 1983.
- Effects of Pollution on Marine Ecosystems, Blanes, Spain, 7-11 October 1985 = Communications Presentees a la Reunion FAO/PNUE sur les Effets de la Pollution Sur FAO/UNEP Meeting on the Effects of Pollution on Marine Ecosystems (1985: Blanes, Spain) Papers Presented at the FAO/UNEP Meeting on the Rome: Food and Agriculture Organization of the United Nations, 1987.*
- Ehlers, P. *Environmental Protection In Shipping What Is Expected Of Europe*. DTSCH. HYDROGR. Z. 1991-92. V44, N2, pp. 107-111, 1991.
- Eisma D. *Transport and Deposition of Suspended Matter in The North Sea and The Relation To Coastal Siltation, Pollution, and Bottom Fauna Distribution*. *Aquatic Sciences*, V3 N2-3, pp. 181-216, 1990.

- Ekau W. *Demersal Fish Fauna Of The Weddell Sea, Antarctica*. Antarctic Science, V2 N2, pp. 129-137, 1990.
- El Gindy, A.A.H., A.A. Hamed. *Wind Pattern and Expected Direction of Floating Pollutant Transport in the Nearshore Alexandria Water*. Physical Oceanography, Oceanography Department Faculty of Science, Alexandria University, Alexandria, Egypt, VII Jounees Etud. Pollutions Lucerne, 1984.
- Eliminating Floatable Refuse in the New York Area*. Joint Hearing Before the Subcommittee on Water Resources, Transportation, and Infrastructure and the Subcommittee on Superfund, Ocean, and Water Protection of the Committee on Environment and Public Works United States Senate One Hundred First Congress First Session on S.506, 1989.
- Elliot, D.H. *Contributions to Antarctic Research III*. V57 Antarctic Series; American Geophysical Union, Washington D.C., 1990.
- Elliot, D.H. *Contributions to Antarctic Research*. American Geophysical Union, Washington D.C., 1991.
- Elliott, A. J., Z. Li. *Thermocline Depths And Water Temperatures At Selected Sites In The N.W. European Shelf Seas*. Marine Pollution Bulletin, V22(6), pp. 282-286, 1991.
- Elmgren R. *Mans Impact On The Ecosystem Of The Baltic Sea - Energy Flows Today And At The Turn Of The Century*. Ambio, V18 N6 pp. 326-332, 1989.
- El-Sayed, S.Z. *Plankton*. Elsevier, New York, New York, 1977.
- Emelianov, E.M.Y. *Geochemistry and Sedimentology of The Mediterranean Sea*. Kluwer Academic Publishers, MA, U.S., 1986.
- England M. *On the Formation of Antarctic Intermediate and Bottom Water in Ocean General Circulation Models*. Journal Of Physical Oceanography, V22 N8, pp. 918-926, 1992.
- England M., J. Godfrey , A. Hirst, M. Tomczak. *The Mechanism For Antarctic Intermediate Water Renewal In A World Ocean Model*. Journal Of Physical Oceanography, V23 N7, pp. 1553-1560, 1993.
- Environmental Acoustics Atlas Of The Caribbean Sea And Gulf Of Mexico*. Marine Environment, V2, 1972.
- Environmental Agenda For The 1990's: A Synthesis of the Eastern Caribbean Country Environmental Profile Series: Antigua and Barbuda, Dominica, Grenada, St. Kitts and Nevis, St. Lucia, St. Vincent and The Grenadines*. St. Michael, Barbados: Caribbean Conservation Association, St. Thomas, V.I., Island Resources Foundation, 1991.
- Environmental Laws of the Commonwealth Caribbean Analysis and Needs Assessment*. A Study by the Caribbean Law Institute, 1992.
- Environmental Management*, March-April, V14 N2:221-227, 1990.

- Environmental-Acoustics Atlas Of The Caribbean Sea and Gulf Of Mexico*. Marine Acoustics, V1, 1986.
- Ernst, W., J.P. Boon, K. Weber. *Occurrence and Fate of Organic Micropollutants In The North Sea. Pollution Of The North Sea, An Assessment*. Eds. W. Salomons, B.L. Bayne, E.K. Duursma, U. Foerstner, pp. 284-299, 1988.
- Estrada M., M. Delgado. *Summer Phytoplankton Distributions In The Weddell Sea*. Polar Biology, V10 N6, pp. 441-449, 1990.
- Eutrophication in The Mediterranean Sea: Receiving Capacity and Monitoring of Long-Term Effects*. Report and Proceedings of A Scientific Workshop, Bologna, Italy, 2-6 March 1987 Sponsored By UNESCO Paris, 1988.
- Evans, R.J. *Death in Hamburg. Society and Politics in the Cholera Years 1830-1910*. Oxford University Press, Oxford, 1987.
- Everaarts J., C. Fischer. *The Distribution Of Heavy Metals (Cu, Zn, Cd, Pb) In The Fine Fraction Of Surface Sediments Of The North Sea*. Netherlands Journal Of Sea Research, V29 N4, pp. 323-331, 1992.
- Fabiano M., P. Povero, R. Danovaro. *Distribution and Composition Of Particulate Organic Matter In The Ross Sea (Antarctica)*. Polar Biology, V13 N8, pp. 525-533, 1993.
- Fahrbach E., G. Rohardt , G. Krause. *The Antarctic Coastal Current In The Southeastern Weddell Sea*. Polar Biology, V12 N2, pp. 171-182, 1992.
- Fahrbach E., M. Knoche, G. Rohardt. *An Estimate Of Water Mass Transformation In The Southern Weddell Sea*. Marine Chemistry, V35 N1-4, pp. 25-44, 1991.
- Farmer C., C. Moore, R. Zika, R. Sikorski. *Effects Of Low And High Orinoco River Flow On The Underwater Light Field of The Eastern Caribbean Basin*. Journal Of Geophysical Research-Oceans, V98, N2, pp. 2279-2288, 1993.
- Farmer, D., D. Lemon. *Dispersion of Dyed Sea-Water Discharged from Moving Vessels in Coastal Waters*. Pacific Marine Science Report, Institute of Ocean Sciences, Patricia Bay, Victoria, BC., 1975.
- Farrell, Jr., C.A. *Disposal of Marine Sewage: An Environmental Management Standard For Naval Ships*. Naval Undersea Center, San Diego, California, June, 1972.
- Federal Regulations. *Subpart A-Implementation of MARPOL 73/78*. Bureau of National Affairs, 1994.
- Feldman, M.H. *Trace Materials in Wastes Disposed to Coastal Waters-Fates, Mechanisms and Ecological Guidance and Control*. National Technical Information Service, Federal Water Quality Administration, Corvallis, Oregon, July, 1970.
- Ferm, R. *Integrated Management Of The Baltic Sea*. Marine Pollution Bulletin, V23, pp. 533-540, 1991.

- Figueiras F., F. Perez, Y. Pazos, A. Rios. *Light and Productivity Of Antarctic Phytoplankton During Austral Summer In An Ice Edge Region In The Weddell-Scotia Sea*. Journal of Plankton Research, V16 N3, pp. 233-253, 1994.
- Final Reports On Research Projects Dealing With The Effects of Pollutants on Marine Communities and Organisms = Rapports Finaux Sur Les Projets De Recherche Traitant Des Effets De Polluants Sur Les Communautés Et...* Athens: UNEP, 1993.
- First Baltic Sea Pollution Load Compilation*. Helsinki: Baltic Marine Environment Protection Commission, Helsinki Commission, 1987.
- Fischer G. *Stable Carbon Isotope Ratios Of Plankton Carbon and Sinking Organic Matter From The Atlantic Sector of The Southern Ocean*. Marine Chemistry, V35 N1-4, pp. 581-596, 1991.
- Fisheries and Pollution To The North Sea Biota*. Water Science and Technology, V24 N10, pp. 147-153, 1991.
- Fitzmaurice M. *The New Helsinki Convention On The Protection Of The Marine Environment Of The Baltic Sea Area*. Marine Pollution Bulletin, V26 N2, pp. 64-67, 1993.
- Fitzmaurice, M. *International Legal Problems of the Environmental Protection of the Baltic Sea*. Graham & Trotman, Norwell, Kluwer Academic Publishers, MA, U.S., 1992.
- Five Years of Solid Waste Planning in New York State*. Waste Management Research Report, V5, N2., 1993.
- Flag Fen 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Flag Fen Treatment Works, United Kingdom, 1992-1994.
- Floatable Pollution in the New Jersey-New York Harbor Complex*. Hearing Before the Subcommittee on Oversight and Investigations of the Committee on Merchant Marine and Fisheries House of Representatives, One Hundred First Congress First Session on Oversight of Legislation Governing the Problem of Pollution in the New York-New Jersey Harbor Complex and a Look at Long-Term Solutions to these Long-Term Problems, 1989.
- Flushing Times Of The North Sea*. Copenhagen, Denmark: International Council for the Exploration of the Sea, 1983.
- Focus on Incinerator Ash Research*. Waste Management Research Report, V1, N2, Spring, 1989.
- Focus on Marine Pollution*. Waste Management Research Report, V3, N2, 1991.
- Focus on Secondary Materials*. Waste Management Research Report, V2, N2, Summer, 1990.
- Focus on Source Reduction*. Waste Management Research Report, V3, N1, 1991.
- Focus on the Relevance of Research*. Waste Management Research Report, V4, N1, 1992.

- Foldvik A., J. Middleton, T. Foster. *The Tides Of The Southern Weddell Sea. Deep-Sea Research Part A. Oceanographic Research Papers*, V37 N8, pp. 1345-1362, 1990.
- Fontana, M.G., *Corrosion Engineering*. McGraw Hill, New York, 1986
- Francis, A. *The Caribbean Basin And Recent Developments In The Law Of The Sea, & Human Rights In The Commonwealth Caribbean: An International Relations Perspective*, St. Augustine, Republic Of Trinidad & Tobago: Institute Of International Relations, 1983.
- Frankenhoff, C., C.A. Matos, E. Towle, J. McEachern, L.T. Giulini. *Environmental Planning and Development In The Caribbean: Analysis and Needs Assessment*. Project Coordinator, Duke E.E. Pollard, Contributors, Editorial Universitaria, Universidad De Puerto Rico, 1977.
- Franklin, A. ed. *Monitoring And Surveillance of Non-Radioactive Contaminants In The Aquatic Environment and Activities Regulating The Disposal Of Wastes At Sea, 1988/89*. Ministry of Agriculture Fisheries and Food, Directorate of Fisheries Research, Aquatic Environment Monitoring Report, N26, pp. 90, 1989.
- Fransz H., J. Mommaerts, G. Radach. *Ecological Modeling of The North Sea*. Netherlands Journal of Sea Research, V28 N1-2, pp. 67-140, 1991.
- Freestone, D. *Protection of Marine Species and Ecosystems in the Wider Caribbean, the Protocol on Specially Protected Areas and Wildlife*. Marine Pollution Bulletin, V22, N12, pp. 579-581, 1991.
- Frost, B.W. *Effects of Size and Concentration of Food Particles on the Feeding Behavior of the Marine Planktonic Copepod Calanus pacificus*. Limnology Oceanography, 17:805-815, 1972.
- Fryxell G. *Marine Phytoplankton At The Weddell Sea Ice Edge - Seasonal Changes At The Specific Level*. Polar Biology, V10 N1, pp. 1-18, 1989.
- Furness, R.W., P.S. Rainbow. *Heavy Metals in the Marine Environment*. CRC Press, Boca Raton, Florida, 1990.
- Gabaldon A. *The Disposal of Toxic or Dangerous Waste In Latin-America and The Caribbean*. Marine Policy, V14 N3, pp. 254-258, 1990.
- Gable F., M. Affs, J. Gentile, D. Aubrey. *Global Climatic Issues In The Coastal Wider Caribbean Region*. Environmental Conservation, V17 N1, pp. 51-60, Spring, 1990.
- Gabrielides, G.P., A. Golik, L. Loizides, M.G. Marino, F. Bingel, M.V. Torreg Rossa. *Man Made Garbage Pollution on the Mediterranean Coastline. Environmental Management And Appropriate Use Of Enclosed Coastal Seas Emecs'-90*. CODEN: International Conference on the Environmental Management of Enclosed Coastal Seas '90: EMECS '90, Kobe, Hyogo Prefect. (Japan), 3-6 August 1990 Marine Pollution Bulletin, V23, 1991.

- Gabrieliadis, G.P. *Proceedings of the FAO/UNEP/IAEA Consultation Meeting on the Accumulation and Transformation of Chemical Contaminants by Biotic and Biotic Processes in the Marine Environment.*, Mediterranean Action Plan, UNEP, MAP Technical Reports Series N59, La Spezia, Italy, pp. 24-28, September, 1990.
- Garrison D., K. Buck . *Protozooplankton In The Weddell Sea, Antarctica - Abundance and Distribution In The Ice-Edge Zone.* Polar Biology, V9 N6, pp. 341-351, 1989.
- Garrison D., K. Buck. *The Biota Of Antarctic Pack Ice In The Weddell Sea and Antarctic Peninsula Regions.* Polar Biology, V10 N3, pp. 211-219, 1989.
- Garrity S., S. Levings. *Marine Debris Along The Caribbean Coast Of Panama.* Marine Pollution Bulletin, V26 N6, pp. 317-324, 1993.
- Gerdes D., M. Klages, W. Arntz, R. Herman, J. Galeron, S. Hain, *Quantitative Investigations On Macrobenthos Communities Of The Southeastern Weddell Sea Shelf Based On Multibox Corer Samples.* Polar Biology, V12 N2, pp. 291-301, 1992.
- Gerlach, S.A. *Nutrients-an Overview.* Institute for Marine Science, University of Kiel, Federal Republic of Germany.
- Geyer, R.A. *Marine Environmental Pollution, 2, Dumping and Mining,* Elsevier Scientific Publishing Company, New York, New York, 1981.
- Geyer, R.A. *Naturally Occurring Hydrocarbon Seeps In The Gulf Of Mexico and The Caribbean Sea.*
- Giam, C.S., H.J. Dou. *Strategies and Advanced Techniques for Marine Pollution Studies : Mediterranean Sea.* NATO Advanced Study Institute Berlin, New York: Springer-Verlag, 1986.
- Gibbs, R.J., M. Angelidis. *Effect of Sludge Digestion on Metal Segregation During Ocean Dumping.* Marine Pollution Bulletin, V20, N10, pp. 503-508, 1989.
- Giovanardi, F., R. Marchetti, G. Montanari, A Rinaldi, R. Barnard. *Eutrophication Problems in the Northern Adriatic Empirical Dispersion Model For Total Phosphorus.* Istituto Ricerca Sulle Acque, C.N.R. Brugherio, Milano, Italia, VII Journees Etud. Pollutions, Lucerne, 1984.
- Glas, P.C.G. *A North Sea Computational Framework For Environmental And Management Studies: An Application For Eutrophication and Nutrient Cycles/* P.C.G. Glas and T.A. Nauta.[Delft, Netherlands]: Delft Hydraulics, 1990.
- Glasby, G.P. *Antarctic Sector of the Pacific.* Elsevier, New York, New York, 1990.
- Glass Products and Production.* Encyclopedia Britannica Macropedia 15th Edition, V13, Encyclopedia Britannica Inc., 1974.
- Gleitz M., U. Bathmann, K. Lochte. *Build Up and Decline Of Summer Phytoplankton Biomass In The Eastern Weddell Sea, Antarctica.* Polar Biology, V14 N6, pp. 413-422, 1994.

- Goenaga C. *The State Of Coral Reefs in The Wider Caribbean*. Interciencia, V16 N1, pp. 12-20, 1991.
- Golchert, H.J. *Waste Disposal From The Point Of View Of Shipping Companies In Particular With Regard To The Situation Aboard*. Schiff Hafen Seewirtsch. 1990.V42, N12, pp. 12-19, 1990.
- Golik, A., 1934-*Tar Pollution In The Mediterranean Sea*. Prepared In Cooperation With IOC. Nairobi, Kenya: Oceans and Coastal Areas Programme Activity Centre, 1988.
- Golik, A., Y. Gertner. *Litter On The Israeli Coastline*. Marine Environmental Research, pp. 1-15, 1992.
- Gordon, A.L. *Oceanography of Antarctic Waters*. Lamont-Doherty Geological Observatory of Columbia University, Palisades, New York, 1971.
- Gramentz, D. *Involvement Of Loggerhead Turtle With The Plastic, Metal, And Hydrocarbon Pollution In The Central Mediterranean*. Marine Pollution Bulletin, V19, pp. 11-13, 1988.
- Graneli E., W. Graneli, M. Rabbani, N. Daugbjerg, G. Fransz, J. Cuzin-Roudy, V.A. Alder. *The Influence of Copepod and Krill Grazing On The Species Composition Of Phytoplankton Communities From The Scotia-Weddell-Sea - An Experimental Approach*. Polar Biology, V13 N3, pp. 201-213, 1993.
- Gray J., K. Clarke, R. Warwick, G. Hobbs. *Detection of Initial Effects of Pollution on Marine Benthos - An Example From The Ekofisk And Eldfisk Oilfields, North Sea*. Marine Ecology-Progress Series, SEP, V66 N3, pp. 285-299, 1990.
- Great Britain. *Department of The Environment. United Kingdom North Sea Action Plan, 1985-1995 : Presented To The Third North Sea Conference Held In The Hague, March 1990*. Department of The Environment, London, 1990.
- Greek Environmental Clean-Up Programme*. Marine Pollution Bulletin, V28, N11, Nov., 1994.
- Grenon, M., M. Batisse, eds. *Futures For The Mediterranean Basin: The Blue Plan*. Foreword By M.K. Tolba, Oxford University Press, New York, 1989.
- Griffin, O.W., R.D. Peltzer, A.M. Reed, R.F. Beck. *Remote Sensing of Surface Ship Wakes*. Naval Engineers Journal, pp. 245-258, May, 1992.
- Grimalt, J., J.M. Bayona, J. Albaiges. *Chemical Markers for the Characterization of Pollutant Inputs in the Coastal Zones*. Instituto de Quimica Bio-Organica, Jorge Girona Salgado, Barcelona, Spain, VII Journees Etud. Pollutions, pp. 18-26, 1984.
- Grogan, W.C., *Compiler. Input Of Contaminants To The North Sea From The United Kingdom*. Final Report Prepared for the Department of the Environment by the Institute of Offshore Engineering. Institute of Offshore Engineering, Heriot-Watt University, Edinburgh, 203pp. 1984 Institute of Offshore Engineering., Heriot-Watt Univ., Edinburgh, EH14 4AS, U.K., 1984.

- Grosch, C.E. *Turbulent Diffusion in a Stratified Fluid with Application to the Ocean*. Ocean and Atmospheric Science, Inc., NTIS, 1972.
- Grossmann S., G. Dieckmann. *Bacterial Standing Stock, Activity, and Carbon Production During Formation and Growth Of Sea Ice In The Weddell Sea, Antarctica*. Applied and Environmental Microbiology, V60 N8, pp. 2746-2753, 1994.
- Groups Ask EPA to Convene Conference on Runoff Pollution in Mississippi, Gulf*. Environmental Reporter, Current Developments, pp. 1839, 1994.
- Gruener, K., R. Reuter, H. Smid. *A New Sensor System For Airborne Measurements Of Maritime Pollution and of Hydrographic Parameters*. Geojournal, V24, N1, pp. 103-117, 1991.
- Guidelines for the Designation of Special Areas and the Identification of Particularly Sensitive Sea Areas*. Draft Assembly Resolution Annex 17, MEPC 31/21.
- Guruswamy, L. *Environmental Management In A North Sea Coastal Zone: Law, Institutions and Policy*. Natural Resources Journal, V25, Albuquerque, New Mexico: University of New Mexico, School of Law, 1985.
- Gutt J. *On The Distribution and Ecology Of Holothurians In The Weddell Sea (Antarctica)*. Polar Biology, V11 N3, pp. 145-155, 1991.
- Guzman H., C. Jimenez. *Contamination of Coral Reefs By Heavy Metals Along The Caribbean Coast of Central America - (Costa-Rica And Panama)* Marine Pollution Bulletin, V24 N11, pp. 554-561, 1992.
- Guzman H., I. Holst. *Effects Of Chronic Oil-Sediment Pollution on The Reproduction of The Caribbean Reef Coral Siderastrea-Siderea*. Marine Pollution Bulletin, V26 N5, pp. 276-282, 1993.
- Guzzella, L., A. De Paolis. *Polycyclic Aromatic Hydrocarbons in Sediments of the Adriatic Sea*. Marine Pollution Bulletin, V28, N3, pp. 159-165, 1994.
- Haas, P.M. *Saving The Mediterranean: The Politics of International Environmental Cooperation*. Columbia University Press, New York, 1990.
- Halewood 1990-1995 Sewage Discharge Monitoring Report*. National Rivers Authority, Halewood Primary Tank Effluent, United Kingdom, 1990-1995.
- Hallberg R. *Environmental Implications of Metal Distribution In Baltic Sea Sediment*. Ambio, V20, N7 pp. 309-316, 1991.
- Halminski, S.J. *Temperature, Salinity, Oxygen, and Phosphate in Waters Off Eastern Central America and Northern South America*. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Environmental Data Service, Key To Oceanographic Records Documentation, N4, Washington, 1975.
- Hammond, R.R., S.A. Jenkins, J.S. Cleveland, J.C. Talcott, A.L. Heath, J. Wasyl, S.G. Goosby, K.F. Schmitt, L.A. Levin. *Coastal Water Clarity Modeling*. SAIC, Technical Report 01-1349-03-4841-000, pp. 491, 1995.

- Hanson, M.B., P. Aguilar. *Marine Debris Bibliography*. NOAA's Marine Debris Information Office, 1989.
- Harden C. *Water Resources Management In Latin-America and The Caribbean - Lee, T.* Professional Geographer, V43 N4, pp. 545-545, 1991.
- Hardisty, J. *The British Seas: An Introduction To The Oceanography and Resources of The North West European Continental Shelf*. London: Routledge, pp. 272, 1990.
- Harms, J. *Marine Plastic Litter as an Artificial Hard Bottom Fouling*. Biologische Anstalt Helgoland, Hamburg, Germany, 44, pp. 503-506, 1990.
- Harris County 1994 Annual Sewage Discharge Monitoring Report*. Harris County Wastewater Treatment Plant, Houston Texas, 1994.
- Hazelworth, J.B. *Oceanographic Conditions In The Caribbean Sea During The Summer of 1971*. Boulder, Colorado Environmental Research Laboratories, For Sale By The Support of Documents., U.S. Government Printing Building Office, 1975.
- Hecla. H.M.S. *Oceanographic Observations In The Eastern Caribbean Sea*. 1971.
- Heezen, B.C., M. Tharp. *Physiographic Diagram of The South Atlantic Ocean, The Caribbean Sea, The Scotia Sea, and The Eastern Margin of The South Pacific Ocean*. New York, Geological Society of America, 1961.
- Heinbokel, J.F. *Studies on the Functional Role of Tintinnids in the Southern California Bight. I. Grazing and Growth Rates in Laboratory Cultures*. Marine Biology V47, pp. 177-189, 1978.
- Heneman, B. *Persistent Marine Debris In The North Sea, Northwest Atlantic Ocean, Wider Caribbean Area, and The West Coast Of Baja California*. Henneman, Burr Bolinas, pp. 160, California, 1988.
- Henwood, T., W. Stuntz. *Evaluation of U.S. Turtle Protective Measures Under Existing Regulations, Including Estimates of Shrimp Trawler Related Turtle Mortality In The Wider Caribbean*. U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 1992.
- Herman R., H. Dahms. *Meiofauna Communities Along A Depth Transect Off Halley Bay (Weddell Sea-Antarctica)*. Polar Biology, V12 N2, pp. 313-320, 1992.
- Hey E. *A Healthy North Sea Ecosystem And A Healthy North Sea Fishery - 2 Sides Of The Same Regulation*. Ocean Development and International Law, V23 N2-3, pp. 217-238, 1992.
- Hey, E., G. Peet, eds. *The Status of The North Sea Environment. Reasons For Concern: Proceedings of the 2nd North Sea Seminar '86*, Rotterdam, V2. pp. 257-284, 1986.
- Hickel, W. *Nitrogen and Phytoplankton Stocks in the German Bight in Relation to the Eutrophication Problem*. Hamburg: Biologist Anstalt Helgoland, 1989.

- Hickel, W., E. Bauerfeind, U. Niermann, H.V. Westernhagen. *Oxygen Deficiency In The South-Eastern North Sea: Sources And Biological Effects*. Hamburg: Biologische Anstalt Helgoland, 1989.
- Highfield, R. *Survey Into North Sea Waste Could Save One Billion Pounds A Year*. Daily Telegraph, 7 November, p.9, SHIP: "challenger"/rv. 1989.
- Hill, H.W. *Currents and Water Masses*. North Sea Science, NATO North Sea Science Conference, MIT Press, Cambridge, Massachusetts, pp. 17-42, 1973.
- Hill, R.T., I.T. Knight, M.S. Anikis, R.R. Colwell. *Benthic Distribution of Sewage Sludge Indicated by Clostridium Perfringens at a Deep-Ocean Dump Site*. Applied and Environmental Microbiology, V59, N1, pp. 47-51, 1993.
- Hiroshi, I. ed. *The Mediterranean Resorts: Method of Environmental Preservation and Development*. Chichukai No Rizoto: Kankyohozon To Kaihatsu, Tokyo, Japan: Process Architecture Publishing Company, 1990.
- Hjulstrom, F. *Transportation of Detritus by Moving Water*. Recent Marine Sediments in Trask, pp. 5-31, 1939.
- Hodges, L. *Environmental Pollution*. Second Edition, Holt, Rinehart and Winston, New York, New York, 1973.
- Hoekstra, M. *Macrowake Features of a Range of Ships*. Report N410461-1-PV, Maritime Research Institute, Netherlands, 1991.
- Holm-Hansen, O., S.Z. El-Sayed. *Primary Production and the Factors Controlling Phytoplankton Growth in the Southern Ocean*. Applications Within Antarctic Biology. Smithsonian Institution, pp. 11-50, Washington, D.C., 1977.
- Horne, R.A.. *Marine Chemistry*. John Wiley and Sons, New York, 1969.
- Hosford, W.F., J.L. Duncan. *The Aluminum Beverage Can*. Scientific American, 271(3), 49, 1994.
- Hovland, F. *Shipboard-generated Waste - Treatment and Disposal Systems on Cruise Vessels*. Waste and Gallery Systems, 1992.
- Hubold G., W. Ekau. *Feeding Patterns of Post-Larval and Juvenile Notothenioids In The Southern Weddell Sea (Antarctica)*. Polar Biology, V10 N4, pp. 255-260, 1990.
- Hunter, J.R., H. Dorr. *Thresholds for Filter Feeding in Northern Anchovy, Engraulis mordax*. CalCOFI Report, VXXIII, 1982.
- Hupkes R. *Pollution Of The North Sea By West European Rivers*. Water Science and Technology, V24 N10: pp. 69-75, 1991.
- Hydes, D. J., H. Edmunds. *NERC North Sea Community Research Programme Qualitative Assessment Of Nutrient Measurements September 1988 To August 1989: Preliminary*

*Report For The Department of The Environment. School of Environmental Sciences, University of East Anglia, Norwich, 1989.*

*Hydrographic Data For Caribbean Sea, and For Pesca Serial Station At 1738'N 67 00'W For 1971. Compiled By G.E. Shanley, Series Title: University of Puerto Rico (Mayaguez Campus), Department of Marine Sciences Data Reference Report , 72-1, 1971.*

Hyman, M. *Modeling Ship Microbubble Wakes.* Dahlgren Division Naval Suffice Warfare Center, Panama City, Florida, 1994.

Hyman, M., J. Kamman, R. Smith, R.C. Nguyen. *Bubble Transport in Ship Wakes With Application to Wake Modification.* Naval Coastal System Center Technical Memorandum, June, 1987.

Hyman, M.C. *Influence of Temperature Stratification on the Development of Surface Ship Micro-Bubble Wakes.* NCSC TN 1017-90, 1989.

Hyman, M.C. *Initial Data Plane Estimation for Surface Ships-II.* NCSC Technical Note, NCSC TN 930-88, 1988.

Hyman, M.C. *Numerical Simulation of the Hydrodynamic Wake of a Surface Ship.* Naval Coastal System Center Technical Note, pp. 544-690, 1990.

Hyman, M.C. *Simulation of the Interaction Between A Wind-Driven Sea State and Surface Ship Wakes.* Naval Coastal System Center Technical Memorandum, CSS TM pp. 584-591. 1992.

Ibrekk H., J. Molvaer, B. Faafeng. *Nutrient Loading To Norwegian Coastal Waters and Its Contribution To The Pollution Of The North Sea.* Water Science and Technology, V24 N10, pp. 239-249, 1991.

*ICES Coordinated Monitoring Programme For Contaminants In Fish And Shellfish, 1978 and 1979, and Six-Year Review of ICES Coordinated Monitoring Programme.* Copenhagen : International Council for the Exploration of the Sea, Series title: Cooperative research report, N126, 1984.

*ICES Coordinated Monitoring Programme, 1977.* Copenhagen, Denmark : International Council for the Exploration of the Sea, Series title: Cooperative research report, N98, 1980.

Ichiye, T. *Experiments On Circulation of The Gulf of Mexico and The Caribbean Sea With A Rotating Tank.,* 1969.

*IDOE Workshop On The Geology And Marine Geophysics of The Caribbean Region And Its Resources (1975: Kingston, Jamaica) Geology, Geophysics and Resources of The Caribbean : Report of The IDOE Workshop on The Geology and Marine Geophysics Of The Caribbean Region and Its Resources, Kingston, Jamaica, 17-22 February, Oceanographic Commission, UNESCO, 1977.*

- Ijlstra T. *North Sea Pollution -Vessel-Source Pollution, Environmental Management and The Establishment of Eezs*. Marine Policy, V17 N2, pp. 130-135, 1993.
- Ijlstra, T. *Laws Of The Sea. Enforcement Of MARPOL: Deficient Or Impossible?* Marine Pollution Bulletin V20(12), pp. 596-597, 1989.
- Ijlstra, T., F. Jong F. *Current Legal Developments. North Sea*. International J. Estuary Coast Law, V3, N3, pp. 246-265, 1988.
- IMO Adopts North Sea Annex V Special Area Status*. North Sea Monitor, 89.4, p.10, 1989.
- IMO Caribbean Project Gets Under Way*. Marine Pollution Bulletin, V28, N11, 1994.
- IMO Special Areas (Antarctic)*. Department of Transportation, U.S. Coast Guard 33 CFR Parts 151 and 155, Final Rule, 1994.
- IMO/FAO/UNESCO/WMO/WHO/IAEA/UN/UNEP. *Review Of Potentially Harmful Substances*. Joint Group of Experts on The Scientific Aspects of Marine Pollution [GESAMP]. Nutrients. GESAMP Reports and Studies N34, pp. 40, 1990.
- Implementation of the Ocean Dumping Ban Act of 1988*. Hearing Before the Subcommittee on Oceanography, Great Lakes and the Outer Continental Shelf and the Subcommittee on Fisheries and Wildlife Conservation and the Environment of the Committee on Merchant Marine and Fisheries House of Representatives, One Hundred Second Congress, First Session on Implementation of the 1988 Ocean Dumping Ban Act and Legislative Reauthorization, Current Permit and Penalty Structures, Trust Accounts and New Treatment Technologies, Research to Assess Effects of Sludge on Marine Life, and Beneficial Uses of Sludge, 1991.
- Incineration Ships Poison North Sea's Sediments*. New Scientist, 119 [1632], p.33, 1988.
- Ingri J., R. Lofvendahl, K. Bostrom. *Chemistry of Suspended Particles In The Southern Baltic Sea*. Marine Chemistry, V32 N1 pp. 73-87, 1991.
- International Atomic Energy Agency. *An Oceanographic Model for the Dispersion of Wastes Disposed of in the Deep Sea*. Joint Group of Experts on the Scientific Aspects of Marine Pollution GESAMP Originally Approved at its Thirteenth Session Held in Geneva, 1983.
- International Convention for the Prevention of Pollution from Ships, 1973*. MARPOL 73.
- International Council For The Exploration of The Sea. Advisory Committee On Marine Pollution*. ICES, Cooperative Research Report, N190, 1992.
- International Council for the Exploration of the Sea. *Working Group for the International Study of the Pollution of the North Sea and its Effects on Living Resources and their Exploitation*. Report of Working Group for the International Study of the Pollution of the North Sea and its Effects on Living Resources and their Exploitation. Charlottenlund, Denmark: International Council for the Exploration of the Sea, 1974.
- International Maritime Organization. *Articles, Protocols, Annexes, Unified Interpretations of the International Convention for the Prevention of Pollution from Ships, 1973, as Modified*

by the Protocol of 1978 Relating Thereto. MARPOL 73/78 Consolidated Edition 1991, IMO London, 1992.

International Maritime Organization. *Guidelines for the Implementation of Annex V of MARPOL 73/78*. International Maritime Organization, London, 1988.

*International Symposium on Biological and Hydrographical Problems of Water Pollution in the North Sea and Adjacent Waters (1967: Helgoland, West Germany)* Internationales Symposium 1967: Vorträge und Diskussionen = International Symposium 1967 : Hamburg: Biologische Anstalt Helgoland, 1968.

*Introduction of Sea Garbage Disposal Regulation*. Marine Pollution Bulletin, V20, N2, 1989.

*Inventory of Caribbean Marine And Coastal Protected Areas*. Washington, D.C.: Department of Regional Development, Executive Secretariat of Economic and Social Affairs, General Secretariat, Organization Of American States: Office Of International Affairs, National Park Service, U.S. Department of the Interior, 1988.

*IOC/FAO/UNEP International Workshop On Marine Pollution In The Caribbean And Adjacent Regions (1976 : Port Of Spain, Trinidad And Tobago)* Report of The IOC/FAO/UNEP International Workshop on Marine Pollution In The Caribbean and Adjacent Regions, Port-of-Spain, Trinidad, 13-17 December 1976, Paris, France: Unesco, 1977.

Jansson, B.O. *Ecosystem Approach to the Baltic problem*. Bulletins From the Ecological Research Committee, Stockholm, Ekologikommitten, Statens, 1972.

Jany,I., K. Scanlon, A. Mauffret. *Geological Interpretation Of Combined Seabeam, Gloria and Seismic Data From Anegada Passage (Virgin-Islands, North Caribbean)*. Marine Geophysical Researches, V12 N3, pp. 173-196, 1990.

*Jefferson 1992-1993 Sewage Discharge Monitoring Report*. Jefferson Department, New Orleans, LA, 1992-1993.

Jeftic, L., M. Bernhard, A. Demetropoulous, F. Fernex, G. Gabrielides, F. Gasparovic, Y. Halim, D. Orhon, L.J. Saliba. *State of The Marine Environment In The Mediterranean Region*. UNEP Regional Seas Report Study, N132, pp. 172, 1990.

Jenkins, B.S. *Density Currents and Turbidity Currents in Waste Disposal in the Ocean*. The University of New South Wales Water Research Laboratory, Manly Vale, N.S.W., Australia, 1970.

Jenkins, S.A., D.L. Inman. *On a Submerged Sphere in a Viscous Fluid Excited by Small-Amplitude Motions*. Journal Fluid Mechanics, V157, pp. 199-224. 1985.

Jenkins, S.A., J. Wasyl. *Resuspension of Estuarial Sediments by Tethered Wings*. Journal of Coastal Research, V6, N4, pp. 961-980, 1990.

Jenkins, S.A., J. Wasyl. *Time Stepped Suspended Transport Model for the Dispersion of Optical Particles in Coastal Waters*. Submitted to Office of Naval Research, Code 1153, pp. 84, 1994.

- Jenkins, S.A., J.A. Nichols, D.W. Skelly. *Coupled Physical-Biological Dispersion Model for the Fate of Suspended Solids in Sewage Discharged into the Ocean*. SIO Reference Series 89-3, pp. 53, 1989.
- Jerlov, N.G. *Factors Influencing The Transparency of The Baltic Waters*. Goteborg, Elanders Boktryckeri Aktiebolag, 1955.
- Joannessen, O.M. *Preliminary Results of Some Oceanographical Observations Carried Out Between Barbados and Tobago*. Marine Sciences Manuscript Report N8, McGill University, November, 1968.
- Johnston, C.S., R.J. Morris, eds. *Oily Water Discharges: Regulatory, Technical And Scientific Considerations*. London: Applied Science Publishers, 1980.
- Johnston, R., P.G.W. Jones. *Inorganic Nutrients In The North Sea*. New York: American Geographical Society, 1965.
- Johnston, R.K., D. Lapota. *A New Approach for Evaluating Biological Toxicity at Aquatic Hazardous Waste Sites*. Reprinted from Proceedings of Sixth Symposium on Coastal and Ocean Management/ASCE, Charleston, pp. 422-436, 1989.
- Joiris C. *Spring Distribution and Ecological Role Of Seabirds and Marine Mammals In The Weddell Sea, Antarctica*. Polar Biology, V11 N7, pp. 415-424, 1991.
- Jones R., J. Amador. *Methane and Carbon Monoxide Production, Oxidation, and Turnover Times in The Caribbean Sea as Influenced by The Orinoco River*. Journal of Geophysical Research-Oceans, V98 N2, pp. 2353-2359, 1993.
- Jones, Dr. N.V. *The North Sea: Perspectives On Regional Environmental Cooperation-The North Sea Environment: Features and Problems*. London , Boston: Graham & Trotman Martinus Nijhoff, 1990.
- Jones, W.P., B.E. Launder. *The Prediction of Laminarization with a two Equation Model of Turbulence*. International J. Heat Mass Transfer, 15, pp. 301, 1972.
- Jonsson P., C. Rappe, L. Kjeller, A. Kierkegaard. *Pulp-Mill Related Polychlorinated Organic Compounds In Baltic Sea Sediments*. Ambio, V22 N1, pp. 37-43, February, 1993.
- Jonsson, I.G., N.A. Carlsen. *Experimental and Theoretical Investigations in an Oscillatory Turbulent Boundary Layer*. J. Hydraulic Research, V14, N1, pp. 45-60, 1976.
- Joy, S. *Effect Of Shipping And Navigation On The Marine Environment*. North Sea Forum. London: Council for Environmental Conservation, pp. 147-151, 1987.
- Kamath, S. *Biological Nitrification/Denitrification of High Sodium Nitrite (Navy Shipyard) Wastewater*. Environmental Pollution, V69, pp. 25-38, 1991.
- Kang, S.H., G.A. Fryxell. *Phytoplankton in the Weddell Sea, Antarctica: Composition, Abundance, and Distribution in the Water-Column Assemblages of the Marginal Ice-Edge Zone During Autral Autumn*. Marine Biology, V116, pp. 335-348, 1993.

- Kasoulides, G.C. *North Sea Declared A 'Special Area'*. Marine Pollution Bulletin, V19 [11], pp. 551-552, 1988.
- Kastelein, R.A., M.S.S. Lavaleije. *Foreign Bodies In The Stomach Of A Female Harbour Porpoise (Phocoena phocoena) From The North Sea*. Aquatic Mammals, V18, N2, pp. 40-46, 1992.
- Katz, C.N., D.B. Chadwick, G.S. Douglas. *Real-Time Fluorescence for Measurements Intercalibrated with GC-MS*. Ocean Proceedings 1991 Conference, pp. 351-358, 1991.
- Katz, C.N., D.B. Chadwick. *Synoptic Mapping and Modeling of a Sewage Outfall Plume*. The Marine Technology Society Ocean Conference Proceedings, pp. 207-215, California, 1993.
- Kautsky H. *The Impact Of Pulp-Mill Effluents On Phyto-benthic Communities In The Baltic Sea*. Ambio, V21 N4m, pp. 308-313, 1992.
- Kavanaugh, M.C., J.C. Leckie, eds. *Particulates in Water, Characterization, Fate, Effects, and Removal*. Advances in Chemistry Series, 1980.
- Kemp, J.F., A.F.M. De Bievre. *A Regional Vessel Traffic Service For The North Sea. The North Sea: Perspectives on Regional Environmental Cooperation*. Eds. D. Freestone, T. Ijlstra, V5, N1-3 pp. 167-179, 1990.
- Kempe, S., G. Liebezeit, V. Dethlefsen, U. eds. *Biogeochemistry and Distribution of Suspended Matter In The North Sea and Implications To Fisheries Biology*. Scientific Common Problems of The Environment United Nations Environment Programme, Hamburg (FRG) pp. 552, 1988.
- Kempe, S., T.C. Jennerjahn. *The Vertical Particle Flux In The Northern North Sea, Its Seasonality And Composition*. Scientific Common Problems of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry and Distribution of Suspended Matter In The North Sea and Implications-To-Fisheries-Biology, pp. 65 229-268, 1988.
- Kennedy, J.F., R.A. Froebel. *Two-Dimensional Turbulent Wakes in Density-Stratified Liquids*. A.S.M.E. Publication, 64-WA/UNT-11, 1964.
- Kennett, J. *Marine Geology*. Prentice-Hall, Inc., Englewood Cliffs, New Jersey, 1982.
- Kersten, M., Klatt, V. *Trace Metal Inventory And Geochemistry Of The North Sea Shelf Sediments*. Scientific, Common Problems Of The Environment United, Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry and Distribution of Suspended Matter In The North Sea And Implications-To-Fisheries-Biology, N65, pp. 289-311, 1988.
- Kester, D.R., W.V. Burt, J.M. Capuzzo, P.K. Park, B.H. Ketchum, I.W. Duedall. *Wastes in the Ocean Volume 5*. John Wiley & Sons, New York, 1985.

- Ketchum, B.H., D.R. Kester, P.K. Park. *Ocean Dumping of Industrial Wastes*. Plenum Press, New York, 1979.
- Keys, J.R. *Antarctic Marine Environments and Offshore Oil*. Commission for the Environment Wellington, New Zealand, 1984.
- Kings Lynn 1992-1994 Sewage Discharge Monitoring Report. National Rivers Authority, Kings Lynn Treatment Works, United Kingdom, 1992-1994.
- Kivi K., H. Kuosa. *Late Winter Microbial Communities In The Western Weddell Sea (Antarctica)*. Polar Biology, V14 N6, pp. 389-399, 1994.
- Kjerfve, B. *Physical Flow Processes in Caribbean Waters Over a Range of Scales*, Belle W. Baruch Institute for Marine Biology and Coastal Research, University of South Carolina.
- Klages N. *Food and Feeding Ecology Of Emperor Penguins In The Eastern Weddell Sea*. Polar Biology, V9 N6, pp. 385-390, 1989.
- Klomp R. *Modeling The Transport And Fate Of Toxics In The Southern North Sea*. Science of The Total Environment, V97-8, pp. 103-114, 1990.
- Kohonen J.T. *Protection of The Marine Environment of The Baltic Sea - An Example On Multilateral Cooperation*. Marine Pollution Bulletin, V23, pp. 541-544, 1991.
- Koroleva, N.D., V.A. Kiselev. *Soviet Marine Pollution Legislation, Prevention of Pollution from Ships and the Law of the Sea Convention*. Soviet Association of Maritime Law.
- Koslow, J.A. *Feeding Selectivity of Schools Northern Anchovy, *Engraulis mordax*, in the Southern California Bight*. Fishery Bulletin, V79, pp. 131-229, 1981.
- Koubbi P. *Influence of The Frontal Zones on Ichthyoplankton and Mesopelagic Fish Assemblages In The Crozet Basin (Indian Sector Of The Southern Ocean)*. Polar Biology, V13 N8, pp. 557-564, 1993.
- Kress N., A. Golik, B. Galil, M. Krom. *Monitoring The Disposal of Coal Fly Ash at a Deep Water Site in The Eastern Mediterranean Sea*. Marine Pollution Bulletin, V26, N8, pp. 447-456, 1993.
- Kristiansen S., E. Syvertsen, T. Farbrot. *Nitrogen Uptake In The Weddell Sea During Late Winter and Spring*. Polar Biology, V12 N2, pp. 245-251, 1992.
- Krohn J., A. Muller, W. Puls. *Pollutant Transport Monitoring And Prediction By Mathematical Modeling-North Sea and Adjacent Estuaries*. Marine Pollution Bulletin, V23, pp.699-702, 1991.
- Kruize R. *Water Pollution Control In Amsterdam - Its Effect On North Sea Environment and The Role of Harbour Reception Facilities*. Water Science and Technology, V24 N10, pp. 155-160, 1991.

- Kruize, R.R. ed. *North Sea Pollution--Technical Strategies For Improvement: Proceedings of The LAWPRC International Conference, Held In Amsterdam*. The Netherlands, 10-14 September 1990, Oxford, New York: Pergamon Press, 1991.
- Kullenberg, G. *Entrainment Velocity in Natural Stratified Sheer Flow*. *Estuary Coastal Marine Science*, 5, pp. 329-338, 1977.
- Kullenberg, G. *Overall Report On The Baltic Open Sea Experiment 1977 (BOSEX)*. Copenhagen, Denmark: International Council for the Exploration of the Sea, 1984.
- Kullenberg, G. *The State Of The Baltic*. Oxford, Pergamon Press, New York, 1981.
- Kuparinen J., P. Bjornsen. *Bottom-Up and Top-Down Controls Of The Microbial Food Web In The Southern Ocean - Experiments With Manipulated Microcosms*: *Polar Biology*, V12 N2, pp. 189-195, 1992.
- La Violette, P.E., ed. *Seasonal and Interannual Variability of the Western Mediterranean Sea*. Coastal and Estuarine Studies Series, V46, AGU, 1994.
- Laane R.W.P.M., J. Vandermeer, A. Devries, A. Vandergiessen. *Monitoring The Progress of Attempts To Reduce Nutrient Load And Inputs of Certain Compounds In The North Sea By 50-Percent*. *Environmental Management*, March-April, V14 N2, pp. 221-227, 1990.
- Laevastu, T. *Surface Water Types of the North Sea and its Characteristics*. *Serial Atlas of the Marine Environment Folio*, 4, American Geographic Society, 1963.
- Laevastu, T. *Synopsis of Information on Oceanography of the North Sea*. FAO, Fisheries Division, Biology Branch, Report FB/60/SZ, 1960.
- Lajczak A., M. Jansson. *Seasonal Variations In Suspended Sediment Yield In The Baltic Sea Drainage Basin*. *Nordic Hydrology*, V24 N1 pp. 53-64, 1993.
- Lancelot C., G. Billen, C. Veth, S. Becquevort S, S.Mathot. *Modelling Carbon Cycling Through Phytoplankton and Microbes In The Scotia-Weddell Sea Area During Sea Ice Retreat*. *Marine Chemistry*, V35 N1-4, pp. 305-324, 1991.
- Lancelot C., S. Mathot, C. Veth, H. Debaar. *Factors Controlling Phytoplankton Ice-Edge Blooms In The Marginal Ice-Zone Of The Northwestern Weddell Sea During Sea Ice Retreat 1988 - Field Observations and Mathematical Modelling*. *Polar Biology*, V13 N6, pp. 377-387, 1993.
- Landmann, G. *An In Situ Camera System, Developed And Tested For The Characterization Of Suspended Matter In The Ocean*. Scientific Common Problems of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry and Distribution of Suspended Matter In The North Sea and Implications-To-Fisheries-Biology. N65 pp. 207-228. 1988.
- Lapota, D., D.E. Rosenberger, D. Duckworth. *A Bioluminescent Dinoflagellate Assay for Detecting Toxicity in Coastal Waters*. *Bioluminescence and Chemiluminescence*, John Wiley & Sons, Chichester, 1994.

- Lapota, D., G.J. Moskowicz, D.E. Rosenberger, J.G. Grovhoug. *The Use of Stimulable Bioluminescence from Marine Dinoflagellates as a Means of Detecting Toxicity*. The Marine Environment, ASTM Publication 1216, pp. 18, 1994.
- Larue D., Warner A. *Sedimentary Basins of The NE Caribbean Plate Boundary Zone and Their Petroleum Potential*. Journal of Petroleum Geology, V14 N3, pp. 275-290, 1991.
- Latin American and Caribbean Commission on Development and Environment. Our Own Agenda*. Latin American and Caribbean Commission on Development and Environment. Washington, D.C.: Inter-American Development Bank, United Nations Development Programme, New York, N.Y., 1990.
- Laubscher R., R. Perissinotto, C. Mcquaid. *Phytoplankton Production and Biomass At Frontal Zones In The Atlantic Sector Of The Southern Ocean*. Polar Biology, V13 N7, pp. 471-481, 1993.
- Lavoie, D.M. *Chemical, Biological, And Physical Measurements From The Mediterranean Sea, Summer 1980*. 1982.
- Law Of The Sea With Emphasis on The Mediterranean Issues*. Thessaloniki: Institute of International Public Law and International Relations of Thessaloniki, 1991.
- Lawrence J., N. Craig. *Strategy For Protecting The North Sea-A Chemical Industry Perspective*. Water Science and Technology, V24 N10, pp.269-276, 1991.
- Laws of the Sea, The Development of International Regimes of the Protection of Marine Areas*. Marine Pollution Bulletin, V22, N1, pp. 6-8, 1991.
- Laws of the Sea, the North Sea Conference: Bremen 1984-London 1987*. Marine Pollution Bulletin, V17, N9, pp. 394-399, 1986.
- Lazaro, B.J., J.C. Lasheras. *Particle Dispersion in a Developing Free Shear Layer, Part I Forced Flow*. J. Fluid Mechanics, V235, pp. 179-221, 1992.
- Lazaro, B.J., J.C. Lasheras. *Particle Dispersion in a Developing Free Shear Layer, Part 2 Forced Flow*. J. Fluid Mechanics, V235, pp. 179-221, 1992.
- Lebedev, V. *Antarctica*. Foreign Languages Publishing House, Moscow, Russia, 1959.
- Lee, A.J. *The North Sea, Setting the Scene*. Eds. P.J. Newman, A.R. Agg, Environmental Protection of the North Sea, Professional Publishing, 1988.
- Lee, A.J. *The Currents and Water Masses of the North Sea*. Oceanography, Marine Biology, 8, pp. 33-71, 1970.
- Lee, A.J., J. Ramster. *The Hydrography of the North Sea. A Review of Our Knowledge in Relation of Pollution Problems*. Helgol. Wiss. Meeresunters, 17, pp. 44-63, 1968.
- Leithe-Eirksen, R. *The Seas of Europe, The Baltic*. Greenpeace.
- Leithe-Eirksen, R. *The Baltic*. Collins & Brown, London, 1992.

- Leong, R.J., O'Connell. *A Laboratory Study of Particulate and Filter Feeding of the Northern Anchovy, Engraulis mordax*. J. Fish. Res. Bd. Canada V26, pp. 557-582, 1969.
- Lewis, R. E., A.M. Riddle. *Sea Disposal: Modelling Studies Of Waste Field Dilution*. Marine Pollution Bulletin, V20, N3, pp. 124-129, 1989.
- Leynaert A., P. Treguer, B. Queguiner, J. Morvan. *The Distribution of Biogenic Silica and the Composition of Particulate Organic Matter in the Weddell-Scotia Sea During Spring 1988*. Marine Chemistry, V35 N1-4, pp. 435-447, 1991.
- Lieberman, S.H., S. Clavell, D.B. Chadwick. *Techniques for Real-Time Environmental Monitoring*. Proceedings of the 16th Meeting U.S.-Japan Marine Facilities Panel: United States Cooperative Program in Natural Resources on Marine Facilities, pp. 495-499, 1989.
- Liebezeit, G. *Distribution Of Dissolved Humic Compounds In The Southern North Sea*. Scientific Common Problems of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry and Distribution of Suspended Matter In The North Sea And Implications-To-Fisheries-Biology, N65 pp. 153-162, 1988.
- Lipton, J., J.W. Gillett. *Uncertainty in Ocean-Dumping Health Risks: Influence of Bioconcentration, Commercial Fish Landings and Seafood Consumption*. Environmental Toxicology and Chemistry, V10, pp. 967-976, 1991.
- Liss, P.S. *Chemistry of the Sea Surface Microlayer*. Eds. J.P. Riley, G. Skirrow, Chemical Oceanography, V2, 2<sup>nd</sup> Edition, Academic Press, New York, Pp. 193-243, 1975.
- Liss, P.S. *Land To Ocean Transport of Contaminants: Comparison of River and Atmospheric Fluxes*. UNEP Regional Seas Reports and Studies, N114/2, 417-446, 1990.
- List, E.J., G. Gartrell, C.D. Winant. *Diffusion and Dispersion in Coastal Waters*. Journal of Hydraulic Engineering, V116, N10, pp. 1158-1179, 1990.
- Lithner, G. *Ronnskarsundersokningen*. Statens Naturvardsverk PM, 497, pp. 1-174, 1974.
- Littler M., P. Taylor, D. Littler. *Complex Interactions In The Control of Coral Donation on A Caribbean Reef Flat*. Oecologia, V80 N3, pp. 331-340, 1989.
- Liverpool 1990-1995 Sewage Discharge Monitoring Report*. National Rivers Authority, Liverpool Primary Tank Effluent to Syphon River, United Kingdom, 1990-1995.
- Lizotte M., C. Sullivan. *Biochemical Composition and Photosynthate Distribution In Sea Ice Microalgae Of McMurdo-Sound, Antarctica - Evidence For Nutrient Stress During The Spring Bloom*. Antarctic Science, V4 N1, pp. 23-30, 1992.
- Lodge D., W. McDowell. *Summary Of Ecosystem-Level Effects Of Caribbean Hurricanes*. Biotropica, V23 N4, pp. 373-378, 1991.
- London Dumping Convention, the First Decade and Beyond*. Provisions of the Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter, 1972

- Leong, R.J., O'Connell. *A Laboratory Study of Particulate and Filter Feeding of the Northern Anchovy, *Engraulis mordax**. J. Fish. Res. Bd. Canada V26, pp. 557-582, 1969.
- Lewis, R. E., A.M. Riddle. *Sea Disposal: Modelling Studies Of Waste Field Dilution*. Marine Pollution Bulletin, V20, N3, pp. 124-129, 1989.
- Leynaert A., P. Treguer, B. Queguiner, J. Morvan. *The Distribution of Biogenic Silica and the Composition of Particulate Organic Matter in the Weddell-Scotia Sea During Spring 1988*. Marine Chemistry, V35 N1-4, pp. 435-447, 1991.
- Lieberman, S.H., S. Clavell, D.B. Chadwick. *Techniques for Real-Time Environmental Monitoring*. Proceedings of the 16th Meeting U.S.-Japan Marine Facilities Panel: United States Cooperative Program in Natural Resources on Marine Facilities, pp. 495-499, 1989.
- Liebezeit, G. *Distribution Of Dissolved Humic Compounds In The Southern North Sea*. Scientific Common Problems of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry and Distribution of Suspended Matter In The North Sea And Implications-To-Fisheries-Biology, N65 pp. 153-162, 1988.
- Lipton, J., J.W. Gillett. *Uncertainty in Ocean-Dumping Health Risks: Influence of Bioconcentration, Commercial Fish Landings and Seafood Consumption*. Environmental Toxicology and Chemistry, V10, pp. 967-976, 1991.
- Liss, P.S. *Chemistry of the Sea Surface Microlayer*. Eds. J.P. Riley, G. Skirrow, Chemical Oceanography, V2, 2<sup>nd</sup> Edition, Academic Press, New York, Pp. 193-243, 1975.
- Liss, P.S. *Land To Ocean Transport of Contaminants: Comparison of River and Atmospheric Fluxes*. UNEP Regional Seas Reports and Studies, N114/2, 417-446, 1990.
- List, E.J., G. Gartrell, C.D. Winant. *Diffusion and Dispersion in Coastal Waters*. Journal of Hydraulic Engineering, V116, N10, pp. 1158-1179, 1990.
- Lithner, G. *Ronnskarsundersokningen*. Statens Naturvardsverk PM, 497, pp. 1-174, 1974.
- Littler M., P. Taylor, D. Littler. *Complex Interactions In The Control of Coral Donation on A Caribbean Reef Flat*. Oecologia, V80 N3, pp. 331-340, 1989.
- Liverpool 1990-1995 Sewage Discharge Monitoring Report*. National Rivers Authority, Liverpool Primary Tank Effluent to Syphon River, United Kingdom, 1990-1995.
- Lizotte M., C. Sullivan. *Biochemical Composition and Photosynthate Distribution In Sea Ice Microalgae Of McMurdo-Sound, Antarctica - Evidence For Nutrient Stress During The Spring Bloom*. Antarctic Science, V4 N1, pp. 23-30, 1992.
- Lodge D., W. McDowell. *Summary Of Ecosystem-Level Effects Of Caribbean Hurricanes*. Biotropica, V23 N4, pp. 373-378, 1991.
- London Dumping Convention, the First Decade and Beyond*. Provisions of the Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter, 1972

(amended 1978, 1980, and 1989), and Decisions made by the Consultative Meeting of its Contracting Parties (1975-1989). Prepared by the Office of the London Dumping Convention, IMO, London, 1991.

- Long, E.R., D.D. McDonald, S.L. Smith, F.D. Calder. *Incidence of Adverse Biological Effects Within Ranges of Chemical Concentrations in Marine and Estuarine Sediments*. Environmental Management, V19, N1, pp. 81-97, Springer-Verlag New York, Inc., 1995.
- Lowestoft 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Lowestoft Treatment Works, United Kingdom, 1992-1994.
- Lundhansen L., M. Pejrup, J. Valeur, A. Jensen. *Gross Sedimentation Rates in The North Sea-Baltic Sea Transition - Effects Of Stratification, Wind Energy Transfer, and Resuspension*. Oceanologica Acta, V16 N3, pp. 205-212, 1993.
- Lyons M., P. Balls, W. Turrell. *A Preliminary Study of The Relative Importance of Riverine Nutrient Inputs To The Scottish North Sea Coastal Zone*. Marine Pollution Bulletin, V26 N11, pp. 620-628, 1993.
- MacIntyre, F. *Chemical Fractionation and Sea-Surface Microlayer Processes*. Ed. E.D. Goldberg, The Sea, V5, Wiley, New York, pp. 245-299, 1974.
- Manos A. *An International Programme For The Protection Of A Semi-Enclosed Sea. The Mediterranean Action Plan*. Marine Pollution Bulletin, V23, pp. 489-496, 1991.
- Manzi, D. *Shipboard-Generated Solid Waste Analysis-Naval Station, Norfolk*, 1994.
- Manzi, D. *Shipboard-Generated Solid Waste Analysis: Phase II Report*. Carderock Division, Naval Surface Warfare Center, Technical Report 63, CR93/01, January, 1994.
- Marine Environmental and Water Quality Models*. Copenhagen: International Council for the Exploration of the Sea, Series title: Cooperative Research Report , N156. 1980.
- Marine Pollution Bulletin, Environmental Perspectives for the Northern Seas*. Proceedings of Environment Northern Seas 1993, V29, N6-12, 1994.
- MARPOL 73/78 Consolidated Edition, 1991*. Articles, Protocols, Annexes, Unified Interpretations of the International Convention for the Prevention of Pollution from Ships, 1973, as modified by the Protocol of 1978 relating thereto, IMO, London, 1992.
- MARPOL Amended to Harmonize Survey and Certification Aspects*. Marine Environment Protection Committee-29th Session.
- Marshall, N. *Understanding The Eastern Caribbean and The Antilles: With Checklists Appended*/Nelson Marshall, St. Michaels, Md.: Th'Anchorage Publisher, 1992.
- Martin J., F. Elbazpoulichet, C. Guieu, M. Loyepilot. *River Versus Atmospheric Input of Material To The Mediterranean Sea - An Overview*. Marine Chemistry, V28 N1-3, pp. 159-182, 1989.

- Martin, J.E., E. Meiburg. *The Accumulation and Dispersion of Heavy Particles in Forced Two-Dimensional Mixing Layers, 1: The Fundamental and Subharmonic Cases*. Physical Fluids, A-6, pp. 1116-1132, 1994.
- Marvin, M., J. Burns, D.G. Capone. *Microbial Degradation of Sweetheart Cups in the Marine Environment*. A Report Submitted to the Sweetheart Cup Company, Inc., Chesapeake Biological Laboratory, University of Maryland, September, 1993.
- Mathot S., J. Dandois, C. Lancelot. *Gross and Net Primary Production In The Scotia-Weddell Sea Sector Of The Southern Ocean During Spring 1988*. Polar Biology, V12 N2, pp. 321-332, 1992.
- Matthews, J.E., T.L. Holcombe. *Regional Geological/Geophysical Study of The Caribbean Sea (Navy Ocean Area Na-9)*. Geophysical Maps of The Eastern Caribbean, 1976.
- Maul, G.A. *English-Spanish Bibliography In Physical Oceanography and Ocean Climate For The Caribbean Sea and Adjacent Regions*. Bibliografia Ingles- Espanol De Oceanografia Fisica Y Clima Oceanico Del Mar Caribe Y, Paris: Intergovernmental Oceanographic Commission, Unesco, 1988.
- Maul, G.A. *Implications Of Climatic Changes In The Wider Caribbean Region: Preliminary Conclusions Of The Task Team Of Experts*. Prepared By George Maul, Kingston, Jamaica: Caribbean Environment Programme, United Nations Environment Programme, CEP Technical Report, Mailliard TD171.5.C3 C46 N3, 1989.
- McConnell B., C. Chambers, M. Fedak. *Foraging Ecology of Southern Elephant Seals In Relation To The Bathymetry and Productivity of The Southern Ocean*. Antarctic Science, V4 N4, pp. 393-398, 1992.
- McCoy, F.W. *Floating Mega-litter In The Eastern Mediterranean*. Marine Pollution Bulletin, V19, N1, pp. 25-28, 1988.
- Mcdonald A. *Coastal Waters, Vol 1, Water Quality Management Plan North Sea - Framework For Analysis* - Koudstaal, R. Environment & Planning, V22 N11:1547-1549, Nov., 1990.
- McDonald S., M. Kennicutt, J. Sericano, T. Wade, H. Liu, S.H. Safe. *Correlation Between Bioassay-Derived P4501A1 Induction Activity and Chemical Analysis Of Clam (Laternula-elliptica)*. Extracts From McMurdo Sound, Antarctica. Chemosphere, V28 N12, pp. 2237-2248, 1994.
- McIntyre, A.D. *Pollution in the North Sea From Oil-Related Industry, an Overview*. Eds. P.J. Newman, A.R. Agg, Environmental Protection of the North Sea, Heinemann, Oxford, pp. 425, 1988.
- McIntyre, A.D. *The Benthos of the Western North Sea*. Rapp. R.-v Reun. Cons. Internationa Expolorer, Mer, 172, pp. 405-417, 1978.
- McIntyre, A.D., J.B. Pearce, eds. *Biological Effects Of Marine Pollution and The Problems of Monitoring: Proceedings From ICES Workshop Held In Beaufort, North Carolina 26 February-2 March 1979*. Copenhagen K, Denmark: Conseil International Pour

- L'exploration de la mer, Series title: Rapports et Proces-Verbaux des Reunions V179, 1980.
- McLeay, D. *Aquatic Toxicity of Pulp and Paper Mill Effluent: A Review*. Environment Canada, Fisheries and Oceans, D. McLeay and Associates Ltd., 1987.
- Meadows, G.A., D. Lyzenga, J. Lyden, R. Beck. *Nonintrusive, Multiple-Point Measurements of Water Surface Slope, Elevation and Velocity*. Proceedings of the Eighteenth ONR Symposium on Naval Hydrodynamics, National Academy Press, pp. 349-360, Washington, D.C., 1990.
- Meadows, L., G. Meadows, A. Troesch, S. Cohen, K.P. Beier, G. Rot. *Lagrangian Velocity Profiles in the Wake of a High Speed Vessel*. Ocean Engineering, V21, N2, pp. 221-242, 1994
- Mediterranean Action Plan and Convention For The Protection of The Mediterranean Sea Against Pollution and Its Related Protocols*. CA CORPORATE AUTHOR(S): UNEP, Nairobi (Kenya). Athens Greece UNEP pp. 61, 1992.
- Mediterranean Coastal Pollution: Proceedings of A Conference Held In Palma, Mallorca, 24-27 September, 1979*. New York: Pergamon Press, 1980.
- Mediterranean Pollution: Proceedings of A Conference Held in Palma, Mallorca, 24-27 September, 1979*. Oxford, New York: Pergamon Press, 1980.
- Mediterranean Sea*. New Encyclopedia Britannica 15th Edition, Encyclopedia Britannica Inc., 1974.
- Melvasalo, T. *Assessment of the Effects of Pollution on the Natural Resources of the Baltic Sea*. Eds. J. Pawlak, K. Grasshoff, L. Thorell, A. Tsiban, Baltic Sea Environment, Proc. 5B, pp. 322-342, 1980.
- Melvasalo, T. ed. *Assessment Of The Effects Of Pollution On The Natural Resources Of The Baltic Sea, 1980*. Helsinki: Baltic Marine Environment Protection Commission, Helsinki Commission, 1981.
- Mercier, J.R., R.I. Hires, M. Wu. *Model Study of the Dilution of Soluble Liquids Discharged From Tankers*. Report No. CD-D-12-74. Department of Transportation, U.S. Coast Guard, Washington, D.C., 1973.
- Merritt, G.E. *Wake Laboratory Experiment*. Cornell Aeronautical Laboratory Inc. New York, 1972.
- Metcalf, W.G. *Hydrographic Station Data Caribbean Sea Atoantis II Cruise 78 And Knorr Cruise 37*. Woods Hole Oceanographic Institution Reference, pp. 74-90, 1974.
- Metcalf, W.G. *Hydrographic Station Data From Atlantis II Cruise 56 To The Southeastern Approaches To The Caribbean Sea*. Woods Hole Oceanographic Institution Reference, 71-13, 1971.
- Metcalf, W.G., et al. *Hydrographic Station Data, Caribbean Sea, Knorr Cruise 25*. 1973.

- Miller R., J. Reinemund. *Energy And Mineral Potential Of The Central American-Caribbean Region -San-Jose, Costa-Rica*. Episodes, V13 N1:28-31, 1990.
- Miller, A.R. *Physical Oceanography of The Mediterranean Sea: A Discourse*. Woods Hole Oceanographic Institution Woods Hole, Massachusetts, U.S., 1964.
- Miller, R. *1913-History and Atlas Of The Fishes Of The Antarctic Ocean*. Carson City, Nevada: Foresta Institute For Ocean and Mountain Studies, 1993.
- Miloradov, M. ed. *Pollution Of The Mediterranean Sea: Proceedings Of An IAWPRC International Regional Conference Held In Split, Yugoslavia, 2-5 October 1985*. International Association on Water Pollution Research and Control Conference 1985: Split, Yugoslavia Pergamon Press, Oxford, New York, 1986.
- Miner, W.E., S.E. Ramber. *A Method of Approximating the Initial Data Plane for Surface Ship Wake Simulations*. Naval Research Laboratory Report 6376, November, 1988.
- Mitchell B., E. Brody, O. Holmhansen, C. McClain, J. Bishop. *Light Limitation Of Phytoplankton Biomass and Macronutrient Utilization In The Southern Ocean*. Limnology and Oceanography, V36 N8, pp. 1662-1677, 1991.
- Mitchell, Dr. R. *Conservation of Marine Benthic Biocenoses in The North Sea and The Baltic: A Framework For The Establishment of A European Network of Marine Protected Areas In The North Sea And The Baltic*. European Committee For The Conservation of Nature and Natural Resources, 1987.
- Moraitou-Apostolopoulou, M., V. Kiortsis. *Mediterranean Marine Ecosystems*. Plenum Press, New York, 1985.
- Moran S., R. Moore, S. Westerlund. *Dissolved Aluminum In The Weddell Sea*. Deep-Sea Research Part A-Oceanographic Research Papers, V39 N3-4a, pp. 537-547, 1992.
- Moran, M.A., R. Benner, R.E. Hodson. *Kinetics of Microbial Degradation of Vascular Plant Material In Two Wetland Ecosystems*. Oecologia V79, pp. 158-167, 1989.
- Moran, M.A., R.E. Hodson. *Dissolved Humic Substances of Vascular Plant Origin In A Coastal Marine Environment*. Limnology Oceanography, 1994.
- Moran, M.A., R.E. Hodson. *Contributions of Three Subsystems of A Freshwater Marsh To Total Bacterial Secondary Productivity*. Microbiology Ecology V24, pp. 161-170, 1992.
- Moran, M.A., R.E. Hodson. *Formation and Bacterial Utilization of Dissolved Organic Carbon Derived From Detrital Lignocellulose*. Limnology Oceanography V34, pp. 1034-1047, 1989.
- Moran, M.A., R.J. Wicks, R.E. Hodson. *Export of Dissolved Organic Matter From A Mangrove Swamp Ecosystem: Evidence From Natural Fluorescence, Dissolved Lignin Phenols, and Bacterial Secondary Production*. Marine Ecological Program Series V76, pp. 175-184, 1991.

- More Measures To Curb North Sea Pollution.* Meeting of International Maritime Organization's Environment Protection Committee [MEPC]. Marine Pollution Bulletin, V20 [12], pp. 591, 1989.
- Most Litter Is Plastic, Beach Study Shows.* Results of Pilot Study of Mediterranean Sea Beaches in 1988, IMO News, p.3, 1990.
- Muench R., H. Fernando, G. Stegen. *Temperature and Salinity Staircases In The Northwestern Weddell Sea.* Journal of Physical Oceanography, V20 N2, pp. 295-306, 1990.
- Muller-Karger F., C. McClain, T. Fisher, W. Esaias, R. Varela. *Pigment Distribution in The Caribbean Sea - Observations From Space.* Progress In Oceanography, V23 N1, pp. 23-64, 1989.
- Mullin, M.M. *Particle Size Analysis on Mixed Paper Pulp, ETA.* Limnology and Oceanography, Vol. 10, pp. 459-462, 1965.
- Naudascher, E. *Flow in the Wake of Self-Propelled Bodies and Related Sources of Turbulence.* J. Fluid Mech. 2, Part 4, 635, 1965.
- NAVSEA. *U.S. Navy Shipboard Solid and Plastics Waste Management Program Plan.* Naval Sea Systems Command, 1993.
- Nedwell D., T. Walker, J. Ellisevans, A. Clarke. *Measurements of Seasonal Rates and Annual Budgets of Organic Carbon Fluxes In An Antarctic Coastal Environment At Signy Island, South Orkney Islands, Suggest A Broad Balance Between Production and Decomposition.* Applied and Environmental Microbiology, V59 N12, pp. 3989-3995, 1993.
- Nehring D., W. Matthaus. *Current Trends in Hydrographic and Chemical Parameters and Eutrophication in The Baltic Sea.* Internationale Revue Der Gesamten Hydrobiologie, V76 N3, pp. 297-316, 1991.
- Nekritz, R. *Bottom Sediment Distribution In The Mediterranean Sea.* Marine Science Department U.S. Naval Oceanographic Office, Washington, D.C., 1963.
- Nelson, D.M., W.O. Jr. Smith. *Sverdrup Revisited - Critical Depths, Maximum Chlorophyll Levels, and The Control Of Southern Ocean Productivity By The Irradiance-Mixing Regime.* Limnology and Oceanography, V36 N8, pp. 1650-1661, 1991.
- Neumann, G. *Ober Die Komplexe Natr Des Seeganges, Teil 1 and 2.* Deut Hydrogr. Zeit., V5, N2/3, pp. 95-110, N5/6, pp. 252-277, 1952.
- Neumann, G., W.J. Jr. Pierson. *Principles of Physical Oceanography.* Prentice-Hall, Inc., Englewood Cliffs, N.J., 1966.
- Newman, P.J., A.R. Agg. *Environmental Protection of the North Sea.* Heinemann Professional Publishing, 1988.
- Nguyen, T.C., M. Hyman. *Simulation of Surface Ship Wakes Using a K-epsilon Turbulence Model.* NCSC Technical Note, NCSC TN 940-88, 1988.

Nguyen, T.D., M. Hyman. *Free Shear Flow Simulations with an Algebraic Stress Model*. NCSC Technical Note, NCSC TN 928-88, 1988.

Nicol S., M. Stolp. *Sinking Rates of Cast Exoskeletons of Antarctic Krill (*Euphausia-superba dana*) and Their Role In The Vertical Flux of Particulate Matter and Fluoride In The Southern Ocean*. Deep-Sea Research Part A-Oceanographic Research Papers, V36 N11, pp. 1753-1762, 1989.

Niels Daan, N. ed. *Data Base Report Of The Stomach Sampling Project 1981*. Copenhagen, Denmark: International Council for the Exploration of the Sea, 1989.

Nielsen, J.N. *Golfstrømmen*. Geografisk Tidsskrift, V28, N1, 1925.

Nielsen, P., I.A. Svendsen, C. Staub. *Onshore-Offshore Sediment Movement on a Beach*. Proceedings of 18th International Conference of Coastal Engineering, pp. 1475-1492, ASCE, New York, 1978.

Nielson, H., ed. *Environment and Pollution Measurement Sensors and Systems*. ECO3 Proceedings, The International Society for Optical Engineering, Washington, V1269, March, 1990.

Nielson, P. *Some Basic Concepts of Wave Sediment Transport*. Series Paper N20, Institute of Hydrodyn. and Hydr. Engineering, Technical University of Denmark, 1979.

Nihoul, J.C. *Hydrodynamic Models Of Shallow Continental Seas: Application To The North Sea*. Nihoul, Neupre, E. Riga, 1982.

Nollkaemper, A. *Laws of the Sea, Legal Protection of the Marine Environment From Pollution of International Watercourses: Recent Developments*. Marine Pollution Bulletin, V26, N6, pp. 298-301, 1993.

Nolting R., H. Debaar, A. Vanbennekomp, A. Masson. *Cadmium, Copper and Iron In The Scotia Sea, Weddell Sea and Weddell Scotia Confluence (Antarctica)*. Marine Chemistry, V35 N1-4, pp. 219-243, 1991.

*Nordic Action Plan On Pollution Of The Seas*. Copenhagen: Nordic Council of Ministers. pp. 105, 1990.

*North Sea becomes MARPOL 'Special Area'*. Marine Pollution Bulletin, 22 [5], pp. 218-219, 1991.

*North Sea to be Made Annex V "special area"*. Marine Environment Protection Committee 28th Session, World Maritime University, 1989.

*North Sea To Be Made Annex V 'Special Area'*. IMO News, [1] , pp. 12. 1990.

*North Sea*. Encyclopedia Americana, V20, Danbury Connecticut, 1990.

*North Sea*. Encyclopedia Britannica Macropedia, 15th Edition, Encyclopedia Britannica Inc., 1974.

- North Sea*. McGraw Hill Encyclopedia of Science and Technology, 7th Edition, V12, McGraw Hill Inc., 1992.
- Northern Europe's Seas, Northern Europe's Environment : Report To The Nordic Council's International Conference On The Pollution Of The Seas, 16-18 October 1989*. Stockholm, Sweden: The Council, 1989.
- Norton, R.L. *Assessment Of Pollution Loads To The North Sea*. National Technical Information Service, 1982.
- Norwich 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Norwich Treatment Works, United Kingdom, 1992-1994.
- Nothig E., B. Vonbodungen, Q. Sui. *Phytoplankton and Protozooplankton Biomass During Austral Summer In Surface Waters of The Weddell Sea and Vicinity*. Polar Biology, V11 N5, pp. 293-304, 1991.
- Nowlin W. *On Water Mass Exchange Between The Southern Ocean and The World Ocean - Emphasis On The Atlantic Sector*. Marine Chemistry, V35 N1-4, pp. 1-7, 1991.
- Nystuen J., C. Andrade. *Tracking Mesoscale Ocean Features In The Caribbean Sea Using Geosat Altimetry*. Journal Of Geophysical Research-Oceans, V98 N5 pp. 8389-8394, 1993.
- O'Connor, T.P., W.V. Burt, I.W. Duedall. *Oceanic Processes in Marine Pollution, Volume 2 Physicochemical Processes and Wastes in the Ocean*. Robert E. Krieger Publishing Company, Inc., Malabar, Florida, 1987.
- Ocean Dumping and Pollution*. Hearings Before the Subcommittee on Oceanography and the Subcommittee on Fisheries and Wildlife Conservation of the Environment of the Committee on Merchant Marine and Fisheries House of Representatives Ninety-Fifth Congress on Ocean Dumping (H.R. 4715, H.R. 5282, H.R. 5851), Ocean Pollution Research (S.1617), Nuclear Waste Disposal, Marine Sanctuaries, Serial No. 95-42, 1978.
- Ocean Refuse and its Impact on the Texas Shoreline*. Hearing Before the Subcommittee on Oversight and Investigations of the Committee on Merchant Marine and Fisheries House of Representatives, Ninety-Ninth Congress Second Session on The Problem of Coastal Pollution Caused by trash Discarded Offshore into the Marine Environment and the Options that Exist on the International, Federal, State, and Local Levels for Responding to it, 1986.
- Oceanographic Characteristics of the Mediterranean*.
- Odegaard H., B. Paulsrud, T. Bilstad, J. Pettersen. *Norwegian Strategies In The Treatment of Municipal Wastewater Towards The Reduction of Nutrient Discharges To The North Sea*. Water Science and Technology, V24 N10, pp. 179-186, 1991.
- Oehlenschlager J., H. Karl. *On The Burden of Fish and Other Marine Food From The Baltic Sea With Inorganic and Organic Pollutants*. Deutsche Lebensmittel-Rundschau, V88 N4, pp. 115-116, 1992.

- Ogden, J.C., E.H. Gladfelter. *Caribbean Coastal Marine Productivity: Results of A Planning Workshop at Discovery Bay Marine Laboratory, University of The West Indies Jamaica*, Paris: Unesco, 1986.
- Okubo, A. *Oceanic Diffusion Diagrams*. Deep-Sea Research, V18, pp. 789, 1971.
- Olson, P.H. *Handling of Waste in Ports*. Marine Pollution Bulletin, V29, N6-12, pp.284-295, Pergamon Press, New York, 1994.
- Olson, T.M., S.E. Gill, C.S. Alig. *Study of Solid and Plastic Waste Management Aboard USS Emory S. Land (AS 39)*. David Taylor Research Center, Bethesda, pp. 45, MD, 1989.
- OPHELLA, International Journal of Marine Biology*. Proceedings of the 8th Symposium of the Baltic Marine Biologists, Marine Biological Laboratory, Helsingor, Denmark, 1984.
- Oslo and Paris Commissions. *Progress Report On The Activities Of The Oslo And Paris Commissions, November 1987 March 1990*. London: Secretariat of the Oslo and Paris Commissions, pp. 51, 1990.
- Otto L., J. Zimmerman, G. Furnes, M. Mork, R. Saetre, G. Becker. *Review Of The Physical Oceanography of The North Sea*. Netherlands Journal of Sea Research, V26 N2-4, pp. 161-238, 1990.
- Owens N., J. Priddle, M. Whitehouse. *Variations In Phytoplanktonic Nitrogen Assimilation Around South Georgi and In The Bransfield Strait (Southern Ocean)*. Marine Chemistry, V35 N1-4, pp. 287-304, 1991.
- Palanques A., D. Drake. *Distribution and Dispersal of Suspended Particulate Matter on The Ebro Continental Shelf, Northwestern Mediterranean Sea*. Marine Geology, V95 N3, pp. 193-206, 1990.
- Pan, B.Y.K., T. Delorea, M. Crist, S. Preiss. *Determination of Navy and Marine Corps Wastewater: Rates and Sources Volume III*. Naval Facilities Engineering Command, Port Hueneme, California, December, 1990.
- Panagopoulos, P., D. Karavellas. *Applicability of Remote Sensing For Survey of Water Quality Parameters In The Mediterranean*. Mediterranean Action Plan, Athens (Greece) IOC, Paris (France), Map Technical Athens Greece UNEP, N67, pp. 140, 1992.
- Paolo F., F. Giuliano, eds. *Semi-Enclosed Seas: Exchange of Environmental Experiences Between Mediterranean And Caribbean Countries*. London, New York Elsevier Applied Science, 1992.
- Paper and Paper Production*. Greenpeace, Washington D.C., 1995.
- Park Y., L. Gamberoni, E. Charriaud. *Frontal Structure and Transport of The Antarctic Circumpolar Current In The South Indian Ocean Sector, 40-80-Degrees-E*. Marine Chemistry, V35 N1-4, pp. 45-62, 1991.
- Park, P.K., T.P. O'Connor, B.H. Ketchum, eds. *Historical and International Considerations*. Plenum, New York, New York, 1980.

- Parker, R.D. *Surface Drifter Data From The Caribbean Sea and Gulf Of Mexico*. 1979.
- Parr, A.E. *A Contribution To The Hydrography Of The Caribbean and Cayman Seas Based Upon The Observations Made By The Research Ship Atlantis, 1933-34*. Bingham Oceanography Collection, Bulletin, V5, 4, pp. 110, New Haven, 1937.
- Pearson, E.A., F.E. DeFraja. *Marine Pollution and Marine Waste Disposal, Proceedings of the 2nd International Congress, San Remo, 17-21st December 1973*. Pergamon Press Ltd Oxford, England, 1975.
- Peng, T.H., W.S. Broecker, G.G. Mathieu, Y.H. Li, A.E. Bainbridge. *Radon Evasion Rates in the Atlantic and Pacific Oceans as Determined During the GEOSECS Program*. J. Geophysics Research, 84, pp. 2471-2486, 1979.
- Perissinotto R., C. Rae. *Occurrence of Anticyclonic Eddies on The Prince-Edward Plateau (Southern Ocean) - Effects on Phytoplankton Biomass and Production*. Deep-Sea Research Part A-Oceanographic Research Papers, V37 N5, pp. 777-793, 1990.
- Perry, R.K. *Bathymetry of The Gulf of Mexico and The Caribbean Sea*. William & Heintz Map Corporation. Tulsa, Oklahoma: American Association of Petroleum Geologists, 1984.
- Peterson, M.H., T.J. Lennox Jr. *Effects of Exposure Conditions on The Corrosion of Mild Steel, Copper, and Zinc In Seawater*. Materials Performance, V23, pp. 15-18, 1984.
- Pimenov N., M. Ivanov. *Bacterial Chemosynthetic Processes in The Water Column of The Baltic Sea in The Winter*. Microbiology, V60 N6, pp. 778-783, 1991.
- Pineschi, L. *The Prevention of Marine Pollution From Ships According To Annex IV To The Protocol on Environmental Protection To The Antarctic Treaty*. Marine Pollution Bulletin, V24, N5, pp. 228-230, 1992.
- Pitter, P., J. Chudoba. *Biodegradability of Organic Substances in the Aquatic Environment*. CRC Press, Inc., Boca Raton, Florida, 1990.
- Pollution Of The Baltic Sea: A Two-Day Seminar On Legal Problems Concerned With Pollution, Uto, 14th - 15th September, 1985*. Stockholm: Stiftelsen Juristforlaget vid Stockholms Universitet, 1988.
- Pollution Prevention, The Navy Needs Better Plans for Reducing Ship Waste Discharges*. U.S. General Accounting Office, Letter Report for Congressional Requesters, National Security and International Affairs Division, November, 1994.
- Pomar M., V. Bruni, T. Maugeri. *Picoplankton Biomass In The Ross Sea (Antarctica)*. Polar Biology, V13 N1, pp. 1-6, 1993.
- Pomeroy, L.R., J.E. Sheldon, W.M. Sheldon. *Changes In Bacterial Numbers and Leucine Assimilation During Estimations of Microbial Respiratory Rates In Seawater By The Precision Winkler Method*. Applied and Environmental Microbiology, V60, pp. 328-332, 1994.

- Progress Reports On Cadmium, Mercury, Copper and Zinc.* Helsinki: Baltic Marine Environment Protection Commission, Helsinki Commission, 1987.
- Pruter, A.T. *Sources, Quantities and Distribution of Persistent Plastics in the Marine Environment.* Marine Pollution Bulletin, V18, N6B, pp. 305-310, 1987.
- Pujos M., O. Javelaud. *Depositional Facies of A Mud Shelf Between The Sinu River and The Darien Gulf (Caribbean Coast of Colombia)-Environmental Factors That Control Its Sedimentation and Origin of Deposits.* Continental Shelf Research, V11 N7, pp. 601-623. 1991.
- Pulp and Paper.* Greenpeace, Washington D.C., 1994.
- Pyewipe 1992-1994 Sewage Discharge Monitoring Report.* National Rivers Authority, Pyewipe Pumping Station, United Kingdom, 1992-1994.
- Radach G., M. Regener, F. Carlotti, W. Kuhn, A. Moll. *Modeling Water Column Processes In The North Sea.* Philosophical Transactions of The Royal Society of London Series A-Physical Sciences and Engineering, V343 N1669, pp. 509-517, 1993.
- Radach, G., J. Berg. *Trends in den Konzentrationen der Nährstoffe und des Phytoplanktons in der Helgolander Bucht (Helgoland Reede Daten).* Berichte der Biologischen Anstalt Helgoland N2., 1986.
- Rao, A.D., S.K. Dube, P.C. Sinha. *Wind Driven Seasonal Ocean Circulation Around Southern India Peninsula.* Indian Journal of Marine Sciences, V21, pp. 196-200, 1992.
- Rapson, W.H. *Paper.* Greenpeace, Washington D.C., 1995.
- Rau G., C. Sullivan, L. Gordon. *Delta-C-13 and Delta-N-15 Variations In Weddell Sea Particulate Organic Matter.* Marine Chemistry, V35 N1-4, pp. 355-369, 1991.
- Raudkivi, A.J. *Loose Boundary Hydraulics.* Third Edition, Pergamon Press, Inc., Maxwell House, Fairview Park, Elmsford, New York, 1990.
- Rawn, A.M. F.R. Bowerman, N.H. Brooks. *Diffusers for Disposal of Seage in Sea Water.* Diffusers for Disposal of Sewage in Sea Water. J. Sanit. Division, ASCE, 86, SA2, pp.65-105, 1960.
- Raymont, J.E.G. *Plankton and Productivity in the Oceans, Volume 2-Zooplankton.* Pergamon Press, Second Edition.
- Raymundo, C.C., M.R. Preston. *The Distribution of Linear Alkylbenzenes in Coastal and Estuarine Sediments of the Western North Sea.* Marine Pollution Bulletin, V24, N3, 1992.
- Reading A. *Caribbean Tropical Storm Activity Over The Past 4 Centuries.* International Journal of Climatology, V10 N4, pp. 365-376, 1990.

- Recent Advances In Selected Areas of Ocean Sciences In The Regions of The Caribbean, Indian Ocean, and The Western Pacific.* Presented at The Fourteenth Session of The IOC Assembly, Unesco, Paris, 1989.
- Redfield, A.C. *On the Proportions of Organic Derivatives in Sea Water and Their Relation to the Composition of Plankton.* James Johnstone Memorial Volume, Liverpool, University Press, pp. 348, 1934.
- Reed, J.B. *Tracking Technologies For Radioactive Waste Shipments.* Denver, Colorado: National Conference of State Legislatures, Series Title: State Legislative Report, V15, N4, 1990.
- Rees H., A. Eleftheriou. *North Sea Benthos - A Review Of Field Investigations Into The Biological Effects of Mans Activities.* Journal Du Conseil, V45 N3; pp. 284-305, 1989.
- Rees, H.L., M.D. Nicholson. *Trends in Lead Levels in the Horse-mussel From the Western North Sea.* Marine Pollution Bulletin, V20, N2, 1989.
- Refuse Disposal Systems.* Encyclopedia Britannica Macropedia 15th Edition, V13, Encyclopedia Britannica Inc., 1974.
- Reid, J.L. *Antarctic Oceanography I.* V53 Antarctic Series, American Geophysical Union, Washington D.C., 1971.
- Reijnders P., K. Lankester. *Status Of Marine Mammals In The North Sea.* Netherlands Journal of Sea Research, V26 N2-4, pp. 427-435, 1990.
- Reinhart, F.M. *Corrosion of Metals and Alloys in the Deep Ocean.* Technical Report R 834, Civil Engineering Laboratory, Naval Construction Battalion Center, Port Hueneme, California, February, 1976.
- Reinhart, F.M., J.F. Jenkins. *The Relationship Between the Concentration of Oxygen in Seawater and the Corrosion of Metals.* U.S. Naval Civil Engineering Laboratory Report, pp. 562-577, 1971.
- Reish D. *Pollution Of The Mediterranean Sea.* Water Environment Research, V65 N5, pp. 611-611, 1993.
- Report Of The ICES Advisory Committee On Marine Pollution, 1991.* Copenhagen, Denmark : International Council for the Exploration of the Sea, N177. 1991.
- Report Of The Mid-Term Expert Consultation On The Joint FAO (GFCM)/UNEP Coordinated Project On Pollution In The Mediterranean, Dubrovnik, 2-3 May 1977.* Expert Consultation on the Joint FAO (GFCM)/UNEP Coordinated Project on Pollution in the Mediterranean (1977 : Dubrovnik) Rome: Food and Agriculture Organization of the United Nations, 1977.
- Report to the Congress on Ocean Dumping Research, January Through December 1975.* Submitted in Compliance with Section 201, Title II of the Marine Protection, Research, and Sanctuaries Act of 1972, ON, U.S. Department of Commerce, 1976.

- Report to the Congress on Ocean Dumping Research, January Through December 1976.*  
Submitted in Compliance with Section 201, Title II of the Marine Protection, Research,  
and Sanctuaries Act of 1972, NOAA, U.S. Department of Commerce, 1977.
- Report/Joint FAO(GFCM)/UNEP Coordinated Project on Pollution in the Mediterranean.*  
Rome. Joint FAO(GFCM)/UNEP Coordinated Project on Pollution in the Mediterranean.  
General Fisheries Council for the Mediterranean, 1976.
- Review Of Contaminants In Baltic Sediments.* Cooperative Research Report, Copenhagen,  
Denmark: International Council for the Exploration of the Sea, 1992.
- Review of Several Plastic Dumping Incidents Involving the Navy, and What the Navy is Doing to  
End These Types of Incidents.* Hearing Before the Subcommittee on Oceanography,  
Great Lakes and the Outer Continental Shelf of the Committee on Merchant Marine and  
Fisheries House of Representatives One Hundred Second Congress, U.S. Government  
Printing Office, 1991.
- Rheinallt, T. *Impact Of Sewage Sludge. Pollution Of The North Sea. An Assessment.* Eds. W.  
Salomons, B.L. Bayne, E.K. Duursma, U. Foerstner, pp. 489-509, 1988.
- Richards, F.A. *Physical, Chemical and Productivity Data From A Survey of The Caribbean Sea  
and The Northeastern Tropical Pacific Ocean.* R. V. Thomas, G. Thompson Cruise 001,  
1970.
- Richards, W.J. *1936-Kinds and Abundances of Fish Larvae In The Caribbean Sea And Adjacent  
Areas.* Seattle, WA: U.S. Department of Commerce, National Oceanic and Atmospheric  
Administration, National Marine Fisheries Service, 1984.
- Ritchie, P., B. Calder. *The Pollution of The Mediterranean Sea.* Pacem In Maribus, Herbert  
Lang, 1972.
- Rivkin R. *Seasonal Patterns of Planktonic Production In Mcmurdo Sound, Antarctica.*  
American Zoologist, V31 N1, pp. 5-16, 1991.
- Roberts H., P. Wilson, A. Lugofernandez. *Biologic and Geologic Responses To Physical  
Processes - Examples Form Modern Reef Systems of The Caribbean-Atlantic Region.*  
Continental Shelf Research, V12 N7-8, 809-834, 1992.
- Robinson, M. *1906-Atlas of Monthly Mean Sea Surface and Subsurface Temperature and Depth  
of The Top of The Thermocline, Gulf of Mexico and Caribbean Sea.* San Diego, Scripps  
Institution of Oceanography, University of California, 1973.
- Robinson, M.K. *Atlas of Monthly Mean Sea Surface and Subsurface Temperature and Depth of  
The Top of The Thermocline Gulf of Mexico and Caribbean Sea.* 1973.
- Roethel F.J., V.T. Breslin. *Ash Disposal, Unique Method of Ash Disposal Can Benefit Marine  
Life.* Solid Waste and Power, 1988.

- Roethel, F.J., V.T. Breslin, K. Aldous. *Mobility of Dioxins and Furans Associated with Stabilized Incineration Residues in the Marine Environment*. Waste Management Institute, Sony Brook, New York.
- Roethel, F.J., V.T. Breslin. *Mobility of Dioxins, Furans, and Trace Metals From Stabilized MSW Combustor Ash in Seawater*. Waste Management Institute, Stony Brook, New York, 1993.
- Roethel, F.J., V.T. Breslin. *Stony Brook's MSW Combustor Ash Demonstration Programs*. Waste Management Institute, Stony Brook, New York.
- Romieu I., H. Weitzenfeld, J. Finkelman. *Urban Air Pollution In Latin America and The Caribbean*. Journal of The Air & Waste Management Association, V41 N9, pp. 1166-1171, 1991.
- Roots O. *Observations on The Transport and Wet Deposition of Airborne Pollutants Over The Baltic Sea and West-Estonian Islands*. *Ambio*, V21 N4 pp. 321-322, 1992.
- Sackett, W.M. *An Evaluation of the Effects of Man-Derived Wastes on the Viability of the Gulf of Mexico*.
- Saetre, R., M. Mork. *The Norwegian Coastal Current*. University Bergen, pp.795, 1981.
- Salemaa H., I. Vuorinen, P. Valipakka. *The Distribution And Abundance of Mysis Populations In The Baltic Sea*. *Annales Zoologici Fennici*, V27 N3 pp. 253-257, 1990.
- Saliba, L.J. *Regional Measures For Marine Pollution Control In The Mediterranean*. Viewpoint, Marine Pollution Bulletin V20(1), pp. 12-17, 1989.
- Salomons, W., B.L. Bayne, E.K. Duursma, U. Forstner. *Pollution of The North Sea: An Assessment*. Berlin, New York: Springer-Verlag, 1988.
- Sand, P.H. *Marine Environment Law in the United Nations Environment Programme, An Emergent Eco-Regime*. Tycooly Publishing, London and New York.
- Sandberg, M. *Baltic Sea Region Environmental Protection: "Eastern" Perspectives And International Cooperation*. Stockholm, Sweden: Almqvist & Wiksell International, 1992.
- Savchuk O., V. Volkova. *Study And Forecast of Baltic Sea Eutrophication With The Help of Simulation Model*. *Doklady Akademii Nauk Sssr*, V314 N4, pp. 1006-1009, 1990.
- Saydam, C., I. Salihoglu. *Dissolved/Dispersed Petroleum Hydrocarbons Suspended Sediment, Plastic Pelagic Tar and Other Litter in the North-Eastern Mediterranean*. Middle East Technical University Institute of Marine Sciences, P.K. 28, Erdemli-Ice 1, Turkey, VII Jounees Etud. Pollutions, Lucerne, 1984.
- Schiewer U. *Eutrophication of The Baltic Sea - Foreword*. *Internationale Revue Der Gesamten Hydrobiologie*, V76 N3 pp. 293-294, 1991.

- Schink, D.R. *Budget For Dissolved Silica In The Mediterranean Sea*. Sponsored by Office of Naval Research, 1967.
- Schlosser P., J. Bullister, R. Bayer. *Studies of Deep Water Formation and Circulation In The Weddell Sea Using Natural and Anthropogenic Tracers*. Marine Chemistry, V35 N1-4, pp. 97-122, 1991.
- Schluter M. *Organic Carbon Flux and Oxygen Penetration Into Sediments Of The Weddell Sea - Indicators For Regional Differences In Export Production*. Marine Chemistry, V35 N1-4, pp. 569-579, 1991.
- Schmidt D. *Mercury in Baltic and North Sea Waters*. *Water Air And Soil Pollution*, V62 N1-2 pp. 43-55, 1992.
- Schneider, H. *Results And Perspectives of The International Conference For The Protection of The North Sea*. *Abh. Naturw. Verein Bremen* 40(3):209-216, 1985.
- Schooley, A.H. *Wake Collapse in a Stratified Fluid: Experimental Exploration of Scaling Characteristics*. *Science* 160, 763, 1968.
- Schooley, A.H. *Wake Collapse in Stratified Fluid*. *Science* 157, 421, 1967.
- Schooley, A.H., R.W. Stewart. *Experiments with a Self-Propelled Body Submerged in a Fluid with a Vertical Density Gradient*. *J. Fluid Mech.* 15, Part I, 83, 1963.
- Schoomaker, J.S., R.R. Hammond, A.L. Heath, J.S. Cleveland. *A Numerical Model for Prediction of Sub-Littoral Optical Visibility*. *SPIE Ocean Optics XII*, pp. 18, 1994.
- Schroeder, W. W. *Physical Environment Atlas of The South Central Insular Shelf, Grand Bahama Island*. Alexander City, Alabama: B. and R. Russell, Educational and Charitable Foundation, Inc., Dauphin Island, Marine Science Programs, University of Alabama System, 1977.
- Schuchert, C., 1858-1942. *Historical Geology Of The Antillean-Caribbean Region, Or, The Lands Bordering The Gulf Of Mexico And The Caribbean Sea*. New York, Hafner Pub. Colorado, 1968.
- Scully-Power, P. *Navy Oceanographer Shuttle Observations STS41-G*. Naval Underwater Systems Center, NUSC Technology, Document 7611, 1986.
- Sebek, V. *The North Sea and The Concept Of Special Areas*. Eds. T. Ijlstra, D. Freestone, The North Sea: Perspectives On Regional Environmental Cooperation, V5, N1-3 pp. 157-166, 1990.
- Seibold, E., W.H. Berger. *The Sea Floor, An Introduction to Marine Geology*. Springer-Verlag Berlin Heidelberg, New York, 1982.
- Seiwell, H.R. *Application of The Distribution of Oxygen To The Physical Oceanography of The Caribbean Sea Region*. *Papers in Physical Oceanography and Meteorol.*, V6, N1, pp. 60, 1938.

- Shiber, J. G. *Plastic Pellets and Tar On Spain's Mediterranean Beaches*. Marine Pollution Bulletin, V18, pp. 84-86, 1987.
- Shillabeer, N. *Benthic Population Studies Of A North Sea Disposal Area Used For Industrial Liquid Waste*. Environmental Pollution, V69 N2-3, pp. 181-191, 1991
- Shimada, H., K. Miida, T. Yokooji. *The Relation Between the Corrosion Behavior of Steels and the Characteristics of Various Ocean Environments*. International Ocean Development Conference, V2, pp. 165-83, 1975.
- Ship's Guide, Shipboard Solid and Plastics Waste Management*. Office of the Chief of Naval Operations In Conjunction with: Commander, Pacific Fleet, Commander, Atlantic Fleet, Naval Sea Systems Command, Naval Supply Systems Command, David Taylor Research Center, 1991.
- Shomura, R.S., M.L. Godfrey. *Proceedings of the Second International Conference on Marine Debris 2-7 April 1989, Honolulu, Hawaii*. ON Technical Memorandum NMFS, 1990.
- Shumacher, M. *Seawater Corrosion Handbook*. Noyes Data Corporation, pp 383, Park Ridge, New Jersey, 1979.
- Side, J. *The North Sea Conference: Bremen 1984 London 1987*. Marine Pollution Bulletin V17(9), pp. 394-399. 1987.
- Siegel V., B. Bergstrom, J. Stromberg, P. Schalk. *Distribution, Size Frequencies and Maturity Stages of Krill, Euphausia-superba, In Relation To Sea-Ice In The Northern Weddell Sea*. Polar Biology, V10 N7, pp. 549-557, 1990.
- Simon, M., A.L. Alldredge, F. Azam. *Bacterial Carbon Dynamics on Marine Snow*. Marine Ecology Programe Series V65, pp. 205-211, 1990.
- Sivkov V., Y. Zhurov. *On The Origin of Accumulations of Suspended Matter in The Deep Basins of The Baltic Sea*. Okeanologiya, V31 N6, pp. 1060-1066, 1991.
- Skjaereth, J.B. *Towards The End Of Dumping In The North Sea*. Ed. W.E. Butler, VIII Anglo Soviet Symposium On The Law Of The Sea and International Shipping. Marine Polity CODEN V16, N2. pp. 130-140, 1992.
- Slaczka, W., E. Andrulewicz, A. Trzosinska. *First Periodic Assessment of the State of the Marine Environment of the Baltic Sea Area, 1980-1985*. Helsinki: Baltic Marine Environment Protection Commission, Helsinki Commission, 1986-1987.
- Slip D., H. Burton. *Accumulation Of Fishing Debris, Plastic Litter, and Other Artefacts, On Heard and Macquarie Islands In The Southern Ocean*. Environmental Conservation, V18 N3, pp. 249-254, 1991.
- Smed, J. *History of International North Sea Research ICES*. J. Sundermann & W. Lenz. North Sea Dynamics, Springer Verlag, Heidelberg, pp. 1-25, 1983.
- Smet M. *Case Studies Of Land-Based Pollution Policy 1 The North Sea*. Marine Policy, V16 N1, pp. 50-51, 1992.

- Smith O., J. Morrison. *Shipboard Acoustic Doppler Current Profiling in The Eastern Caribbean Sea, 1985-1986*. Journal of Geophysical Research-Oceans, V94 N7, pp. 9713-9719, 1989.
- Smith, D.C., M. Simon, A.L. Alldredge, F. Azam. *Intense Hydrolytic Enzyme Activity on Marine Aggregates and Implications For Rapid Particle Dissolution*. Nature, V359, pp. 139-142, 1992.
- Smith, R.W., M.C. Hyman. *Connective-Diffusive Bubble Transport in Ship Wakes*. Naval Coastal Systems Technical Note pp. 857-887, 1987.
- Smookler, A., C. Alig. *The Navy's Shipboard Waste Management Research & Development Program*. Naval Engineers Journal, 1992.
- Somer, E. *Heavy Metals in the Baltic*. ICES C.M., 1977.
- Spalding 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Spalding Treatment Works, United Kingdom, 1992-1994.
- Specter C., D. Gayle. *Managing Technology Transfer For Coastal Zone Development - Caribbean Experts Identify Major Issues*. International Journal of Remote Sensing, V11 N10, pp. 1729-1740, 1990.
- Squires, M.D. *Facilities For The Reception Of Wastes: A Shipping Industry View*. Workgroup Noordzee. Proceedings of the 2nd North Sea Seminar '86, Rotterdam, 1, 2, 3 October 1986, edited by E. Hey and G. Peet. V2. pp.195-211, Amsterdam, 1986.
- Stagg R. *North Sea Task Force Biological Effects Monitoring Programme*. Water Science and Technology, V24 N10, pp. 87-98, 1991.
- Stebbing, A.R.D., V. Dethlefsen, M. Carr. *Biological Effects o Contaminants In The North Sea: Results of The ICES/IOC Bremerhaven Workshop*. Amelinghausen: Inter-Research, 1992.
- Stefels J., H. Revier. *Protecting The North Sea Environment - Trends and Issues*. Water Science and Technology, V24 N10, pp. 277-281, 1991.
- Stephan K. ed. *Biogeochemistry and Distribution Of Suspended Matter In The North Sea and Implications To Fisheries Biology*. Hamburg: Geologisch-Palaontologischen Institutes der Universitat Hamburg, 1988.
- Stewart, M. *Numerical Prediction of Thermal Ship Wakes*. Naval Research Laboratory Memorandum Report 4955, September, 1987.
- Storer, R. *Standard Guide E 12 18. 1992 Annual Book of ASTM Standards, American Society for testing & Materials*. ASTM, Philadelphia, pp. 874-883, 1992.
- Studies of The Pollution Of The Baltic Sea*. ICES/SCOR Working Group on the Study of Pollution of the Baltic. Charlottenlund, Denmark: International Council for the Exploration of the Sea, 1977.

- Study of The Risk For Accidents and The Related Environmental Hazards From The Transportation of Chemicals By Tankers in The Baltic Sea Area: Report on A Study of The Transportation Pattern For Chemicals in The Baltic.* Helsinki: Baltic Marine Environment Protection Commission, Helsinki Commission, 1990.
- Sturges, W. *Numerical Modeling Studies of The Gulf Of Mexico and The Caribbean Sea Using The Bryan-Cox Model.* Florida State University, Department of Oceanography, The New Orleans, U.S. Department of The Interior, Minerals Management Service, Gulf of Mexico OCS Region, 1991.
- Sullivan, S.M. *A Review of the Biological Information Relating to OTEC Operation.* Oceanic Engineering Operations Interstate Electronics Corporation, Anaheim, CA.
- Summer, A.B., A. Smookler, C. Alig. *The Navy's Shipboard Waste Management R&D Program.* Session 2C, Naval Engineers Journal, 1992.
- Sundermann, J. *Suspended Particulate Matter In The North Sea - Field Observations and Model Simulations.* Philosophical Transactions of The Royal Society of London Series A-Physical Sciences and Engineering, V343 N1669:423-430, 1993.
- Sundermann, J., W. Lenz, eds. *North Sea Dynamics.* Berlin, New York: Springer-Verlag, 1983.
- Surface Sediment Distribution, Mediterranean Sea.* Prepared by Scripps Institution of Oceanography for U.S. Fleet Numerical Weather Central [S.I.: s.n.], 1970 (Presidio of Monterey, California: Defense Language Institute, Systems Development Agency), 1970.
- Survey of The Status of Oil Pollution Control In The South American Sub-Region of The Wider Caribbean Region.* Prepared In Co-Operation With IMO, Nairobi, Kenya: Oceans and Coastal Areas Programme Activity Centre, United Nations Environment Programme, 1990.
- Sverdrup, H.U. *On Vertical Circulation In The Ocean Due To The Action Of The Wind With Application To Conditions Within The Antarctic Circumpolar Current.* Cambridge, University Press, 1983.
- Sverdrup, H.U., M.W. Johnson, R.H. Fleming. *The Oceans.* Prentice Hall, New Jersey, 1942.
- Sverdrup, H.U., M.W. Johnson, R.H. Fleming. *The Oceans, Their Physics, Chemistry, and General Biology.* Prentice-Hall, Inc., Englewood Cliffs, N.J.
- Swanson, R.L. *The Incongruity of Policies Regulating New York City's Sewage Sludge: Lessons for Coastal Management.* Coastal Management, V21, pp. 299-312, 1993.
- Swanson, R.L., A. Valle-Levinson, C.J. Decker. *Recreation vs. Waste Disposal: The Use and Management of Jamaica Bay.* Long Island Historical Journal, V5, N1, pp. 21-41, Fall, 1992.
- Swanson, R.L., A. Valle-Levinson. *Meteorological Conditions That Kept Long Island and New Jersey Beaches Free of Floatables During the Summer of 1989.* State University of New York at Stony Brook, Baywood Publishing Co., Inc., 1990.

Swanson, R.L., R.L. Zimmer. *Meteorological Conditions Leading to the 1987 and 1988 Washups of Floatable Wastes on New York and New Jersey Beaches and Comparison of these Conditions with the Historical Record*. Estuarine, Coastal and Shelf Science, 30, pp. 59-78, 1990.

Swanson, R.L., R.R. Young, S.S. Ross. *An Analysis of Proposed Shipborne Waste Handling Practices Aboard United States Navy Vessels*. Marine Sciences Research Center, University at Stony Brook, Stony Brook, New York, 1994.

Swanson, R.L., T.M. Bell, J. Kahn, J. Olha. *Use Impairments and Ecosystem Impacts of the New York Bight*. Chemistry and Ecology, V5, pp. 99-127, 1991.

*Symposium on Investigations and Resources of The Caribbean Sea and Adjacent Regions, 1968: Willemstad, Curacao Papers on Fishery Resources*. Contribuciones Sobre Recursos Pesqueros. Edited By Marine Biology and Environment Branch, Fishery Resources Division, FAO. Rome, Food and Agriculture Organization of The United Nations, 1971.

*Symposium on Investigations and Resources of The Caribbean Sea and Adjacent Regions, Papers on Physical and Chemical Oceanography*. Willemstad, Curacao, 1968. Papers On Physical and Chemical Oceanography, Marine Geology and Geophysics and Marine Biology. Paris, Unesco, 1971.

*Symposium On Progress In Marine Research In The Caribbean And Adjacent Regions (1976 : Caracas, Venezuela) Symposium on Progress In Marine Research In The Caribbean and Adjacent Regions*. Caracas, Venezuela, 12-16 July 1976 : Papers On Oceanography, Meteorology, Geology and Geo-Physics = Symposia Sobre Adelantos En. Rome: Food and Agriculture Organization of The United Nations, 1978.

Szekielda, K.H., D. McGinnis, B. Ambroziak, P. McClain, D. Clark. *Satellite Observations Over The North Sea*. Eds. S. Kempe, G. Liebezeit, V. Dethlefsen, U. Harms, Scientific Common Problems of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry And Distribution of Suspended Matter In The North Sea and Implications-To-Fisheries-Biology. 1988. N65 pp. 1-33, 1988.

Tay, K.L. *The Role and Application of Environmental Bioassay Techniques in Support of the Impact Assessment and Decision-Making Under the Ocean Dumping Control Act in Canada*. Marine Environmental Branch, Environmental Protection, Conservation and Protection, Environment Canada, 1988.

*Ten Years After the Signing of the Helsinki Convention: National Statements by the Contracting Parties on the Achievements in Implementing the Goals of the Convention on the Protection of the Marine Environment*. Helsinki, Baltic Marine Environment Protection Commission, Helsinki Commission, 1984.

Tennekes, H., Lumley, J.L. *A First Course in Turbulence*. MIT Press, Cambridge, Massachusetts., pp. 300, 1972.

- Texas General Land Office. *The Gulf of Mexico as a Special Area Under MARPOL Annex V. Report Compiling Information in Support of Designation.* For Consideration by United States Delegation to the International Maritime Organization, 1987.
- Thorade, H. *Die Geschwindigkeit van Trifistromungen and die Ekman'sche Theorie.* Ann. d. Hydr. u. Marit. Meteorol., V42, pp. 379, 1914.
- Tijuana Monthly Sewage Discharge Report.* Comision de Servicios De Agua Del Estado Sistema De Alejamiento Y Tratamiento De Aguas Residuales De La Ciudad De Tijuana, BC, Mexico, 1994.
- Tilzer M., W. Gieskes, R. Heusel, N. Fenton. *The Impact Of Phytoplankton On Spectral Water Transparency In The Southern Ocean - Implications For Primary Productivity.* Polar Biology, V14 N2, pp. 127-136, 1994.
- Time That Solid Waste Remains In The Sea.* Sherkin Comment, N6, pp. 21, Medwaves, 19 January, 1990.
- Title I of the Ocean Dumping Ban Act of 1988.* Hearings Before the Subcommittee on Oversight and Investigations, Subcommittee on Fisheries and Wildlife Conservation and the Environment and the Subcommittee on Oceanography and the Great Lakes of the Committee on Merchant Marine and Fisheries House of Representatives, One Hundred First Congress First Session on Implementation of Title I of Public Law 100-66, The Ocean Dumping Ban Act of 1988, 1989.
- Tomascik T. *Settlement Patterns of Caribbean Scleractinian Corals On Artificial Substrata Along A Eutrophication Gradient, Barbados, West-Indies.* Marine Ecology-Progress Series, V77 N2-3, pp. 261-269, 1991.
- Topping, G., J.M. Bowers, P.G.W. Jones. *A Review of The Past And Present Measurements of Selected Trace Metals In Sea Water In The Oslo Commission And ICNAF/NAFO Areas.* Copenhagen: International Council for the Exploration of the Sea, 1980.
- Toxicity and Bioaccumulation of Selected Substances in Marine Organisms.* FAO/UNEP Meeting on Toxicity and Bioaccumulation of Selected Substances in Marine Organisms (1984: Rovinj, Yugoslavia) Report of the FAO/UNEP Meeting on Rovinj, Yougoslavia, 5-9 November 1984. Rome: Food and Agriculture Organization of the United Nations, 1985.
- Traverso, E. *Effects of Rolling Direction and Tensile Stress on the Corrosion of Naval Carbon Steel.* Natural Marine Environments-I: Natural Seawater, British Corrosion Journal, V14 (2), pp. 97-102, 1979.
- Treatment and Discharge of Industrial Wastewater In The Mediterranean Area: Report On A WHO Workshop, Venice, 10-14 June 1985.* Copenhagen: World Health Organization, Regional Office for Europe, 1986.
- Treguer P., B. Queguiner. *Biochemistry and Circulation of Water Masses In The Southern Ocean.* Based on Papers Presented At The International Symposium, Brest, France, 2-6 July 1990-Preface. Marine Chemistry, V35 N1-4:R11-R12, 1991.

- Treguer P., Jacques G. *Dynamics Of Nutrients and Phytoplankton, and Fluxes Of Carbon, Nitrogen and Silicon In The Antarctic Ocean*. Polar Biology, V12 N2, pp. 149-162, 1992.
- Tromp D, R. Coenen. *The 3rd International Conference On The Protection Of The North Sea - Review and Challenges*. Water Science and Technology, V24 N10:R15-R20, 1991.
- Turkstra E, M. Scholten, C. Bowmer, H. Schobben. *A Comparison of The Ecological Risks From Fisheries and Pollution To The North Sea Biota*. Water Science And Technology, V24 N10:147, 1991.
- U.S. Coast Guard. *Adoption of Amendments to the Annex of the Protocol of 1978 Relating to the International Convention for the Prevention of Pollution From Ships, 1973 (Designation of the Gulf of Aden as a Special Area)*. MEPC25/20, 1994.
- U.S. Coast Guard. *IMO Special Areas, Final Rule*. Draft, Antarctic Designation, 1994.
- U.S. Environmental Protection Agency (EPA). *Designation of the Gulf of Mexico*. The Oceanic Society, 1990.
- U.S. Environmental Protection Agency (EPA). *Evaluation of Dredged Material Proposed for Ocean Disposal - Testing Manual*. EPA Washington D.C. and Department of the Army, Washington D.C., 1991.
- U.S. Environmental Protection Agency (EPA). *Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Freshwater Organisms*. First Edition, EPA/600/4-89/001, 1988.
- U.S. Environmental Protection Agency (EPA). *Technical Support Document for Water Quality-Based Toxics Control*. Office of Water, EPA 505/2-90-001, March, 1991.
- U.S. Navy Climatic Study Of The Caribbean Sea And Gulf Of Mexico*. United States, Naval Oceanography Command Detachment, Asheville, N.C. V1: West Caribbean Sea and Central American Waters, 1985.
- U.S. Navy Climatic Study Of The Caribbean Sea and Gulf Of Mexico*. United States. Naval Oceanography Command Detachment, Asheville, N.C. V4: Gulf Of Mexico and Gulf Of Tehuantepec, 1986.
- Uchupi, E. *Bathymetric Atlas of The Atlantic, Caribbean, and Gulf Of Mexico*. Compiled, Woods Hole, Massachusetts, Woods Hole Oceanographic Institution, 1971.
- Underwood, P.C. *New Law of The Sea For The Caribbean - An Examination of Marine Law and Policy Issues In The Lesser Antilles* - Gold, E. Marine Policy, V13 N4, pp. 353-355, 1989.
- UNEP. *Assessment of Organotin Compounds As Marine Pollutants In The Mediterranean = Evaluation Des Composés Organostanniques En Tant Que Polluants Du Milieu Marin En Méditerranée*. Athens: MAP Technical Reports Series N33, UNEP, 1989.

- UNEP. *Assessment of The State Of Pollution Of The Mediterranean Sea By Cadmium and Cadmium Compounds = Evaluation De L'etat De La Pollution De La Mer Mediterranee Par Le Cadmium Et Les Composes De Cadmium*. MAP Technical Reports Series N34 Athens: UNEP, 1989.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Mercury and Mercury Compounds = Evaluation De L'etat De La Pollution De La Mer Mediterranee Par Le Mercure Et Les Composes Mercuriels*. Athens: UNEP, 1987.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Organohalogen Compounds = Evaluation De L'etat De La Pollution De La Mer Mediterranee Par Les Composes Organohalogenes*. MAP Technical Reports Series N39 UNEP, 1990.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Organophosphorus Compounds = Evaluation De L'etat De La Pollution De La Mer Mediterranee Par Les Composes Organophosphores*. MAP Technical Reports Series N58, UNEP, 1988.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Persistent Synthetic Materials Which May Float, Sink or Remain In Suspension*. MAP Technical Reports Series N56, Mediterranean Action Plan, MEDPOL N56, pp. 111, 1991.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Petroleum Hydrocarbons = Evaluation De L'etat De La Pollution De La Mer Mediterranee Par Les Hydrocarbures De Petrole*. Athens: UNEP, 1988.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Radioactive Substances = Evaluation De L'etat De La Pollution De La Mer Mediterranee Par Les Substances Radioactives*. Athens: UNEP, 1992.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Persistent Synthetic Materials Which May Float, Sink Or Remain In Suspension*. Map Technical Report Series, Athens-Greece, UNEP, N56, pp. 111, 1991.
- UNEP. *Assessment of The State of Pollution of The Mediterranean Sea By Radioactive Substances*. Athens: UNEP, 1992.
- UNEP. *Regional Seas Programme In Latin America and Wider Caribbean/United Nations Environment*, UNEP Regional Seas Reports and Studies N22 Rev. 22, 1985.
- UNEP. *State Of The Mediterranean Marine Environment*. United Nations Environment Programme, Mediterranean Action Plan, Athens: UNEP, 1989.
- UNEP. *Technical Annexes to the Report on the State of the Marine Environment*. UNEP Regional Seas Reports and Studies N114/2, 1990.
- UNESCO. *Coastal Ecosystems of Latin America and the Caribbean*. Objectives, Priorities and Activities of Unesco's COMAR Project for the Latin American and Caribbean Region, Caracas, Venezuela, 1983.

- United States. Naval Oceanography Command Detachment, Asheville, N.C. *U.S. Navy Climatic Study Of The Caribbean Sea And Gulf Of Mexico*, V3: Florida Coastal Waters and Southwest Atlantic, 1986.
- Van Pagee, J. A.. *Transport and Water Quality Modeling In The Southern North Sea In Relation To Coastal Pollution Research and Control*. Delft, Netherlands: Waterloopkundig Laboratorium, 1986.
- van Pagee, J.A. *Mathematical Modelling As A Tool For Assessment Of Coastal Water Pollution: Case Study*, North Sea Delft, Netherlands: Delft Hydraulics, 1989.
- van Pagee, J.A. *North Sea Pollution: The Use of Modeling Techniques for Impact Assessment of Waste Inputs*. Delft Hydraulics, Delft, Netherlands, 1989.
- Vanderloeff M., A. Vanbennekom. *Weddell Sea Contributes Little To Silicate Enrichment In Antarctic Bottom Water*. Deep-Sea Research Part A-Oceanographic Research Papers, V36 N9. pp. 1341-1357, 1989.
- Vanderloeff M., G. Berger. *Scavenging and Particle Flux-Seasonal and Regional Variations In The Southern Ocean (Atlantic Sector)*. Marine Chemistry, V35 N1-4, pp. 553-567, 1991.
- Vauk, G., G. Dahlmann, E. Hartwig, J. Ranger, B. Reineking, E. Schrey, E. Vauk, E. Hentzelt. *Recording of Oil Victims on The German North Sea Coast And Results of Oil Analyses As Well As Investigations of Burdening of The German Bight By Ship's Refuse*. Umweltbundesamt, Berlin FRG. TEXTE, N30, pp. 153, 1987.
- Vauk, G.J.M., E. Schrey. *Litter Pollution from Ships in the German Bight*. Marine Pollution Bulletin, V18, N6B, pp. 316-319, 1987.
- Venizelos, L.E. *The Endangered Loggerheads Of Zakynthos (Greece): A Part Of The Mediterranean Sea Turtle Conservation Issue*. The Formation of MEDASSET (Mediterranean Association to Save the Sea Turtles). Thalassoraphica. V13, N1, pp. 53-59, 1990.
- Verlencar X., V. Dhargalkar. *Primary Productivity and Nutrients In The Indian Sector of The Southern Ocean*. Indian Journal of Marine Sciences, V21 N1, pp. 6-12, 1992.
- Vernette G., A. Mauffret, C. Bobier, L. Briceno. *Mud Diapirism, Fan Sedimentation and Strike-Slip Faulting, Caribbean Colombian Margin*. Tectonophysics, V202 N2-4, pp. 335-349, 1992.
- Vesecky, J.F., R.H. Stewart. *The Observation of Ocean Surface Phenomena Using Imagery From the SEASAT Synthetic Aperture Radar: An Assessment*. Journal Geophysics, Research, V87(C5), pp. 3397-3430, 1982.
- Veth C. *The Structure and Evolution of The Top Layers of The Water Column Across The Marginal Ice Zone During Spring 1988 In The Scotia-Weddell Sea Sector of The Southern Ocean*. Marine Chemistry, V35 N1-4, pp. 63-76, 1991.

- Veth C., C. Lancelot, S. Ober. *On Processes Determining The Vertical Stability Of Surface Waters In The Marginal Ice Zone of The North-Western Weddell Sea and Their Relationship With Phytoplankton Bloom Development*. Polar Biology, V12 N2, pp. 237-243, 1992.
- Viale, D., N. Verneau, Y. Tison. *Stomach Obstruction In A Spermwhale Beached On The Lavezzi Islands: Macropollution In The Mediterranean*. Journal de Recherche Oceanographique, Union des Oceanographes de France, 16 [3/4] , 100-102. In French, English abstract, 1992.
- Voipio, A. *The Baltic Sea*. Elsevier Scientific Publication Co., New York, New York, 1981.
- von Wachenfeldt, T., L. Edler, eds. *Symposium of The Baltic Marine Biologists (8th: 1983 Lund, Sweden) Proceedings of The 8th Symposium Of The Baltic Marine Biologists Recent Changes in The Baltic Ecosystem*. Denmark Marine Biological Laboratory, 1984.
- Waite T., R. Szymczak. *Particulate Iron Formation Dynamics In Surface Waters of The Eastern Caribbean*. Journal of Geophysical Research. Oceans, V98 N2, pp. 2371-2383, 1993.
- Walney 1990-1995 Sewage Discharge Monitoring Report*. National Rivers Authority, Walney Channel Sewage Treatment Works U.K., 1990-1995.
- Walton, D.W.H. *Antarctic Science*. Cambridge University Press, Great Britain, 1987.
- Wangersky, P.J., S.R. Aston, J.T. Baker, B.B. Benson, M. Branica, J.N. Butler, R. Chester, R.A. Duce, E.K. Duursma, D. Dyrssen, M. Ehrhardt, J.M. Gieskes, N. Handa, A. Jensen, I.R. Kaplan, R. Kilho Park, Y. Kitano, K. Kremling, A.P. Lisitzin, P.S. Liss, C.S. Martens, A. Nissenbaum, H.G. Ostlund, H. Postma, A. Walton, R.F. Weiss, M. Whitfield, J. Willey, T.R.S. Wilson, R.G. Zika. *Marine Chemistry, An International Journal for Studies of all Chemical Aspects of the Marine Environment*. Elsevier Science Publishers, B.V., Amsterdam, V19, 1986.
- Warrington 1990-1995 Sewage Discharge Monitoring Report*. National Rivers Authority, Warrington South, Final Effluent Tank Outlet Bestel Dean Sampler, U.K., 1990-1995.
- Waste Discharge Into The Marine Environment: Principles and Guidelines For The Mediterranean Action Plan*. Prepared in Collaboration with the Institute of Sanitary Engineering, Polytechnic of Milan, Italy, 1st ed. Oxford [Oxfordshire], New York: Pergamon Press, 1982.
- Waste Disposal - Establish NOPC*. Hearings Before the Subcommittee on Oceanography and the Committee on Merchant Marine and Fisheries House of Representatives Ninety-Eighth Congress First Session on Hazardous Waste Disposal - H.R. 1700, Establishing a National Oceans Policy Commission - H.R. 2853, Serial No. 98-15, 1983.
- Wastes in Marine Environments, Summary*. Congress of the United States, Office of Technology Assessment, Washington D.C., U.S., 1987.
- Wastewater Chemistry Laboratory. *Discharge Specifications for the Point Loma Ocean Outfall. City of San Diego*. Metropolitan Wastewater Department, 1994.

- Watermann, B, H. Kranz. *Pollution and Fish Diseases in The North Sea - Some Historical Aspects*. Marine Pollution Bulletin, V24 N3, pp. 131-138, 1992.
- Weast, R. *CRC Handbook of Chemistry and Physics, 52nd Edition*. The Chemical Rubber Company, July, 1971.
- Weber, C. *Short-Term Methods for Estimating the Acute Toxicity of Effluents & Receiving Waters to Marine & Estuarine Organisms*. Environmental Monitoring & Support Laboratory, Cincinnati, 1988.
- Wefer G., G. Fischer. *Annual Primary Production and Export Flux In The Southern Ocean From Sediment Trap Data*. Marine Chemistry, V35 N1-4, pp. 597-613, 1991.
- Wenk, V.D. *Water Pollution: Domestic Wastes*. A Technology Assessment Methodology, National Technical Information Service, V6, 1971.
- West, J.M. *Basic Corrosion and Oxidation*. John Wiley and Sons, pp. 122, New York, 1980.
- Westerlund S., P. Ohman. *Cadmium, Copper, Cobalt, Nickel, Lead, and Zinc In The Water Column Of The Weddell Sea, Antarctica*. Geochimica Et Cosmochimica Acta, V55 N8, pp. 2127-2146,. 1991.
- Westerlund S., P. Ohman. *Iron In The Water Column Of The Weddell Sea*. Marine Chemistry, V35 N1-4, pp. 199-217, 1991.
- Westing, A.H. *Comprehensive security for the Baltic: An Environmental Approach*. London, SAGE, Newbury Park, California, 1989.
- White M., U. Piatkowski. *Abundance, Horizontal and Vertical Distribution Of Fish In Eastern Weddell Sea Micronekton*. Polar Biology, V13 N1, pp. 41-53, 1993.
- Widnes 1990-1995 Sewage Monitoring Report*. National Rivers Authority, Widnes Final Effluent, Hale Bank, U.K., 1995.
- Wilburn, A.M. *Current Velocity and Hydrographic Observations In The Straits Of Florida, The Caribbean Sea And Offshore Of The Antillean Archipelago: Subtropical Atlantic Climate Study (STACS) 1984 and 1985*, Miami, Florida: National Oceanic and Atmospheric Administration, Environmental Research Laboratories, Atlantic Oceanographic and Meteorological Laboratory, Springfield, VA, 1987.
- Wilde, P. *Environmental Monitoring and Assessment Program at Potential OTEC Sites*. Lawrence Berkeley Laboratory, University of California Berkeley, California, U.S.
- Wilde, P. *Status of OTEC Environmental Monitoring Program*. Lawrence Berkeley Laboratory, Berkeley, California, U.S.
- Wilkinson C., A. Cheshire. *Comparisons of Sponge Populations Across The Barrier Reefs Of Australian And Belize - Evidence For Higher Productivity In The Caribbean*. Marine Ecology-Progress Series, V67 N3, pp. 285-294, 1990.
- Will The Mediterranean Be Saved?* Outlook, SHIP: "series". N66, pp. 9-10, 1989.

- Williams S. *Experimental Studies of Caribbean Seagrass Bed Development*. Ecological Monographs, V60 N4, pp. 449-469, 1990.
- Williams, P.M. *Chemical and Microbiological Studies of Sea-Surface Films in the Southern Gulf of California and Off the West Coast of Baja California*. Marine Chemistry, V19, pp. 17-98, 1986.
- Williams, R., J. Aiken. *Optical Measurements From Underwater Towed Vehicles Deployed From Ships Of Opportunity In The North Sea*. Belhaven, Washington: International Society for Optical Engineering. pp. 198, SPIE Proceedings V1269, 1990.
- Wirth, H., M.G. Wiesner. *Sedimentary Facies In The North Sea*. Scientific Common Problems of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry and Distribution of Suspended Matter In The North Sea And Implications-To-Fisheries-Biology, N65 pp. 269-287, 1988.
- Wirth, H., R. Seifert. *Mineralogy, Geochemistry And SEM Observations Of Suspended Particulate Matter From The North Sea, Spring 1984*. Eds. S. Kempe, G. Liebezeit, V. Dethlefsen, U. Harms, Scientific Common Problems Of The Environment United Nations Environment Programme, Hamburg (FRG) International Carbon Unit, Biogeochemistry And Distribution Of Suspended Matter In The North Sea And Implications-To-Fisheries-Biology, N65 pp. 163-181, 1988.
- Wisbech 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Wisbech Treatment Works, U.K., 1992-1994.
- Witham 1992-1994 Sewage Discharge Monitoring Report*. National Rivers Authority, Witham Treatment Works, U.K., 1992-1994.
- Wolanski, E., R. Gibbs, P. Ridd, A. Mehta. *Settling of Ocean-dumped Dredged Material, Townsville, Australia*. Estuarine, Coastal and Shelf Science, 35, pp. 473-489, 1992.
- Wolfe, D. *Persistent Plastics and Debris in the Ocean: an International Problem of Ocean Disposal*. Marine Pollution Bulletin, V18, N6B, pp. 303-305, 1987.
- Wolfe, D.A. ed. *Plastics in the Sea, Selected Papers from the Sixth International Ocean Disposal Symposium*. Marine Pollution Bulletin, Pergamon Press, Oxford, New York, V18, V6B, 1986.
- Worf, D.L. *Biological Monitoring for Environmental Effects*. D.C. Heath and Company, U.S., 1980.
- Workshop On Pollution Of The Mediterranean (5th: 1980: Cagliari, Italy) Ves Journees D'etudes Sur Les Pollutions Marines En Mediterranee: Cagliari, 9-13 Octobre 1980/ Commission Internationale Pour L'exploration Scientifique De La Mer Mediterranee, Monaco, Et Programme Des Nations Monaco: Secretariat General De La Commission, 1981.*
- Workshop On Pollution Of The Mediterranean (7th : 1984 : Lucerne, Switzerland) Viies Journees D'etudes Sur Les Pollutions Marines En Mediterranee: Lucerne, 11-13*

- Octobre 1984. Commission Internationale Pour L'exploration Scientifique de la mer Mediterranee, Monaco, Programme, Monaco: Secretariat general de la C.I.E.S.M, 1985.
- World Bank. *The Environmental Program For The Mediterranean: Preserving A Shared Heritage and Managing A Common Resource*. World Bank, Luxembourg: European Investment Bank, Washington, D.C., USA, 1990.
- World Health Organization, Regional Office for Europe. *The Hazards To Health And Ecological Effects Of Pollution Of The North Sea: Report On A Working Group Convened By The Regional Office For Europe Of The World Health Organization : Bilthoven, 6-8 December 1972*. Copenhagen: The Office, 1973.
- World Health Organization. *The Hazards to Health and Ecological Effects of Persistent Substances in the Environment, Sources, Turnover in the Environment, and Ecological Effects of Arsenic, Cadmium, Lead, Manganese and Mercury*. World Health Organization, Regional Office For Europe, Stockholm, 1973.
- Wulff F., A. Niemi. *Priorities For The Restoration of The Baltic Sea - A Scientific Perspective*. *Ambio*, V21 N2, pp. 193-195, 1992.
- Wulff F., L. Rahm, P. Jonsson, L. Brydsten. *A Mass-Balance Model of Chlorinated Organic Matter For The Baltic Sea - A Challenge For Ecotoxicology*. *Ambio*, V22 N1, pp. 27-31, 1993.
- Wulff F., L. Rahm. *Accumulation of Chlorinated Organic Matter In The Baltic Sea From 50 Years Of Use - A Threat To The Environment*. *Marine Pollution Bulletin*, V26 N5 pp. 272-275, 1993.
- Wust, G. *1890-Stratification and Circulation in The Antillean-Caribbean Basins*. New York, Columbia University Press, 1964.
- Wyatt, J.R. *Tar Ball Pollution in the Western Caribbean*. University of the West Indies, Mona, Kingston, 7, Jamaica, 1982.
- Xavier, P. ed. *The Mediterranean*. London: Collins & Brown, 1991.
- Yanchinski, S. *UN Strides Into Caribbean Pollution*. *New Scientist*, 1980.
- Yoneda, Y., Y.Yoshida. *The Relation Between the Sardine and the Food Plankton-I. On the Food Intake by Sardinops melanosticta*. *Bulletin of the Japanese Society of Scientific Fisheries*, V21, N2, 1955.
- Yurkovskis A., F. Wulff, L. Rahm, A. Andruzaitis. *A Nutrient Budget of The Gulf Of Riga, Baltic Sea*. *Estuarine Coastal and Shelf Science*, V37 N2, pp. 113-127, 1993.
- Zanke, U. *Seegang Erzeugte Kolke am Bauwerken, Sonderforschungsbereich 79, Teilsprojekt B9*. Pergamon Press, Inc., Maxwell House, Fairview Park, Elmsford, New York, 1977
- Zar, J.H. *Biostatistical Analysis*. Prentice Hall, Englewood Cliffs, 1984.

- Zika R., P. Milne, O. Zafiriou. *Photochemical Studies of The Eastern Caribbean - An Introductory Overview*. Journal Of Geophysical Research-Oceans, V98.
- Zmarzly, D.L., T.D. Stebbins, D. Pasko, R.M. Duggan, K.L. Barwick. *Spatial Patterns and Temporal Succession in Soft-Bottom Macroinvertebrate Assemblages Surrounding an Ocean Outfall on the Southern San Diego Shelf: Relation to Anthropogenic and Natural Events*. Marine Biology Laboratory, San Diego, California, 1993.
- Zmudzinski, L. *Environmental Quality in the Baltic Region*. University of Education, Slupsk, Poland, in *Comprehensive Security for the Baltic*, SAGE Publications, 1989.
- Zoffmann, C. *Microbial and Nutrient Pollution Along the Coasts of Alicante, Spain*. Marine Pollution Bulletin, V20, N2, 1989.
- Zutic, V., G. Cauwet, A. Monaco. *Role of Organic Aggregates in the Transport of Pollutants in Mediterranean Estuaries*. Center for Marine Research, "Rudjer Boskovic" Institute Zagreb, Yugoslavia, VII Jounees Etud. Pollutions, Lucerne, 1984.